

RESEARCH



Exact relations for Green's functions in linear PDE and boundary field equalities: a generalization of conservation laws

Graeme W. Milton^{1*}  and Daniel Onofrei²

*Correspondence:

milton@math.utah.edu

¹Department of Mathematics,
University of Utah, Salt Lake City,
UT, USA

Full list of author information is
available at the end of the article

Abstract

Many physical equations have the form $\mathbf{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{E}(\mathbf{x}) - \mathbf{h}(\mathbf{x})$ with source $\mathbf{h}(\mathbf{x})$ and fields \mathbf{E} and \mathbf{J} satisfying differential constraints, symbolized by $\mathbf{E} \in \mathcal{E}, \mathbf{J} \in \mathcal{J}$ where \mathcal{E}, \mathcal{J} are orthogonal spaces. We show that if $\mathbf{L}(\mathbf{x})$ takes values in certain nonlinear manifolds \mathcal{M} , and coercivity and boundedness conditions hold, then the infinite body Green's function (fundamental solution) satisfies exact identities. The theory also links Green's functions of different problems. The analysis is based on the theory of exact relations for composites, but without assumptions about the length scales of variations in $\mathbf{L}(\mathbf{x})$, and more general equations, such as for waves in lossy media, are allowed. For bodies Ω , inside which $\mathbf{L}(\mathbf{x}) \in \mathcal{M}$, the "Dirichlet-to-Neumann map" giving the response also satisfies exact relations. These boundary field equalities generalize the notion of conservation laws: the field inside Ω satisfies certain constraints that leave a wide choice in these fields, but which give identities satisfied by the boundary fields, and moreover provide constraints on the fields inside the body. A consequence is the following: if a matrix-valued field $\mathbf{Q}(\mathbf{x})$ with divergence-free columns takes values within Ω in a set \mathcal{B} (independent of \mathbf{x}) that lies on a nonlinear manifold, we find conditions on the manifold, and on \mathcal{B} , that with appropriate conditions on the boundary fluxes $\mathbf{q}(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}(\mathbf{x})$ (where $\mathbf{n}(\mathbf{x})$ is the outward normal to $\partial\Omega$) force $\mathbf{Q}(\mathbf{x})$ within Ω to take values in a subspace \mathcal{D} . This forces $\mathbf{q}(\mathbf{x})$ to take values in $\mathbf{n}(\mathbf{x}) \cdot \mathcal{D}$. We find there are additional divergence-free fields inside Ω that in turn generate additional boundary field equalities. Consequently, there exist partial null Lagrangians, functionals $F(\mathbf{w}, \nabla\mathbf{w})$ of a vector potential \mathbf{w} and its gradient that act as null Lagrangians when $\nabla\mathbf{w}$ is constrained for $\mathbf{x} \in \Omega$ to take values in certain sets \mathcal{A} , of appropriate nonlinear manifolds, and when \mathbf{w} satisfies appropriate boundary conditions. The extension to certain nonlinear minimization problems is also sketched.

Keywords: Green's functions, Exact relations, Inverse problems, Boundary field equalities, Inhomogeneous media

Mathematics Subject Classification: 35J08, 35J25, 65N21

1 Introduction

Many important linear equations of physics in an inhomogeneous medium of infinite extent in \mathbb{R}^d can be written as a system of second-order linear partial differential equations:

$$\sum_{i=1}^d \frac{\partial}{\partial x_i} \left(\sum_{j=1}^d \sum_{\beta=1}^m L_{i\alpha j\beta}(\mathbf{x}) \frac{\partial u_{\beta}(\mathbf{x})}{\partial x_j} \right) = f_{\alpha}(\mathbf{x}), \quad \alpha = 1, 2, \dots, m, \quad (1.1)$$

for the m -component potential $\mathbf{u}(\mathbf{x})$ given the m -component source term $\mathbf{f}(\mathbf{x})$. If the integral of $\mathbf{f}(\mathbf{x})$ over \mathbb{R}^d is zero, these can be reexpressed as

$$J_{i\alpha}(\mathbf{x}) = \sum_{j=1}^d \sum_{\beta=1}^m L_{i\alpha j\beta}(\mathbf{x}) E_{j\beta}(\mathbf{x}) - h_{i\alpha}(\mathbf{x}), \quad E_{j\beta}(\mathbf{x}) = \frac{\partial u_{\beta}(\mathbf{x})}{\partial x_j}, \quad \sum_{i=1}^d \frac{\partial J_{i\alpha}(\mathbf{x})}{\partial x_i} = 0, \quad (1.2)$$

where, counter to the usual convention, we find it convenient to let the divergence act on the first index of \mathbf{J} , and to let the gradient in $\mathbf{E} = \nabla \mathbf{u}$ be associated with the first index of \mathbf{E} , and $\mathbf{h}(\mathbf{x})$ is chosen so

$$\sum_{i=1}^d \frac{\partial h_{i\alpha}(\mathbf{x})}{\partial x_i} = f_{\alpha}(\mathbf{x}). \quad (1.3)$$

Assuming we are looking for solutions where $\mathbf{E}(\mathbf{x})$ and $\mathbf{J}(\mathbf{x})$ are square integrable in \mathbb{R}^d , integration by parts shows that

$$\int_{\mathbb{R}^d} \sum_{i=1}^d \sum_{\alpha=1}^m J_{i\alpha}(\mathbf{x}) E_{i\alpha}(\mathbf{x}) \, d\mathbf{x} = 0. \quad (1.4)$$

Thus, $\mathbf{E}(\mathbf{x})$ and $\mathbf{J}(\mathbf{x})$ belong to orthogonal spaces: \mathcal{E} the set of square-integrable fields $\mathbf{E}(\mathbf{x})$ such that $\mathbf{E} = \nabla \mathbf{u}$ for some m -component potential \mathbf{u} , and \mathcal{J} the set of square-integrable fields $\mathbf{J}(\mathbf{x})$ such that $\nabla \cdot \mathbf{J} = 0$. With these definitions, Eq. (1.2) take the equivalent, more abstract, form

$$\mathbf{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{E}(\mathbf{x}) - \mathbf{h}(\mathbf{x}), \quad \mathbf{J} \in \mathcal{J}, \quad \mathbf{E} \in \mathcal{E}, \quad \mathbf{h} \in \mathcal{H}. \quad (1.5)$$

where $\mathcal{H} = \mathcal{E} \oplus \mathcal{J}$ consists of square-integrable $d \times m$ matrix-valued fields. When $m = 1$, we can interpret these equations as conductivity equations, with \mathbf{J} as the current field, $\mathbf{f} = \nabla \cdot \mathbf{h}$ as a source of current, $\mathbf{E} = -\nabla V$ as the electric field, and $V = -u$ as the electrical potential. Then the $\sigma_{ij}(\mathbf{x}) = L_{i1j1}(\mathbf{x})$ are the elements of the conductivity tensor field $\boldsymbol{\sigma}(\mathbf{x})$.

As shown in the Appendix (see also [26] Chap. 2, [35] Chap. 1, [27, 28] and the appendix of [29]), this structure (1.5) is suitable for a multitude of additional linear physical equations too, including wave equations. They can be formulated in a Hilbert space \mathcal{H} of square-integrable fields in \mathbb{R}^d taking values in some tensor space \mathcal{T} , where \mathcal{H} can be split into two orthogonal subspaces \mathcal{E} and \mathcal{J} , i.e., $\mathcal{H} = \mathcal{E} \oplus \mathcal{J}$. This splitting is typically such that the operator Γ_1 that projects onto \mathcal{E} is local in Fourier space, i.e., if $\mathbf{E} = \Gamma_1 \mathbf{A}$ then the Fourier components $\widehat{\mathbf{E}}(\mathbf{k})$ and $\widehat{\mathbf{A}}(\mathbf{k})$ of \mathbf{E} and \mathbf{A} are related via $\widehat{\mathbf{E}}(\mathbf{k}) = \Gamma_1(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k})$ for some operator $\Gamma_1(\mathbf{k})$ that projects onto a subspace $\mathcal{E}_{\mathbf{k}} \subset \mathcal{T}$. The Appendix gives examples of fields $\mathbf{E} \in \mathcal{E}$ having some components that beside those involving $\nabla \mathbf{u}$, just involve \mathbf{u} alone: examples are the acoustic, electrodynamic, or elastodynamic wave equations in possibly lossy (energy absorbing) media [32, 34, 35] at constant (possibly complex) frequency. The lossy nature of the moduli at constant frequency ensures the coercivity we need for well posedness. Also the fields $\mathbf{E} \in \mathcal{E}$ could have higher-order gradients: a classic example is the Kirchhoff plate equation (see, e.g., [26] Sect. 2.3, and references therein) where one takes $\mathbf{E}(\mathbf{x})$ to be the linearized plate curvature $\mathbf{E} = \nabla \nabla u$, in which $u(\mathbf{x})$ is the (infinitesimal)

vertical deflection of the plate. Furthermore, some components of the fields $\mathbf{E} \in \mathcal{E}$ are not necessarily derivatives of potentials but could also involve, say, divergence-free vector fields (and the corresponding components of the fields $\mathbf{J} \in \mathcal{J}$ would then be gradients of potentials). Such mixed formulations are useful in quasistatic and wave equations in lossy media when one wants to reformulate the problem so that $\mathbf{L}(\mathbf{x})$ is real and positive definite [7, 32, 34, 35]. Again, examples are given in the Appendix.

To begin we are considering an inhomogeneous medium of infinite extent in \mathbb{R}^d , where the material moduli are contained in a tensor $\mathbf{L}(\mathbf{x})$. Some types of boundedness and coercivity constraints on $\mathbf{L}(\mathbf{x})$ are usually needed to ensure that Eq. (1.5) always have a unique weak solution for $\mathbf{E}(\mathbf{x})$ [and hence $\mathbf{J}(\mathbf{x})$] for any given source field $\mathbf{h}(\mathbf{x})$ with say compact support. Later we will consider the equations in a bounded domain Ω . Then the concept of weak solution implies that, further throughout the paper, unless more regularity is specified, the tangential component of the $\mathbf{E}(\mathbf{x})$ along the boundary can be viewed in the $H^{\frac{1}{2}}$ sense and the normal component of $\mathbf{J}(\mathbf{x})$ over the boundary in the $H^{-\frac{1}{2}}$ sense.

In an infinite medium in the governing Eq. (1.5) the uniquely determined field \mathbf{E} depends linearly on the source term \mathbf{h} . So assuming \mathbf{E} depends continuously on \mathbf{h} , as it should in any physical problem of interest, the Schwartz kernel theorem implies that we can informally write

$$\mathbf{E}(\mathbf{x}) = \int_{\mathbb{R}^d} \mathbf{G}(\mathbf{x}, \mathbf{x}') \mathbf{h}(\mathbf{x}') \, d\mathbf{x}', \tag{1.6}$$

where the integral kernel $\mathbf{G}(\mathbf{x}, \mathbf{x}')$ (possibly a generalized function) is the Green’s function for the problem that depends on both \mathbf{x} and \mathbf{x}' , and not just on $\mathbf{x} - \mathbf{x}'$, because the medium is inhomogeneous.

Remark 1.1 In general the existence of a continuous Green Function for problem (1.5) may be difficult to prove. Thus, for the sake of clarity of exposition, further in the paper δ will denote a smooth approximation of the Dirac delta distribution and \mathbf{G} will denote an approximate Green function (still called the Green’s function), i.e., the solution of problem (1.5) with source δ (i.e., a smooth approximation of the Dirac delta distribution).

The Green’s function can also be considered as a linear map $\mathbf{G} : \mathcal{H} \rightarrow \mathcal{H}$, or more specifically, $\mathbf{G} : \mathcal{H} \rightarrow \mathcal{E}$. The main objective in this paper is to show that when $\mathbf{L}(\mathbf{x})$ is constrained to take values in certain nonlinear manifolds \mathcal{M} , then the Green’s function kernel $\mathbf{G}(\mathbf{x}, \mathbf{x}')$ satisfies some exact identities for every $\mathbf{x} \neq \mathbf{x}'$. The manifold \mathcal{M} , with dimension k_0 , need not have codimension one.

When we say \mathcal{T} is a tensor space, we mean that there is a natural inner product $\langle \mathbf{A}, \mathbf{B} \rangle_{\mathcal{T}}$ between any $\mathbf{A}, \mathbf{B} \in \mathcal{T}$ and for every d -dimensional rotation \mathbf{R} there exists an associated linear operator $\mathbf{Q}(\mathbf{R})$ acting on \mathcal{T} such that $\langle \mathbf{Q}(\mathbf{R})\mathbf{A}, \mathbf{Q}(\mathbf{R})\mathbf{B} \rangle_{\mathcal{T}} = \langle \mathbf{A}, \mathbf{B} \rangle_{\mathcal{T}}$ for all $\mathbf{A}, \mathbf{B} \in \mathcal{T}$. Thus, for example, \mathcal{T} could consist of vectors that have a combination of scalars, vectors, second-order tensors, or higher-order tensors (or even tensors of “half integer” order, like spins in quantum mechanics) as elements. However, the tensorial nature of \mathcal{T} is rather moot in this paper as we are not concerned with the action of rotations on elements of \mathcal{T} . Indeed, (1.2) with $m = d$ could be regarded as the linear elasticity equations with $L_{i\alpha j\beta}$ being the fourth-order elasticity tensor $C_{i\alpha j\beta}$ (that annihilates any antisymmetric component of \mathbf{E}) and with $\mathbf{J}(\mathbf{x})$ and $\mathbf{E}(\mathbf{x})$ being the second-order tensor stress and displacement gradient fields. But mathematically it is the same problem when the components of \mathbf{u} rep-

resent different physical scalar potentials, e.g., such as temperature, electrical potential, and pressure, that are invariant under rotations. Then for any \mathbf{x} , $\mathbf{J}(\mathbf{x})$ and $\mathbf{E}(\mathbf{x})$ are not second-order tensors, but rather triplets of vector fields.

Our paper also presents a broad theory of boundary field equalities that generalize the notion of a conservation law. These boundary field equalities imply, for example, that the “Dirichlet-to-Neumann map” (DtN map) governing the response of inhomogeneous bodies satisfies certain exact identities when the tensor field $\mathbf{L}(\mathbf{x})$ inside Ω takes values in a certain nonlinear manifold \mathcal{M} . These identities generalize the exact identities satisfied by the effective tensor \mathbf{L}_* in the theory of exact relations for composites when $\mathbf{L}(\mathbf{x})$ inside the period cell takes values in \mathcal{M} .

The classic conservation law says that if a vector field $\mathbf{u}(\mathbf{x})$ with d -components satisfies $\nabla \cdot \mathbf{u} = 0$ inside a body Ω , then the integral of $\mathbf{u} \cdot \mathbf{n}$ over the surface $\partial\Omega$ of Ω is zero: here \mathbf{n} denotes the outward normal of Ω . This naturally leads to the question: can one make other assumptions about the fields inside a body (still leaving many degrees of freedom in the choice of these fields) that imply exact “boundary field equalities” among the fields at the boundary for suitable boundary conditions? Of course, these boundary conditions should not be such that they trivially imply the boundary field equalities, independent of any assumption about the fields inside the body. *We emphasize that, in general, our boundary field equalities do not result from integration by parts, but rather arise through algebraic properties of the underlying operators. Thus, it is an entirely new idea to obtaining identities satisfied by the boundary fields.*

A divergence-free field satisfies a differential constraint, but additional algebraic constraints are also possible. An example of the latter type of boundary field equality, discussed in [35] Sect. 1.5, and implicit in the work of Milgrom [23] (see also [26] Chap. 6, and references therein) is the following one. Consider in a body Ω the primary equations (1.2) with no source term, i.e., $h_{ij}(\mathbf{x}) = 0$ for all i, j , and with $L_{i\alpha j\beta}(\mathbf{x}) = \delta_{ij}A_{\alpha\beta}(\mathbf{x})$. Assume that Ω contains just two phases in any configuration where the $m \times m$ matrix-valued field $\mathbf{A}(\mathbf{x})$ takes the value $\mathbf{A}^{(1)}$ in phase 1 and $\mathbf{A}^{(2)}$ in phase 2, in which $\mathbf{A}^{(1)}$ and $\mathbf{A}^{(2)}$ are real, symmetric, positive definite matrices. Associated with this problem are boundary fields: the vector potential $\mathbf{u}(\mathbf{x})$ on $\partial\Omega$ (that may be obtained by integrating over the surface the tangential values of $\mathbf{E}(\mathbf{x})$) and the vector-valued flux $\mathbf{n} \cdot \mathbf{J}$. The key observation is that there exists a congruence transformation that simultaneously diagonalizes $\mathbf{A}^{(1)}$ and $\mathbf{A}^{(2)}$, i.e., a matrix \mathbf{W} such that $\mathbf{W}\mathbf{A}^{(1)}\mathbf{W}^T$ and $\mathbf{W}\mathbf{A}^{(2)}\mathbf{W}^T$ are simultaneously diagonal. To do this we choose $\mathbf{W} = \mathbf{Q}(\mathbf{A}^{(2)})^{-1/2}$ where \mathbf{Q} satisfying $\mathbf{Q}\mathbf{Q}^T = \mathbf{I}$ is taken to diagonalize $(\mathbf{A}^{(2)})^{-1/2}\mathbf{A}^{(1)}(\mathbf{A}^{(2)})^{-1/2}$. Then one obtains an equivalent set of decoupled conductivity equations:

$$\tilde{J}_{i\alpha}(\mathbf{x}) = \sigma_{\alpha}(\mathbf{x})\tilde{E}_{i\alpha}(\mathbf{x}), \quad \tilde{E}_{i\alpha}(\mathbf{x}) = \frac{\partial \tilde{u}_{\alpha}(\mathbf{x})}{\partial x_i}, \quad \sum_{i=1}^d \frac{\partial \tilde{J}_{i\alpha}(\mathbf{x})}{\partial x_i} = 0, \quad (1.7)$$

indexed by $\alpha = 1, 2, \dots, m$, with

$$\begin{aligned} \tilde{\mathbf{J}}(\mathbf{x}) &= \mathbf{J}(\mathbf{x})\mathbf{W}^T, \quad \tilde{\mathbf{E}}(\mathbf{x}) = \mathbf{E}(\mathbf{x})\mathbf{W}^{-1}, \quad \tilde{\mathbf{u}}(\mathbf{x}) = (\mathbf{W}^T)^{-1}\mathbf{u}(\mathbf{x}), \\ \sigma_{\alpha}(\mathbf{x}) &= \{\mathbf{Q}(\mathbf{A}^{(2)})^{-1/2}\mathbf{A}^{(1)}(\mathbf{A}^{(2)})^{-1/2}\mathbf{Q}^T\}_{\alpha\alpha} \quad \text{for } \mathbf{x} \text{ in phase 1,} \\ &= 1 \quad \text{for } \mathbf{x} \text{ in phase 2.} \end{aligned} \quad (1.8)$$

It is then clear that if the boundary values of $\mathbf{u}(\mathbf{x})$ are prescribed so that $\tilde{\mathbf{u}}(\mathbf{x}) = (\mathbf{W}^T)^{-1}\mathbf{u}(\mathbf{x})$ only has one nonzero component, then certainly the flux $\mathbf{n} \cdot \tilde{\mathbf{J}}(\mathbf{x}) = \mathbf{n} \cdot \mathbf{J}(\mathbf{x})\mathbf{W}^T$ will only have one matching nonzero component. In other words the flux $\mathbf{n} \cdot \mathbf{J}(\mathbf{x})$ is of the form $\alpha(\mathbf{x})\mathbf{v}$ where

only the scalar $\alpha(\mathbf{x})$ varies on the surface of Ω . These constraints on the flux $\mathbf{n} \cdot \mathbf{J}(\mathbf{x})$ for the prescribed $\mathbf{u}(\mathbf{x})$ are an example of a boundary field equality. Another example of a boundary field equality is that given in [48] for an elastic body containing two isotropic elastic phases having the same shear modulus, where the boundary field equality involves the volume fraction (and thus may be used in an inverse way to determine this volume fraction).

These simple examples serve to give an idea of boundary field equalities, but the general theory developed here goes far beyond them. Rather than say considering inside Ω the constitutive law $\mathbf{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{E}(\mathbf{x})$ with the constraint that $\mathbf{L}(\mathbf{x}) \in \mathcal{M}$, we may eliminate $\mathbf{L}(\mathbf{x})$ and just view these relations as a nonlinear local constraint on the fields. Then, we obtain results like the following. Suppose we are given a $d \times s$ matrix-valued field $\mathbf{Q}(\mathbf{x})$ such that $\nabla \cdot \mathbf{Q} = 0$ [so each column of $\mathbf{Q}(\mathbf{x})$ represents a single divergence-free vector field]. Let $\mathbf{q}(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}(\mathbf{x})$ be the associated s -component flux $\mathbf{q}(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}(\mathbf{x})$ at the boundary of Ω [where $\mathbf{n}(\mathbf{x})$ is the outward normal on $\partial\Omega$]. With $\mathbf{Q}(\mathbf{x})$ constrained to take values in a subset \mathcal{B} of some r -dimensional nonlinear manifold [where \mathcal{B} does not depend on \mathbf{x}], we find conditions on \mathcal{B} and nonlocal linear constraints on the boundary flux \mathbf{q} (specified in Sect. 8) which forces the nonuniquely determined field $\mathbf{Q}(\mathbf{x})$ inside Ω to lie in a subspace \mathcal{D} and hence which forces $\mathbf{q}(\mathbf{x})$ to lie in $\mathbf{n}(\mathbf{x}) \cdot \mathcal{D}$ [where $\mathbf{n}(\mathbf{x}) \cdot \mathcal{D}$ is obtained by applying $\mathbf{n}(\mathbf{x}) \cdot$ to each element of \mathcal{D}]. Of course, it should not be the case that $\mathcal{B} \subset \mathcal{D}$, since otherwise the result would be trivial.

Alternatively, we may express \mathbf{Q} in terms of the elements gradient of t -component potential \mathbf{w} , and we obtain results like the following. Suppose $\nabla \mathbf{w}(\mathbf{x}) \in \mathcal{A}$ for all \mathbf{x} , where \mathcal{A} is a subset of a nonlinear manifold (that does not depend on \mathbf{x}), then we find conditions on \mathcal{A} and nonlocal linear constraints on the surface potential $\mathbf{w}(\mathbf{x})$, $\mathbf{x} \in \partial\Omega$, (specified in Sect. 8) which forces the nonuniquely determined field $\nabla \mathbf{w}(\mathbf{x})$ inside Ω to lie in a subspace \mathcal{C} , and this then places restrictions on the tangential derivatives of the surface potential. Again, it should not be the case that $\mathcal{A} \subset \mathcal{C}$, since otherwise the result would be trivial. To better understand the significance of the constraint that inside Ω , $\nabla \mathbf{w}(\mathbf{x}) \in \mathcal{C}$, let \mathbf{N} be any $d \times t$ matrix normal to the space \mathcal{C} , i.e., such that $\text{Tr}(\mathbf{C}\mathbf{N}^T) = 0$ for all $\mathbf{C} \in \mathcal{C}$. Then $\text{Tr}[\nabla \mathbf{w}(\mathbf{x})\mathbf{N}^T] = 0$ which implies $\nabla \cdot [\mathbf{N}\mathbf{w}(\mathbf{x})] = 0$, i.e., $\mathbf{N}\mathbf{w}(\mathbf{x})$ is a divergence-free field. We deduce that

$$\int_{\partial\Omega} \mathbf{n}(\mathbf{x}) \cdot [\mathbf{N}\mathbf{w}(\mathbf{x})] \, dS = 0, \tag{1.9}$$

which implies the additional boundary field equality:

$$\int_{\partial\Omega} \mathbf{n}(\mathbf{x}) \otimes \mathbf{w}(\mathbf{x}) \, dS \in \mathcal{C}. \tag{1.10}$$

An explanation of why there exist such subsets \mathcal{B} and \mathcal{A} of nonlinear manifolds with this property, and a prescription for obtaining them and the appropriate boundary conditions, will be given in Sect. 8.

Somewhat related questions have been the focus of attention in the homogenization community. Luc Tartar [45] raised (in a more general setting) essentially this fundamental question: if one has a sequence of fields $\mathbf{u}_\epsilon(\mathbf{x})$ such that $\nabla \mathbf{u}_\epsilon(\mathbf{x})$ takes values in a set \mathcal{A} , then what is the range of values that the weak limits of $\nabla \mathbf{u}_\epsilon(\mathbf{x})$ can take? [Alternatively, if one has a sequence of fluxes $\mathbf{Q}_\epsilon(\mathbf{x})$ such that $\nabla \cdot \mathbf{Q}_\epsilon(\mathbf{x}) = 0$ and $\mathbf{Q}_\epsilon(\mathbf{x})$ takes values in a set \mathcal{B} and converges weakly to $\mathbf{Q}_0(\mathbf{x})$, then what is the range of values that $\mathbf{Q}_0(\mathbf{x})$ can take?]. For a sample of work addressing this thorny problem, see, for example, the papers [12, 37, 51] and references therein. Here we are tackling the question of whether, for certain sets \mathcal{A} ,

one can deduce additional constraints on each element $\nabla \mathbf{u}_\epsilon(\mathbf{x})$ in the entire sequence, namely that $\nabla \mathbf{u}_\epsilon(\mathbf{x}) \in \mathcal{C}$, when $\mathbf{u}_\epsilon(\mathbf{x})$ satisfies appropriate boundary conditions. This is important as the homogenization approach is not suitable in applications where there is no separation of length scales. In cases where it is applicable, we deduce the nontrivial result that the weak limit of $\nabla \mathbf{u}_\epsilon(\mathbf{x})$ lies in \mathcal{C} (that is, essentially, a corollary of the theory of exact relations for composites). We emphasize that our boundary field equalities are generally not simply an application of integration by parts, not even at a qualitative level. Rather they follow from algebraic identities.

We mention too that beyond boundary field equalities there are boundary field inequalities and some of these go beyond just using convexity and the divergence theorem [27, 28].

Although we do not explicitly address this in the paper, we mention here that our results on boundary field equalities enable us to introduce and give examples of what we call “partial null Lagrangians.” Null Lagrangians include, for example, functions $F(\mathbf{x}, \mathbf{w}, \nabla \mathbf{w})$ for which the corresponding integral

$$\mathcal{W}(\mathbf{w}) = \int_{\Omega} F(\mathbf{x}, \mathbf{w}(\mathbf{x}), \nabla \mathbf{w}(\mathbf{x})) \, d\mathbf{x}, \quad (1.11)$$

has the property that $\mathcal{W}(\mathbf{w}_0 + \boldsymbol{\phi}) = \mathcal{W}(\mathbf{w}_0)$ for any choice of $\boldsymbol{\phi} \in C_0^\infty(\Omega)$ and for any choice of $\mathbf{w}_0 \in C^1(\overline{\Omega})$. Null Lagrangians of this form have been completely characterized by Olver and Sivaloganathan [41]. It is well known (see the references in [41]) that when $F(\mathbf{x}, \mathbf{w}, \nabla \mathbf{w}) = \Psi(\nabla \mathbf{w})$, then $\Psi(\nabla \mathbf{w})$ is a null Lagrangian if and only if it is an affine combination of subdeterminants of $\nabla \mathbf{w}$ of all orders. Other classes of null Lagrangian have been characterized by Murat [38–40] (see also Pedregal [42]). Null Lagrangians are also instrumental in the construction of polyconvex functions and play a fundamental role in the calculus of variations and in establishing the existence and uniqueness of minimizers to large classes of “energy functions” (see, for example, [3], the recent review [5], and references therein). Additionally, they are an important tool for establishing bounds on the effective moduli of composite materials, through the “translation method,” or equivalently, the method of “compensated compactness” (as summarized in the books [1, 6, 26, 47, 49]). In the liquid crystal community it is well known that if a t -component vector field $\mathbf{w}(\mathbf{x})$ takes values in \mathcal{A} , where \mathcal{A} consists of vectors of unit length, so that $|\mathbf{w}(\mathbf{x})| = 1$, then $(\nabla \mathbf{w}) \cdot \mathbf{w} = \nabla(\mathbf{w} \cdot \mathbf{w}) = 0$. For any given d -component vector field $\mathbf{a}(\mathbf{x})$, the function $F(\mathbf{x}, \mathbf{w}, \nabla \mathbf{w}) = \mathbf{a}(\mathbf{x}) \cdot (\nabla \mathbf{w}) \cdot \mathbf{w}$ is then an example of what we call a partial null Lagrangian: its integral can be exactly computed under the constraint that $\mathbf{w}(\mathbf{x}) \in \mathcal{A}$ for all $\mathbf{x} \in \Omega$ once one knows the boundary fields (and, in this example, the integral is zero and independent of the boundary fields).

More generally, we call $F(\mathbf{x}, \mathbf{w}, \nabla \mathbf{w})$ a partial null Lagrangian on a subset \mathcal{A} (independent of \mathbf{x}) of a nonlinear manifold in the space of pairs of t -component vectors and $d \times t$ matrices, if for every $\boldsymbol{\phi} \in C_0^\infty$ and $\mathbf{w}_0 \in C^1(\overline{\Omega})$ satisfying

$$(\mathbf{w}_0(\mathbf{x}), \nabla \mathbf{w}_0(\mathbf{x})) \in \mathcal{A}, \quad (\mathbf{w}_0(\mathbf{x}) + \boldsymbol{\phi}(\mathbf{x}), \nabla(\mathbf{w}_0(\mathbf{x}) + \boldsymbol{\phi}(\mathbf{x}))) \in \mathcal{A}, \quad \forall \mathbf{x} \in \Omega, \quad (1.12)$$

and with the surface fields $\mathbf{w}_0(\mathbf{x})$, $\mathbf{x} \in \partial\Omega$, satisfying appropriate nonlocal boundary conditions, one has $\mathcal{W}(\mathbf{w}_0 + \boldsymbol{\phi}) = \mathcal{W}(\mathbf{w}_0)$. With \mathcal{A} , and the boundary conditions on \mathbf{w}_0 chosen so it forces $\nabla(\mathbf{w}_0 + \boldsymbol{\phi})$ to lie in a subspace \mathcal{C} , we obtain functions $F(\mathbf{w}, \nabla \mathbf{w})$ that are partial null Lagrangians, but not null Lagrangians. Specifically, since $\mathbf{N}\mathbf{w}(\mathbf{x})$ is a divergence-free field [see the text preceding (1.9)], an obvious partial null Lagrangian is any component

of the t -component function,

$$\mathbf{F}(\mathbf{w}, \nabla \mathbf{w}) = [\mathbf{N}\mathbf{w}] \cdot \nabla \mathbf{w}, \tag{1.13}$$

and we have

$$\int_{\Omega} \mathbf{F}(\mathbf{w}(\mathbf{x}), \nabla \mathbf{w}(\mathbf{x})) \, d\mathbf{x} = \int_{\partial\Omega} [\mathbf{n}(\mathbf{x}) \cdot \mathbf{N}\mathbf{w}(\mathbf{x})] \mathbf{w}(\mathbf{x}) \, dS. \tag{1.14}$$

In two dimensions, if \mathbf{N}_1 and \mathbf{N}_2 are two $2 \times t$ matrices normal to the space \mathcal{C} , then the fact that $\nabla \cdot [\mathbf{N}_1 \cdot \mathbf{w}(\mathbf{x})] = 0$ allows us to find a potential $W_1(\mathbf{x})$ such that

$$\mathbf{R}^\perp \mathbf{N}_1 \cdot \mathbf{w}(\mathbf{x}) = \nabla W_1, \quad \mathbf{R}^\perp = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \tag{1.15}$$

Furthermore, $\mathbf{n}(\mathbf{x}) \cdot [\mathbf{N}_1 \mathbf{w}(\mathbf{x})]$ gives us the tangential derivatives of $\nabla W_1(\mathbf{x})$, which when integrated gives the surface potential $W_1(\mathbf{x})$, $\mathbf{x} \in \partial\Omega$. Then

$$F(\mathbf{x}, \mathbf{w}, \nabla \mathbf{w}) = [\mathbf{N}_2 \mathbf{w}(\mathbf{x})] \cdot \mathbf{R}^\perp \mathbf{N}_1 \mathbf{w}(\mathbf{x}) \tag{1.16}$$

is a partial null Lagrangian and we have

$$\int_{\Omega} F(\mathbf{w}(\mathbf{x}), \nabla \mathbf{w}(\mathbf{x})) \, d\mathbf{x} = \int_{\partial\Omega} [\mathbf{n}(\mathbf{x}) \cdot \mathbf{N}_2 \mathbf{w}(\mathbf{x})] W_1(\mathbf{x}) \, dS. \tag{1.17}$$

We remark that, in the context of this paper, the constraint that $\mathbf{Q}(\mathbf{x}) \in \mathcal{B}$, or equivalently that $\nabla \mathbf{w} \in \mathcal{A}$, is automatically satisfied when there are appropriate materials inside Ω with a bounded and coercive tensor field $\mathbf{L}(\mathbf{x}) \in \mathcal{M}$ for all $\mathbf{x} \in \Omega$. Then, if the appropriate boundary conditions are satisfied, the partial null Lagrangians place integral constraints on the fields inside Ω .


As our work has as its basis the theory of exact relations for composite materials, let us briefly review this.

2 A brief review of exact relations in composites

In this setting, one typically starts with a tensor field $\mathbf{L}(\mathbf{y})$ that is periodic in \mathbf{y} and which is a linear map from \mathcal{T} to \mathcal{T} , where \mathcal{T} is some q -dimensional inner product space. Here $\mathbf{L}(\mathbf{y})$ may represent the conductivity tensor, dielectric tensor, elasticity tensor, or a wealth of other physical tensor fields (see, for example, [26] Chap. 12). In homogenization theory one often considers a body Ω filled by a material having tensor field $\mathbf{L}(\mathbf{x}/\epsilon)$ and in the limit $\epsilon \rightarrow 0$ the body often responds to external fields (that are independent of ϵ) as if it were filled with a homogeneous medium with tensor \mathbf{L}_* that is known as the effective tensor of the medium. In many problems the problem of determining \mathbf{L}_* can be formulated as a problem in the abstract theory of composites. The setting is a Hilbert space \mathcal{H} , say of periodic fields that are square integrable in the unit cell of periodicity and which take values in \mathcal{T} . It has a splitting into three orthogonal spaces $\mathcal{H} = \mathcal{U} \oplus \mathcal{E} \oplus \mathcal{J}$. For example, in the conductivity problem $q = d$, \mathcal{U} is the space of d -dimensional vector fields that are constant (independent of \mathbf{y} , where \mathbf{y} can be thought of as a microscale spatial coordinate) and \mathcal{E} represents gradients of periodic scalar-valued potentials, while \mathcal{J} denotes those periodic fields that have zero divergence and zero average value over the unit cell. To determine the effective tensor \mathbf{L}_* one prescribes a field $\mathbf{E}_0 \in \mathcal{U}$ and solves the equations

$$\mathbf{J}_0 + \mathbf{J} = \mathbf{L}(\mathbf{E}_0 + \mathbf{E}), \quad \text{with } \mathbf{J}_0 \in \mathcal{U}, \quad \mathbf{E} \in \mathcal{E}, \quad \mathbf{J} \in \mathcal{J}, \tag{2.1}$$

where the action of $\mathbf{L} : \mathcal{H} \rightarrow \mathcal{H}$ is defined by $\mathbf{B} = \mathbf{L}\mathbf{A}$ with $\mathbf{B}(\mathbf{y}) = \mathbf{L}(\mathbf{y})\mathbf{A}(\mathbf{y})$ (i.e., \mathbf{L} acts locally in space). Of course \mathbf{L} needs to be such that these equation have a unique solution




for \mathbf{E} (hence uniquely giving \mathbf{J}_0 and \mathbf{J}) for any $\mathbf{E}_0 \in \mathcal{U}$. Clearly \mathbf{J}_0 depends linearly on \mathbf{E}_0 and it is this linear relation that defines the effective tensor: $\mathbf{J}_0 = \mathbf{L}_* \mathbf{E}_0$. This formulation which stems from ideas in [9, 19] was crystallized in [24, 25], see also [26] Chap. 12.

In the field of composites there are a myriad of results on what are known as exact relations: microstructure independent formulae satisfied by effective moduli. A canonical example is Dykhne’s result [10] for 2D conductivity that if the determinant of the local (anisotropic) conductivity tensor $\mathbf{L}(\mathbf{x}) = \boldsymbol{\sigma}(\mathbf{x})$ is constant, then the effective conductivity tensor $\mathbf{L}_* = \boldsymbol{\sigma}_*$ has the same determinant. More generally, as illustrated in Fig. 1, in the theory of exact relations, one wants to find nonlinear manifolds \mathcal{M} (of dimension less than q^2) in the space $L(\mathcal{T})$ of linear maps $\mathcal{T} \rightarrow \mathcal{T}$ such the effective tensor \mathbf{L}_* lies in \mathcal{M} whenever $\mathbf{L}(\mathbf{y})$ lies in \mathcal{M} for all \mathbf{y} (and generally $\mathbf{L}(\mathbf{y})$ also satisfies some sort of boundedness and coercivity properties necessary to ensure that \mathbf{L}_* exists and is unique). In the two-dimensional conductivity example \mathcal{M} consists of 2×2 symmetric matrices $\boldsymbol{\sigma}$ such that $\det(\boldsymbol{\sigma}) = c$, where c is a constant parameterizing the manifold.

The general theory of exact relations was founded by Grabovsky [13], and his insight is summarized in Fig. 2. He realized that if an exact relation held for all composites, then it must certainly hold for layered geometries. For lamination with a vector \mathbf{n} perpendicular to the layers, so that $\mathbf{L}(\mathbf{y})$ is just a function of the single variable $\mathbf{n} \cdot \mathbf{y}$, it is convenient to introduce the fractional linear transformation provided by [25, 52]

$$W_{\mathbf{n}}(\mathbf{L}) = [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\boldsymbol{\Gamma}(\mathbf{n})]^{-1}(\mathbf{L} - \mathbf{L}_0), \quad (2.2)$$



in which $\Gamma(\mathbf{n})$ is a certain tensor dependent on \mathbf{L}_0 and the lamination direction \mathbf{n} . $W_n(\mathbf{L})$ applied to the local tensor field $\mathbf{L}(\mathbf{y})$ and effective tensor \mathbf{L}_* gives a new tensor field $\mathbf{K}_n(\mathbf{y}) = W_n(\mathbf{L}(\mathbf{y}))$ and $\mathbf{K}_n^* = W_n(\mathbf{L}_*)$ that are related simply by a linear average $\mathbf{K}_n^* = \langle \mathbf{K}_n \rangle$ (where the angular brackets denote a volume average of $\mathbf{K}_n(\mathbf{y})$ over its unit cell of periodicity). Thus, the relation

$$\mathbf{L}_* = W_n^{-1}(\langle W_n(\mathbf{L}) \rangle), \tag{2.3}$$

determines the effective tensor \mathbf{L}_* . Furthermore, noting that $W_n(\mathbf{L}_0) = 0$, one can take \mathbf{L}_0 to be on the manifold \mathcal{M} to ensure that $W_n(\mathcal{M})$ passes through the origin. (There are other linear lamination formula [2, 46], but it is unclear if the general theory of exact relations can be developed using them, or their generalizations). Therefore the exact relation in these new coordinates must be a linear relation: $\mathbf{K}_n^* \in \mathcal{K}_n$ when $\mathbf{K}_n(\mathbf{y}) \in \mathcal{K}_n$ where the tensor subspace \mathcal{K}_n defines the exact relation: $\mathcal{M} = W_n^{-1}(\mathcal{K}_n)$. Thus, \mathcal{K}_n has the same dimension k_0 as \mathcal{M} . As \mathcal{K}_n remains linear as \mathbf{n} is varied one sees that \mathcal{K}_n must have the property that $W_m[W_n^{-1}(\mathcal{K}_n)] = \mathcal{K}_m$ for all unit vectors \mathbf{m} and \mathbf{n} . As $W_m[W_n^{-1}(\cdot)]$ is a nonlinear transformation, the image of a linear subspace under the transformation is generally a “curved” manifold, so \mathcal{K}_n has to be rather special for its image to be a linear subspace rather than a “curved” manifold. Through perturbation analysis with \mathbf{m} close to \mathbf{n} , Grabovsky established that \mathcal{K}_n must be independent of \mathbf{n} , $\mathcal{K}_n = \mathcal{K}_0$ for all \mathbf{n} , and he established that \mathcal{K}_0 must satisfy the algebraic constraint that for every unit vector $\mathbf{m} \in \mathbb{R}^d$ one has

$$\mathbf{B}\Psi(\mathbf{m})\mathbf{B} \in \mathcal{K}_0, \quad \text{for all } \mathbf{B} \in \mathcal{K}_0, \tag{2.4}$$

where the left-hand side of (2.4) is to be regarded as the composition of three linear maps each mapping \mathcal{T} to \mathcal{T} (or as the product of three $q \times q$ matrices if one takes a basis in \mathcal{T} and

represents each map by a matrix acting on the basis element), and $\Psi(\mathbf{m}) : \mathcal{T} \rightarrow \mathcal{T}$ depends on \mathbf{L}_0 and the differential constraints on the fields relevant to the physical problem under consideration. Explicitly, $\Psi(\mathbf{m})$ is given by

$$\Psi(\mathbf{m}) = \langle \Gamma(\mathbf{n}) \rangle_{\mathbf{n}} - \Gamma(\mathbf{m}) \quad (2.5)$$

where the angular brackets $\langle \cdot \rangle_{\mathbf{n}}$ denote a possibly weighted average over the sphere $|\mathbf{n}| = 1$ (for example, one could take a weighting concentrated at $\mathbf{n} = \mathbf{n}_0$, giving $\langle \Gamma(\mathbf{n}) \rangle_{\mathbf{n}} = \Gamma(\mathbf{n}_0)$) and $\Gamma(\mathbf{k})$ is given by

$$\Gamma(\mathbf{k}) = [\Gamma_1(\mathbf{k})\mathbf{L}_0\Gamma_1(\mathbf{k})]^{-1}\Gamma_1(\mathbf{k}), \quad (2.6)$$

and the inverse is to be taken on the subspace $\mathcal{E}_{\mathbf{k}}$ onto which $\Gamma_1(\mathbf{k})$ projects.

As a simple example, for two-dimensional conductivity with $\mathbf{L}_0 = \sigma_0\mathbf{I}$ as our reference tensor, one sees that the 2×2 matrix

$$\Psi(\mathbf{m}) = (\mathbf{I} - 2\mathbf{m}\mathbf{m}^T)/2\sigma_0 \quad (2.7)$$

is trace free and symmetric. We can then take \mathcal{K}_0 as the subspace of trace-free and symmetric 2×2 matrices. These have the property that the product of three of them (but not just two of them) is again trace free and symmetric: assuming without loss of generality that one matrix is diagonal, we have

$$\begin{pmatrix} a & 0 \\ 0 & -a \end{pmatrix} \begin{pmatrix} b & c \\ c & -b \end{pmatrix} \begin{pmatrix} d & e \\ e & -d \end{pmatrix} = \begin{pmatrix} abd + ace & abe - acd \\ abe - acd & -abd - ace \end{pmatrix}, \quad (2.8)$$


which is again trace free and symmetric, and thus, the algebraic condition (2.4) is satisfied. Then the associated manifold $\mathcal{M} = W_{\mathbf{n}}^{-1}(\mathcal{K}_0)$ consists of 2×2 symmetric matrices with determinant σ_0^2 , and this is the manifold corresponding to the Dykhne [10] exact relation.

Grabovsky's pioneering work, developed further with Sage in [18], provided essential clues that led to the breakthrough result [17] establishing conditions that guarantee an exact relation holds for all composites, and not just laminates. Using carefully devised perturbation expansions that had their basis in [31] Sect. 5, coupled with analytic continuation arguments, one sees [17] that finding exact relations which hold for all composite geometries is tied with identifying tensor subspaces \mathcal{K} such that for all Fourier vectors $\mathbf{k} \neq 0$ one has

$$\mathbf{B}_1\Psi(\mathbf{k})\mathbf{B}_2 \in \mathcal{K}, \quad \text{for all } \mathbf{B}_1, \mathbf{B}_2 \in \mathcal{K}, \quad (2.9)$$

where $\Psi(\mathbf{k})$ only depends on $\mathbf{k}/|\mathbf{k}|$, i.e., $\Psi(\mathbf{k}) = \Psi(\mathbf{m})$ with $\mathbf{m} = \mathbf{k}/|\mathbf{k}|$ and $\Psi(\mathbf{m})$ is the same operator as in (2.5). The space \mathcal{K}_0 then could be \mathcal{K} or it may be just those symmetric or Hermitian matrices in \mathcal{K} . (Previously in [26] Chap. 17, \mathcal{K}_0 and \mathcal{K} had been labeled as \mathcal{K} and $\overline{\mathcal{K}}$, respectively. We choose to drop the overline in $\overline{\mathcal{K}}$ to simplify notation, as this space will be the focus of our analysis). Recently, Grabovsky [16] found a relation that holds for laminate geometries but not more general composites, so the condition (2.4) is not sufficient to guarantee that an exact relation holds for all microstructures: one needs to use (2.9).

The general theory of exact relations is very rich and Grabovsky and collaborators have systematically explored, and with tremendous effort, exact relations for a wide variety of physically important problems, including conductivity with the Hall effect, elasticity, piezoelectricity, thermoelasticity, and thermoelectricity: [15] gives a comprehensive review; see also [26] Chap. 17, and [14]. To simplify the algebra, they assume \mathcal{K} has



rotational invariance properties, so removing this assumption may yield a plethora of additional exact relations. The theory of exact relations encompasses links between effective tensors: an example of such a link, for an isotropic two-phase composite, is Levin’s result [22] that the effective thermal expansion coefficient is known once the effective bulk modulus is measured.

Now one may ask: is there something deeper and more general behind these exact relations? Indeed, it is the purpose of this paper to reveal that there is something deeper. As indicated by the argument presented in Fig. 3, exact relations should apply not only to effective tensors \mathbf{L}_* of periodic composites, but also to Dirichlet-to-Neumann maps of bodies containing inhomogeneous media with inhomogeneities that are not necessarily small compared to the dimensions of Ω . We formulate the problem slightly differently: in place of fields in \mathcal{U} are source terms, and we no longer require the subspaces \mathcal{E} and \mathcal{J} to be comprised of periodic fields, but rather fields that are square integrable over \mathbb{R}^d .

3 Functional framework

Definition 3.1 Let $\mathcal{H} = \mathcal{J} \oplus \mathcal{E}$ be a tensor space of functions with \mathbb{R}^d as domain (e.g., $L^2(\mathbb{R}^d) \otimes \mathcal{T}$, for a corresponding tensor space \mathcal{T}) such that the projection Γ_1 onto \mathcal{E} acts locally in Fourier space, i.e., if $\mathbf{E} = \Gamma_1 \mathbf{A}$ then the Fourier components $\widehat{\mathbf{E}}(\mathbf{k})$ and $\widehat{\mathbf{A}}(\mathbf{k})$ of \mathbf{E} and \mathbf{A} are related via $\widehat{\mathbf{E}}(\mathbf{k}) = \Gamma_1(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k})$ for some operator $\Gamma_1(\mathbf{k})$ that projects onto a subspace $\mathcal{E}_{\mathbf{k}} \subset \mathcal{T}$.

In the case of the primary equations (1.2), \mathcal{E} can be taken as the set of square-integrable fields $\mathbf{E}(\mathbf{x})$ such that $\mathbf{E} = \nabla \mathbf{u}$ for some m -component potential \mathbf{u} , \mathcal{J} can be taken as the set of square-integrable fields $\mathbf{J}(\mathbf{x})$ such that $\nabla \cdot \mathbf{J} = 0$, $\mathcal{E}_{\mathbf{k}}$ consists of rank-one $d \times m$ matrices of the form $\mathbf{k} \otimes \mathbf{a}$, where $\mathbf{a} \in \mathbb{C}^m$, and consequently the action of $\Gamma_1(\mathbf{k})$ is given by $\Gamma_1(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k}) = \mathbf{k} \otimes (\mathbf{k} \cdot \widehat{\mathbf{A}}(\mathbf{k})) / |\mathbf{k}|^2$ for $\mathbf{k} \neq 0$ and $\Gamma_1(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k}) = 0$ when $\mathbf{k} = 0$. For wider classes of partial differential equations, involving higher-order derivatives, an explicit formula for $\Gamma_1(\mathbf{k})$ is given, for example, in Sect. 12.2 of [26], and in [27, 28].

Let q denote the dimension of \mathcal{T} , and let \mathcal{L} be the space of linear operators $\mathbf{A} : \mathcal{H} \rightarrow \mathcal{H}$. Consider $\mathbf{L}, \mathbf{L}_0 \in \mathcal{L}$ defined as $\mathbf{L}(\mathbf{R})(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{R}(\mathbf{x})$ and, respectively, $\mathbf{L}_0(\mathbf{R})(\mathbf{x}) = \mathbf{L}_0\mathbf{R}(\mathbf{x})$, for $\mathbf{R} \in \mathcal{H}$ where $\mathbf{L}(\mathbf{x}), \mathbf{L}_0 \in L^\infty(\mathbb{R}^d) \otimes L(\mathcal{T})$ with \mathbf{L}_0 denoting a given constant tensor.

(Thus, \mathbf{L} acts locally in space while \mathbf{L}_0 acts locally both in space and in Fourier space). Assume that \mathbf{L} is self-adjoint, bounded, and coercive, i.e., there exist constants $\beta_0 > \alpha_0 > 0$ such that

$$\beta_0 \mathbf{I} \geq \mathbf{L}(\mathbf{x}) \geq \alpha_0 \mathbf{I} > 0, \quad \text{for all } \mathbf{x} \in \mathbb{R}^d, \quad (3.1)$$

where the inequalities hold in the sense of the associated quadratic forms. We emphasize that many physical problems where \mathbf{L} is not self-adjoint, including those where \mathbf{L} is complex and symmetric with a positive definite imaginary part, can be converted to equivalent problems taking the required form (1.5), where the new \mathbf{L} is self-adjoint, bounded and coercive (see, [7, 25] Sect. 18, [6] Chap. 13, [26] Sect. 12.11, [32, 35] Sect. 5.2). We will next consider linear PDEs admitting a formulation in the following canonical form (see [35]),

$$\mathbf{J} = \mathbf{L}\mathbf{E} - \mathbf{h}, \quad \text{with } \mathbf{J} \in \mathcal{J}, \quad \mathbf{E} \in \mathcal{E}, \quad \mathbf{h} \in (\mathbf{L} - \mathbf{L}_0)\mathcal{H}. \quad (3.2)$$

This is a restricted form of (1.5) since in general $\mathbf{L} - \mathbf{L}_0$ may be singular and thus $(\mathbf{L} - \mathbf{L}_0)\mathcal{H}$ does not equal \mathcal{H} . The boundedness and coercivity conditions (3.1) ensure that problem (3.2) has a unique solution for every $\mathbf{h} \in (\mathbf{L} - \mathbf{L}_0)\mathcal{H}$. Since $\mathbf{h} \in (\mathbf{L} - \mathbf{L}_0)\mathcal{H}$ we can equivalently let $\mathbf{h} = -(\mathbf{L} - \mathbf{L}_0)\mathbf{s}$ with $\mathbf{s} \in \mathcal{H}$ and consider the equation

$$\mathbf{J} = \mathbf{L}\mathbf{E} + (\mathbf{L} - \mathbf{L}_0)\mathbf{s}, \quad \text{with } \mathbf{J} \in \mathcal{J}, \quad \mathbf{E} \in \mathcal{E}, \quad \mathbf{s} \in \mathcal{H}. \quad (3.3)$$

Definition 3.2 Following [26] let us introduce the following operator $\Gamma : \mathcal{H} \rightarrow \mathcal{H}$, defined by $\Gamma \mathbf{A} = \mathbf{E}$ if and only if $\mathbf{E} \in \mathcal{E}$ and $\mathbf{A} - \mathbf{L}_0 \mathbf{E} \in \mathcal{J}$. These equations are easily solved by going to Fourier space and one sees that Γ is univalued, with action in Fourier space given by the following lemma:

Lemma 3.1 *The operator Γ introduced in Definition 3.2, is self-adjoint and acts locally in Fourier space. Explicitly, if $\mathbf{E} = \Gamma \mathbf{A}$ then the Fourier components $\widehat{\mathbf{E}}(\mathbf{k})$ and $\widehat{\mathbf{A}}(\mathbf{k})$ of \mathbf{E} and \mathbf{A} are related via $\widehat{\mathbf{E}}(\mathbf{k}) = \Gamma(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k})$, where $\Gamma(\mathbf{k})$ is defined by (2.6).*

To establish the lemma, suppose $\mathbf{E} \in \mathcal{E}$ and $\mathbf{A} - \mathbf{L}_0 \mathbf{E} \in \mathcal{J}$. This, and the orthogonality of \mathcal{E} and \mathcal{J} , implies that for all \mathbf{k} the Fourier components $\widehat{\mathbf{E}}(\mathbf{k})$ and $\widehat{\mathbf{A}}(\mathbf{k}) - \mathbf{L}_0 \widehat{\mathbf{E}}(\mathbf{k})$ lie in $\mathcal{E}_{\mathbf{k}}$ and its orthogonal complement, respectively. Recalling that $\Gamma_1(\mathbf{k})$ projects onto $\mathcal{E}_{\mathbf{k}}$, we obtain

$$0 = \Gamma_1(\mathbf{k})[\widehat{\mathbf{A}}(\mathbf{k}) - \mathbf{L}_0 \widehat{\mathbf{E}}(\mathbf{k})] = \Gamma_1(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k}) - [\Gamma_1(\mathbf{k})\mathbf{L}_0\Gamma_1(\mathbf{k})]\Gamma_1(\mathbf{k})\widehat{\mathbf{E}}(\mathbf{k}), \quad (3.4)$$

and this is easily solved for $\widehat{\mathbf{E}}(\mathbf{k})$, yielding $\widehat{\mathbf{E}}(\mathbf{k}) = \Gamma(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k})$, where $\Gamma(\mathbf{k})$ is defined by (2.6). The expression for $\Gamma(\mathbf{k})$ can equivalently be rewritten as

$$\Gamma(\mathbf{k}) = \Gamma_1(\mathbf{k})[\Gamma_1(\mathbf{k})\mathbf{L}_0\Gamma_1(\mathbf{k})]^{-1}\Gamma_1(\mathbf{k}), \quad (3.5)$$

(where the inverse is to taken on the space $\mathcal{E}_{\mathbf{k}}$) which is evidently self-adjoint, as $\Gamma_1(\mathbf{k})$ and \mathbf{L}_0 are self-adjoint. Note also that $\mathbf{L}_0\Gamma(\mathbf{k})$ can also be interpreted as a nonorthogonal projection onto $\mathbf{L}_0\mathcal{E}$ along \mathcal{J} .

Let $\mathbf{M} : \mathcal{T} \rightarrow \mathcal{T}$ be a self-adjoint positive semidefinite operator. Using the definitions of $\mathbf{L} \in \mathcal{L}$ and $\mathbf{L}_0 \in \mathcal{L}$, we define the operator $\mathbf{K} : \mathcal{H} \rightarrow \mathcal{H}$ as $\mathbf{K}(\mathbf{R})(\mathbf{x}) = \mathbf{K}(\mathbf{x})\mathbf{R}(\mathbf{x})$ with

$$\mathbf{K}(\mathbf{x}) = W_{\mathbf{M}}(\mathbf{L}(\mathbf{x})) = [\mathbf{I} + (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbf{M}]^{-1}(\mathbf{L}(\mathbf{x}) - \mathbf{L}_0) \in L(\mathcal{T}). \quad (3.6)$$

It follows from Grabovsky's Definition 3.17 and Lemma 3.18 [15] that $\mathbf{I} + (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbf{M}$ is invertible when $\mathbf{M}\mathbf{L}_0\mathbf{M} \leq \mathbf{M}$, and so under this assumption the fractional linear transformation $W_{\mathbf{M}} : L(\mathcal{T}) \rightarrow \mathcal{L}(\mathcal{T})$ given by (3.6) is well defined.

For $\lambda \in [0, 1]$ consider the sequence of operators $L_\lambda : \mathcal{H} \rightarrow \mathcal{H}$ defined by

$$L_\lambda = L_0 + \lambda[\mathbf{I} + (1 - \lambda)(\mathbf{L} - L_0)\mathbf{M}]^{-1}(\mathbf{L} - L_0), \tag{3.7}$$

and note that $L_{\lambda=1} = L$ and $L_{\lambda=0} = L_0$. We also point out that L_λ is just a homothety by λ in the W variables, i.e., $W_{\mathbf{M}}(L_\lambda) = \lambda W_{\mathbf{M}}(L)$. In particular, in the context of composites L_λ with $\mathbf{M} = \Gamma(\mathbf{n})$ is in fact the effective tensor of a laminate in direction \mathbf{n} of L and L_0 with a volume fraction λ of L .

Here we assume these operators L_λ are well defined (which is a consequence of Theorem 4.2 under some restrictions on L and L_0). Let $\Psi \in \mathcal{L}$, be defined by $\Psi = \mathbf{M} - \Gamma$. As Γ acts locally in Fourier space, if $\mathbf{B} = \Psi\mathbf{A}$, then the Fourier components $\widehat{\mathbf{B}}(\mathbf{k})$ and $\widehat{\mathbf{A}}(\mathbf{k})$ of \mathbf{B} and \mathbf{A} satisfy a local relation $\widehat{\mathbf{B}}(\mathbf{k}) = \Psi(\mathbf{k})\widehat{\mathbf{A}}(\mathbf{k})$ with $\Psi(\mathbf{k}) = \mathbf{M} - \Gamma(\mathbf{k})$ taking values in $L(\mathcal{T})$. Assume there exists a subspace $\mathcal{K} \subset L(\mathcal{T})$ such that for all \mathbf{k}

$$\mathcal{K}\Psi(\mathbf{k})\mathcal{K} \doteq \{\mathbf{B}_1\Psi(\mathbf{k})\mathbf{B}_2, \text{ for } \mathbf{B}_1, \mathbf{B}_2 \in \mathcal{K}\} \subset \mathcal{K}, \tag{3.8}$$

in which $\mathbf{B}_1\Psi(\mathbf{k})\mathbf{B}_2$ is the composition of the three maps $\mathbf{B}_2 : \mathcal{T} \rightarrow \mathcal{T}$, $\Psi(\mathbf{k}) : \mathcal{T} \rightarrow \mathcal{T}$, and $\mathbf{B}_1 : \mathcal{T} \rightarrow \mathcal{T}$. If (3.8) holds for all \mathbf{k} , then it clearly holds if $\Psi(\mathbf{k})$ is replaced by any tensor \mathbf{A} in the subspace \mathcal{A} spanned by the $\Psi(\mathbf{k})$ as \mathbf{k} varies. Hence, (3.8) can be rewritten as

$$\mathcal{K}\mathcal{A}\mathcal{K} \doteq \{\mathbf{B}_1\mathbf{A}\mathbf{B}_2, \text{ for } \mathbf{B}_1, \mathbf{B}_2 \in \mathcal{K}, \mathbf{A} \in \mathcal{A}\} \subset \mathcal{K}. \tag{3.9}$$

Spaces \mathcal{K} having this property have been called an associative \mathcal{A} -multialgebra by Grabovsky [15]. Instead of testing that (3.8) holds for all $\Psi(\mathbf{k})$ as \mathbf{k} varies, it suffices to test it for a basis of \mathcal{A} .

Next, let us denote by $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_q$ a basis of \mathcal{T} . For given q functions $\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_q \in \mathcal{H}$, consider the following linear map $\mathbb{S} : \mathcal{T} \rightarrow \mathcal{H}$ defined by

$$\mathbb{S}\mathbf{e}_i = \mathbf{s}_i \quad \text{for all } i \in \{1, \dots, q\}. \tag{3.10}$$

We let $\mathbb{S}(\mathbf{x})$ denote the associated field taking for each \mathbf{x} values in $L(\mathcal{T})$ such that $\mathbb{S}(\mathbf{x})\mathbf{e}_i = \mathbf{s}_i(\mathbf{x})$ for all i . This field can be considered to lie in the space $\mathfrak{H} = L^2(\mathbb{R}^d) \otimes L(\mathcal{T})$, endowed with the inner product

$$\langle \mathbb{A}, \mathbb{B} \rangle_{\mathfrak{H}} = \sum_{i=1}^q \langle \mathbb{A}\mathbf{e}_i, \mathbb{B}\mathbf{e}_i \rangle_{\mathcal{H}}. \tag{3.11}$$

Note that any linear operator $\mathbf{F} : \mathcal{H} \rightarrow \mathcal{H}$, such as \mathbf{K} or Ψ , has a natural extension to an operator on \mathfrak{H} : we define

$$\mathbf{F}\mathbb{A} = \mathbb{B} \text{ iff } \mathbf{F}(\mathbb{A}\mathbf{e}_i) = \mathbb{B}\mathbf{e}_i \quad \text{for all } i, \tag{3.12}$$

where, to simplify notation, we use the same symbol for the operator acting on \mathfrak{H} as for the operator acting on \mathcal{H} .

4 The central theorem

Define \mathcal{S} as a subspace of $L(\mathcal{T})$ such that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$. For example, (3.8) implies \mathcal{S} could be taken as $\mathcal{Q} = \mathcal{A}\mathcal{K}$, defined as the space spanned by $\Psi(\mathbf{k})\mathcal{K}$ as \mathbf{k} varies. A natural choice for \mathcal{S} is the largest subspace with the property that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$, although it is then not clear how easily that subspace can be computed. The central theorem of this paper states:

Theorem 4.1 Consider problem (3.2) and let $\mathbf{L}_0, \mathbf{M}, \mathbf{L}$ satisfy all the conditions presented in the previous section. Assume that the following conditions hold:

$$\mathbf{K}(\mathbf{x}) = W_{\mathbf{M}}(\mathbf{L}(\mathbf{x})) \in \mathcal{K} \quad \text{for all } \mathbf{x}, \quad (4.1)$$

$$\mathbf{L}_\lambda \text{ is bounded and coercive on } \mathcal{H} \text{ for all } \lambda \in [0, 1], \quad (4.2)$$

$$\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_q \in \mathcal{H} \text{ are such that } \mathbb{S}(\mathbf{x}) \in \mathcal{S} \quad \text{for all } \mathbf{x}. \quad (4.3)$$

where \mathbf{L}_λ was defined at (3.7) and $W_{\mathbf{M}}$ was defined at (3.6). Next, consider the set of q sources $\mathbf{h}_i = (\mathbf{L} - \mathbf{L}_0)\mathbf{s}_i$, $i = 1, 2, \dots, q$, and let $(\mathbf{E}_i, \mathbf{J}_i)$ ($i = \{1, \dots, q\}$) denote the unique solution of the problem (3.2) for each of the sources $\mathbf{h}_1, \mathbf{h}_2, \dots, \mathbf{h}_q$ respectively. For each solution pair $(\mathbf{E}_i, \mathbf{J}_i)$ define the corresponding polarization field via $\mathbf{P}_i = \mathbf{J}_i - \mathbf{L}_0\mathbf{E}_i$ and introduce the operator $\mathbb{P} : \mathcal{T} \rightarrow \mathcal{H}$ defined by

$$\mathbb{P}\mathbf{e}_i = \mathbf{P}_i \quad \text{for all } i \in \{1, \dots, q\}. \quad (4.4)$$

Associated with \mathbb{P} is the field $\mathbb{P}(\mathbf{x})$ taking values for each \mathbf{x} in $L(\mathcal{T})$ such that $\mathbb{P}(\mathbf{x})\mathbf{e}_i = \mathbf{P}_i(\mathbf{x})$ for all i . Then

$$\mathbb{P}(\mathbf{x}) \in \mathcal{K} \quad \text{for all } \mathbf{x}. \quad (4.5)$$

Proof Our proof of this result has much in common with the proof establishing sufficient conditions for an exact relation to hold for all composite geometries (see [17, 26] Sect. 17.3, [15] Sect. 4.5, and [16]). From (4.2) we have that for any given $\mathbf{h} \in (\mathbf{L} - \mathbf{L}_0)\mathcal{H}$ there exists unique $\mathbf{E}_\lambda \in \mathcal{E}$ and $\mathbf{J}_\lambda \in \mathcal{J}$ that solve (3.2) with $\mathbf{L} = \mathbf{L}_\lambda$. We choose to define the polarization field \mathbf{P}_λ as,

$$\mathbf{P}_\lambda = \mathbf{J}_\lambda - \mathbf{L}_0\mathbf{E}_\lambda = (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{E}_\lambda - \mathbf{h}. \quad (4.6)$$

[Note that the polarization field \mathbf{P}_λ is not $(\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{E}_\lambda$]. Then from Definition 3.2 we have that

$$\mathbf{\Gamma}\mathbf{P}_\lambda = \mathbf{\Gamma}(\mathbf{J}_\lambda - \mathbf{L}_0\mathbf{E}_\lambda) = -\mathbf{E}_\lambda. \quad (4.7)$$

Indeed, (4.7) follows from $-\mathbf{E}_\lambda \in \mathcal{E}$ and $\mathbf{J}_\lambda - \mathbf{L}_0\mathbf{E}_\lambda - \mathbf{L}_0(-\mathbf{E}_\lambda) = \mathbf{J}_\lambda \in \mathcal{J}$. Hence, we obtain

$$\begin{aligned} [\mathbf{I} + (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{\Gamma}]\mathbf{P}_\lambda &= \mathbf{P}_\lambda - (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{E}_\lambda \\ &= (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{E}_\lambda - \mathbf{h} - (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{E}_\lambda \\ &= -\mathbf{h}. \end{aligned} \quad (4.8)$$

Thus, from (4.8) together with the uniqueness of $\mathbf{E}_\lambda, \mathbf{J}_\lambda$, we have that, for all $\lambda \in [0, 1]$,

$$[\mathbf{I} + (\mathbf{L}_\lambda - \mathbf{L}_0)\mathbf{\Gamma}]^{-1} : (\mathbf{L} - \mathbf{L}_0)\mathcal{H} \rightarrow \mathcal{H} \text{ is a well-defined linear operator.} \quad (4.9)$$

With $\lambda = 1$, this result is Grabovsky's corollary 3.19 [15] but this follows in our case from different arguments than in his book. Next, for $\lambda = 1$ in (4.8) we obtain

$$\mathbf{P} = -[\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}]^{-1}\mathbf{h}, \quad (4.10)$$

where in (4.10) and in what follows we use \mathbf{P} instead of $\mathbf{P}_{\lambda=1}$. This may be equivalently rewritten as follows (see [17] Sect 3.2, or [26] Sect. 14.9, for a similar approach)

$$\begin{aligned} \mathbf{P} &= -[\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}]^{-1}\mathbf{h} \\ &= -[\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M} + (\mathbf{L} - \mathbf{L}_0)(\mathbf{\Gamma} - \mathbf{M})]^{-1}\mathbf{h} \\ &= [\mathbf{I} - [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1}(\mathbf{L} - \mathbf{L}_0)(\mathbf{M} - \mathbf{\Gamma})]^{-1}[\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1}(\mathbf{L} - \mathbf{L}_0)\mathbf{s} \\ &= (\mathbf{I} - \mathbf{K}\Psi)^{-1}\mathbf{K}\mathbf{s}, \end{aligned} \quad (4.11)$$

where $\mathbf{h} = -(\mathbf{L} - \mathbf{L}_0)\mathbf{s}$, $\mathbf{K} \in \mathcal{L}$ was introduced at (3.6), and $\Psi = \mathbf{M} - \Gamma$ was defined at (3.8). Using the notation introduced immediately after (4.4), equality (4.11) can be equivalently written as

$$\mathbb{P} = (\mathbf{I} - \mathbf{K}\Psi)^{-1}\mathbf{K}\mathbb{S}, \tag{4.12}$$

where $\mathbb{S} : \mathcal{T} \rightarrow \mathcal{H}$ was defined at (3.10). We choose to regard \mathbb{S} and \mathbb{P} as fields in \mathfrak{H} and regard \mathbf{K} and Ψ in (4.12) as operators acting in \mathfrak{H} , defined according to (3.12). Similarly we can define fields $\mathbb{E}(\mathbf{x})$ and $\mathbb{J}(\mathbf{x})$ via

$$\mathbb{E}(\mathbf{x})\mathbf{e}_i = \mathbf{E}_i(\mathbf{x}), \quad \mathbb{J}(\mathbf{x})\mathbf{e}_i = \mathbf{E}_i(\mathbf{x}), \tag{4.13}$$

and the governing equations become

$$\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x}) + (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbb{S}(\mathbf{x}), \text{ with } \mathbb{E} \in \mathfrak{E}, \quad \mathbb{J} \in \mathfrak{J}, \tag{4.14}$$

in which \mathfrak{E} is comprised of fields \mathbb{E} such that $\mathbb{E}\mathbf{e}_i \in \mathcal{E}$ for all i , while \mathfrak{J} is comprised of fields \mathbb{J} such that $\mathbb{J}\mathbf{e}_i \in \mathcal{J}$ for all i .

Consider the following sequence of related fields:

$$\mathbb{P}_\lambda = (\mathbf{I} - \lambda\mathbf{K}\Psi)^{-1}\lambda\mathbf{K}\mathbb{S}. \tag{4.15}$$

For small λ the Neumann series for $\lambda\mathbf{K}\Psi$ is convergent, and we have

$$\begin{aligned} \mathbb{P}_\lambda &= (\mathbf{I} - \lambda\mathbf{K}\Psi)^{-1}\lambda\mathbf{K}\mathbb{S} \\ &= \sum_{j=0}^{\infty} \lambda^{j+1}(\mathbf{K}\Psi)^j\mathbf{K}\mathbb{S}. \end{aligned} \tag{4.16}$$

It is to be emphasized that in this expansion \mathbf{K} and Ψ are operators: they act on the field in \mathfrak{H} to the right of them. Related expansions in the theory of composites were first introduced in [31], Sect. 5, for the conductivity problem, and their convergence properties, allowing for possibly nonsymmetric conductivity tensors, were studied in [8]. They also form the basis of accelerated iterative Fast Fourier transform (FFT) techniques for evaluating the fields in composites and the associated effective tensors [11] (see also [26] Sects. 14.9 and 14.10) that generally converge faster than the iterative FFT techniques first proposed in [36]. However, the application that motivates their introduction in our paper is their essential role in the theory of exact relations in composites [17].

To prove $\mathbb{P}_\lambda(\mathbf{x})$ takes values in \mathcal{K} when $\mathbb{S}(\mathbf{x})$ takes values in \mathcal{S} , one proceeds by induction. Define the partial sums

$$\mathbb{P}_\lambda^m = \sum_{j=0}^m \lambda^{j+1}(\mathbf{K}\Psi)^j\mathbf{K}\mathbb{S}, \quad \mathbb{Q}_\lambda^m = \Psi \sum_{j=0}^m \lambda^{j+1}(\mathbf{K}\Psi)^j\mathbf{K}\mathbb{S}. \tag{4.17}$$

Clearly these fields, which are in \mathfrak{H} , are related by

$$\mathbb{P}_\lambda^{m+1} = \lambda\mathbf{K}\mathbb{Q}_\lambda^m + \lambda\mathbf{K}\mathbb{S}, \quad \mathbb{Q}_\lambda^{m+1} = \Psi\mathbb{P}_\lambda^m. \tag{4.18}$$

Assume for some m that for every \mathbf{x} , $\mathbb{P}_\lambda^m(\mathbf{x}) \in \mathcal{K}$. This is clearly true when $m = 0$ by the definition of \mathcal{S} . Then the Fourier components of \mathbb{P}_λ^m also lie in \mathcal{K} . It follows that the Fourier components of \mathbb{Q}_λ^m , and hence also the values of $\mathbb{Q}_\lambda^m(\mathbf{x})$, lie in \mathcal{Q} , defined (as in the beginning of this section) as the space spanned by $\Psi(\mathbf{k})\mathcal{K}$ as \mathbf{k} varies. Then since $\mathbf{A}\mathcal{Q} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$, we deduce that $\mathbb{P}_\lambda^{m+1}(\mathbf{x})$ lies in \mathcal{K} for all \mathbf{x} .

We conclude that there exists $\lambda_0 > 0$ such that for $\lambda < \lambda_0$, $\mathbb{P}_\lambda(\mathbf{x})$ defined by (4.15) lies in \mathcal{K} for all \mathbf{x} . This implies that for small enough λ we have

$$f_{\mathbb{F}}(\lambda) = \langle (\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\mathbb{S}, \mathbb{F} \rangle_{\mathfrak{H}} = 0, \text{ for all } \mathbb{F} \in \mathfrak{H} \text{ with } \mathbb{F}(\mathbf{x}) \in \mathcal{K}^\perp \text{ for all } \mathbf{x}, \quad (4.19)$$

in which \mathcal{K}^\perp is the orthogonal complement of \mathcal{K} in the space $L(\mathcal{T})$ with respect to the inner product defined (analogously to (3.11)) by

$$\langle \mathbf{A}, \mathbf{B} \rangle_{L(\mathcal{T})} = \sum_{i=1}^q \langle \mathbf{A} \mathbf{e}_i, \mathbf{B} \mathbf{e}_i \rangle_{\mathcal{T}} \quad \text{for all } \mathbf{A}, \mathbf{B} \in L(\mathcal{T}). \quad (4.20)$$

Next we will prove that $(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\mathbb{S}$ is analytic on an open set $D \subset \mathbb{C}$ with $[\frac{\lambda_0}{2}, 1] \subset D$. Indeed, note that the operator function $(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} : \mathbb{C} \rightarrow \mathcal{L}$ is analytic for λ such that $\frac{1}{\lambda} \in \rho(\mathbf{K}\Psi)$ (where $\rho(\mathbf{K}\Psi)$ denotes the resolvent set of $\mathbf{K}\Psi$) and therefore the function $(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\mathbb{S}$ will be analytic on this set of λ values (see [21] Chap. 17).

Thus, we observe that using the openness of $\rho(\mathbf{K}\Psi)$ it is enough to show that $[1, \frac{2}{\lambda_0}] \subset \rho(\mathbf{K}\Psi)$, as this will imply that there exists an open set D_1 (bounded above and below by positive numbers) with $[1, \frac{2}{\lambda_0}] \subset D_1 \subset \rho(\mathbf{K}\Psi)$ and in turn this will give the existence of an open set $D \subset \left\{ \lambda, \frac{1}{\lambda} \in D_1 \right\}$ with $[\frac{\lambda_0}{2}, 1] \subset D$ such that $(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\mathbb{S}$ is analytic on D .

We have that

$$\begin{aligned} (\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K} &= [\mathbf{I} - \lambda [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1} (\mathbf{L} - \mathbf{L}_0)\Psi]^{-1} \lambda \mathbf{K} \\ &= [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M} - \lambda (\mathbf{L} - \mathbf{L}_0)(\mathbf{M} - \Gamma)]^{-1} [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{M}] \lambda \mathbf{K} \\ &= \lambda [\mathbf{I} + (1 - \lambda)(\mathbf{L} - \mathbf{L}_0)\mathbf{M} + \lambda (\mathbf{L} - \mathbf{L}_0)\Gamma]^{-1} (\mathbf{L} - \mathbf{L}_0) \\ &= \lambda [\mathbf{I} + \lambda [\mathbf{I} + (1 - \lambda)(\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1} (\mathbf{L} - \mathbf{L}_0)\Gamma]^{-1} [\mathbf{I} \\ &\quad + (1 - \lambda)(\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1} (\mathbf{L} - \mathbf{L}_0) \\ &= [\mathbf{I} + (\mathbf{L}_\lambda - \mathbf{L}_0)\Gamma]^{-1} (\mathbf{L}_\lambda - \mathbf{L}_0), \quad \text{for } \lambda \in (0, 1], \end{aligned} \quad (4.21)$$

Then, using the fact that by definition, for $\lambda \neq 0$, we have

$$\mathbf{L}_\lambda - \mathbf{L}_0 = \lambda [\mathbf{I} + (1 - \lambda)(\mathbf{L} - \mathbf{L}_0)\mathbf{M}]^{-1} (\mathbf{L} - \mathbf{L}_0), \quad (4.22)$$

or equivalently,

$$(\mathbf{L}_\lambda - \mathbf{L}_0) = \lambda (\mathbf{L} - \mathbf{L}_0) - (1 - \lambda)(\mathbf{L} - \mathbf{L}_0)\mathbf{M}(\mathbf{L}_\lambda - \mathbf{L}_0), \quad (4.23)$$

which implies

$$\mathbf{L}_\lambda - \mathbf{L}_0 = (\mathbf{L} - \mathbf{L}_0)[\lambda \mathbf{I} - (1 - \lambda)\mathbf{M}(\mathbf{L}_\lambda - \mathbf{L}_0)]. \quad (4.24)$$

We see that $(\mathbf{L}_\lambda - \mathbf{L}_0)\mathcal{H} \subset (\mathbf{L} - \mathbf{L}_0)\mathcal{H}$ and this together with (4.9) gives

$$(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K} \in \mathcal{L} \quad \text{for all } \lambda \in [0, 1], \quad (4.25)$$

and this implies

$$(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\Psi \in \mathcal{L} \Rightarrow -\mathbf{I} + (\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \in \mathcal{L} \Rightarrow (\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \in \mathcal{L} \text{ for } \lambda \in (0, 1].$$

The last result above implies that $[1, \infty) \subset \rho(\mathbf{K}\Psi)$ which in turn as explained above implies that there exists an open set D with $[\lambda_0/2, 1] \subset D$ such that $(\mathbf{I} - \lambda \mathbf{K}\Psi)^{-1} \lambda \mathbf{K}\mathbb{S}$ is analytic on D .

This together with (4.19) and by analytic continuation in the complex plane implies that

$$f_{\mathbb{F}}(1) = \langle (\mathbf{I} - \mathbf{K}\Psi)^{-1} \mathbf{K}\mathbb{S}, \mathbb{F} \rangle_{\mathfrak{H}} = 0, \quad \text{for all } \mathbb{F} \in \mathfrak{H} \text{ with } \mathbb{F}(\mathbf{x}) \in \mathcal{K}^{\perp} \quad \text{for all } \mathbf{x}. \tag{4.26}$$

Thus, as desired, we conclude that $\mathbb{P} = (\mathbf{I} - \mathbf{K}\Psi)^{-1} \mathbf{K}\mathbb{S}$ takes values in \mathcal{K} when $\mathbb{S}(\mathbf{x}) \in \mathcal{S}$ for all \mathbf{x} . □

The fundamental algebraic property (3.8) is clearly an algebraic property of the subspace \mathcal{K} comprised of operators \mathbf{K} mapping \mathcal{T} to \mathcal{T} and is independent of what basis $\mathbf{e}_i, i = 1, 2, 3, \dots, q$, for \mathcal{T} we may choose. Up to now we could apply our theory when the field $\mathbb{S}(\mathbf{x})$ took values in \mathcal{S} . Suppose instead that, for some nonsingular mapping $\mathbf{D} : \mathcal{T} \rightarrow \mathcal{T}$, $\mathbb{S}(\mathbf{x})\mathbf{D}^{-1}$, regarded as the composition of the two maps $\mathbf{D}^{-1} : \mathcal{T} \rightarrow \mathcal{T}$ and $\mathbb{S}(\mathbf{x}) : \mathcal{T} \rightarrow \mathcal{T}$, took values in \mathcal{S} . In this case we can introduce a new basis

$$\mathbf{e}'_i = \mathbf{D}\mathbf{e}_i, \quad i = 1, 2, 3, \dots, q, \tag{4.27}$$

and define $\mathbb{S}'(\mathbf{x})$ as that field taking values in $L(\mathcal{T})$ such that

$$\mathbb{S}'(\mathbf{x})\mathbf{e}'_i = \mathbf{s}_i = \mathbb{S}(\mathbf{x})\mathbf{e}_i = \mathbb{S}(\mathbf{x})\mathbf{D}^{-1}\mathbf{e}'_i, \tag{4.28}$$

implying $\mathbb{S}'(\mathbf{x}) = \mathbb{S}(\mathbf{x})\mathbf{D}^{-1}$. Accordingly, we need to introduce the field $\mathbb{P}'(\mathbf{x})$ as that field taking values in $L(\mathcal{T})$ such that

$$\mathbb{P}'(\mathbf{x})\mathbf{e}'_i = \mathbf{P}_i = \mathbb{P}(\mathbf{x})\mathbf{e}_i = \mathbb{P}(\mathbf{x})\mathbf{D}^{-1}\mathbf{e}'_i, \tag{4.29}$$

implying $\mathbb{P}'(\mathbf{x}) = \mathbb{P}(\mathbf{x})\mathbf{D}^{-1}$. Our theorem says that $\mathbb{P}'(\mathbf{x})$ takes values in \mathcal{K} when $\mathbb{S}'(\mathbf{x})$ takes values in \mathcal{S} , and so we conclude that for all nonsingular $\mathbf{D} : \mathcal{T} \rightarrow \mathcal{T}$ and for any $\mathbf{x} \in \mathbb{R}^d$,

$$\mathbb{P}(\mathbf{x})\mathbf{D}^{-1} \in \mathcal{K} \quad \text{when } \mathbb{S}(\mathbf{x}')\mathbf{D}^{-1} \in \mathcal{S} \text{ for all } \mathbf{x}', \tag{4.30}$$

or equivalently that

$$\mathbb{P}(\mathbf{x}) \in \mathcal{K}\mathbf{D} \quad \text{when } \mathbb{S}(\mathbf{x}') \in \mathcal{S}\mathbf{D} \text{ for all } \mathbf{x}', \tag{4.31}$$

where

$$\begin{aligned} \mathcal{S}\mathbf{D} &= \{\mathbf{A} \in L(\mathcal{T}) \mid \mathbf{A} = \mathbf{S}\mathbf{D} \text{ for some } \mathbf{S} \in \mathcal{S}\}, \\ \mathcal{K}\mathbf{D} &= \{\mathbf{A} \in L(\mathcal{T}) \mid \mathbf{A} = \mathbf{K}\mathbf{D} \text{ for some } \mathbf{K} \in \mathcal{K}\}. \end{aligned} \tag{4.32}$$

It is not immediately clear when the assumption of the central theorem that \mathbf{L}_{λ} is bounded and coercive for all $\lambda \in [0, 1]$ is satisfied. The following theorem gives a simple condition on \mathbf{M} that guarantees this.

Theorem 4.2 *Let \mathbf{M} be such that $\mathbf{M}\mathbf{L}_0\mathbf{M} \leq \mathbf{M}$, and assume that \mathbf{L}, \mathbf{L}_0 are coercive. Then \mathbf{L}_{λ} defined at (3.7) is bounded and coercive for all $\lambda \in [0, 1]$.*

Proof The boundedness of \mathbf{L}_{λ} can be seen to be a corollary of Grabovsky’s Definition 3.17 and Lemma 3.18 [15]. Then we remark that condition $\mathbf{M}\mathbf{L}_0\mathbf{M} \leq \mathbf{M}$ implies the fact that the family of self-adjoint operators \mathbf{L}_{λ} is continuous in $\lambda \in [0, 1]$. Therefore, eigenvalues of \mathbf{L}_{λ} depend continuously on λ .

When $\lambda = 1$, or $\lambda = 0$, we have $\mathbf{L}_{\lambda} = \mathbf{L}$ and, respectively, $\mathbf{L}_{\lambda} = \mathbf{L}_0$ which are bounded and coercive by hypothesis.

Thus, if there exists a λ for which \mathbf{L}_{λ} is not coercive, then there exists $\underline{\lambda} \in (0, 1)$ so that at least one of the eigenvalues of $\mathbf{L}_{\underline{\lambda}}$ is zero, which in turn will imply that $\mathbf{L}_{\underline{\lambda}}$ is a singular

matrix. Hence, to prove coercivity for the family L_λ for all $\lambda \in (0, 1)$ it is sufficient to show that L_λ is invertible for all $\lambda \in (0, 1)$.

In this regard, if we consider

$$L'_\lambda = L_0^{-1/2} L_\lambda L_0^{-1/2}, \quad L' = L_0^{-1/2} L L_0^{-1/2}, \quad M' = L_0^{1/2} M L_0^{1/2}, \tag{4.33}$$

(3.7) becomes

$$L'_\lambda = I + \lambda [I + (1 - \lambda)(L' - I)M']^{-1}(L' - I), \tag{4.34}$$

where the condition $ML_0M \leq M$ is rewritten for M' as $(M')^2 \leq M'$ and we also have $L' > 0$. Then invertibility of L_λ is equivalent to invertibility of L'_λ which in turn is equivalent to the invertibility of

$$F = [I + (1 - \lambda)(L' - I)M']L'_\lambda = (1 - \lambda)(I - M') + L'(\lambda I + (1 - \lambda)M'). \tag{4.35}$$

The eigenvalues of M' are between 0 and 1 and therefore the eigenvalues of $\lambda I + (1 - \lambda)M'$ will always be between λ and 1. Thus, invertibility of F is equivalent to the invertibility of

$$F' = D + L', \quad D = (1 - \lambda)(I - M')(\lambda I + (1 - \lambda)M')^{-1}. \tag{4.36}$$

The eigenvalues of the self-adjoint operator D are clearly nonnegative, and therefore, the operator $D + L'$ is positive definite and hence invertible. \square

Remark 4.3 Here we show that the condition (3.8) simplifies in the case where for $\mathbf{k} \neq 0$, $\Gamma(\mathbf{k})$ only depends on $\mathbf{k}/|\mathbf{k}|$, and that M can be eliminated from the condition. By subtracting the conditions implied by (3.8) that

$$\mathcal{K}(M - \Gamma(\mathbf{n}))\mathcal{K} \subset \mathcal{K}, \quad \mathcal{K}(M - \Gamma(\mathbf{m}))\mathcal{K} \subset \mathcal{K}, \tag{4.37}$$

we get that $\mathcal{K}(\Gamma(\mathbf{n}) - \Gamma(\mathbf{m}))\mathcal{K} \subset \mathcal{K}$ for all unit vectors \mathbf{m} and \mathbf{n} . Hence, if (4.37) holds it will still hold if M is replaced by $\langle \Gamma(\mathbf{n}) \rangle_{\mathbf{n}}$ where the angular brackets $\langle \cdot \rangle_{\mathbf{n}}$ denote a possibly weighted average over the sphere $|\mathbf{n}| = 1$. The formula (2.6) for $\Gamma(\mathbf{k})$ implies $0 \leq \Gamma(\mathbf{n}) \leq L_0^{-1}$. So with the choice $M = \langle \Gamma(\mathbf{n}) \rangle_{\mathbf{n}}$ we have that $M \leq L_0^{-1}$ which as $M \geq 0$ then implies the condition that $ML_0M \leq M$. Other choices of M may be useful too, as given \mathcal{K} , both M and L_0 determine the manifold $\mathcal{M} = W_M^{-1}(\mathcal{K}_0)$, where \mathcal{K}_0 consists of all self-adjoint maps in \mathcal{K} .

5 Exact identities satisfied by the Green's function

Consider a point \mathbf{x}^0 and take $\mathbf{h}(\mathbf{x})$ to be proportional to δ , which as conveyed in Remark 1.1 denotes a smooth approximation of a Dirac delta function localized at $\mathbf{x} = \mathbf{x}^0$:

$$\mathbf{h}(\mathbf{x}) = \mathbf{h}^0 \delta(\mathbf{x} - \mathbf{x}^0), \quad \text{with } \mathbf{h}^0 = -(\mathbf{L}(\mathbf{x}^0) - L_0)\mathbf{s}^0, \tag{5.1}$$

where the amplitude $\mathbf{s}^0 \in \mathcal{T}$ is prescribed. We also recall that \mathbf{G} denotes in this paper an approximate Green's function (see Remark 1.1), and here and in the next two sections, we assume that $\mathbf{L}(\mathbf{x})$ is smooth enough.

Then, with appropriate decay conditions at infinity imposed so that the Green's function (fundamental solution) exists and is unique, (1.6) and (4.6) informally imply

$$\mathbb{P}(\mathbf{x}) = (\mathbf{L}(\mathbf{x}^0) - L_0)\mathbf{s}^0 \delta(\mathbf{x} - \mathbf{x}^0) - (\mathbf{L}(\mathbf{x}) - L_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - L_0)\mathbf{s}^0. \tag{5.2}$$

With the tensor $\mathbf{s}^0 \in \mathcal{T}$ replaced by a succession of q tensors $\mathbf{s}_1^0, \mathbf{s}_2^0, \dots, \mathbf{s}_q^0$, each in \mathcal{T} , and defining the linear map $\mathbf{S}^0 : \mathcal{T} \rightarrow \mathcal{T}$ via $\mathbf{S}^0 \mathbf{e}_i = \mathbf{s}_i^0$, we obtain

$$\begin{aligned} \mathbb{P}(\mathbf{x})\mathbf{e}_i &= (\mathbf{L}(\mathbf{x}^0) - L_0)\mathbf{s}_i^0 \delta(\mathbf{x} - \mathbf{x}^0) - (\mathbf{L}(\mathbf{x}) - L_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - L_0)\mathbf{s}_i^0 \\ &= [(\mathbf{L}(\mathbf{x}^0) - L_0)\delta(\mathbf{x} - \mathbf{x}^0) - (\mathbf{L}(\mathbf{x}) - L_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - L_0)]\mathbf{S}^0 \mathbf{e}_i, \end{aligned} \tag{5.3}$$

informally implying

$$\mathbb{P}(\mathbf{x}) = \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{S}^0, \tag{5.4}$$

with

$$\mathbf{T}(\mathbf{x}, \mathbf{x}^0) = (\mathbf{L}(\mathbf{x}^0) - \mathbf{L}_0)\delta(\mathbf{x} - \mathbf{x}^0) - (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - \mathbf{L}_0). \tag{5.5}$$

For fixed \mathbf{x} and \mathbf{x}^0 , with $\mathbf{x} \neq \mathbf{x}^0$ we can consider $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ as a map from \mathcal{T} to \mathcal{T} , and given any $\mathbf{S}^0 : \mathcal{T} \rightarrow \mathcal{T}$ we can choose sources \mathbf{s}_i^0 such that $\mathbf{S}^0 \mathbf{e}_i = \mathbf{s}_i^0$. Define $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathcal{S} = \{\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{S} \mid \mathbf{S} \in \mathcal{S}\}$ where $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{S}$ is the composition of the two maps, and \mathcal{S} , as defined at the beginning of Sect. 4, is a subspace of $L(\mathcal{T})$ such that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$. Then our theorem says that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathcal{S} \subset \mathcal{K}$ for all \mathbf{x}, \mathbf{x}^0 , with $\mathbf{x} \neq \mathbf{x}^0$. Alternatively we can view $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ as a map from $L(\mathcal{T})$ to $L(\mathcal{T})$ defined as $\mathbf{B} = \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{A}$ iff $\mathbf{b}_i = \mathbf{B}\mathbf{e}_i$ and $\mathbf{a}_i = \mathbf{A}\mathbf{e}_i$ satisfy $\mathbf{b}_i = \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{a}_i$ for $i = 1, 2, \dots, q$. Viewed in this way, $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ maps \mathcal{S} to a subset of \mathcal{K} . More generally, (4.31) implies $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ maps $\mathcal{S}\mathbf{D}$ to a subset of $\mathcal{K}\mathbf{D}$ for all nonsingular $\mathbf{D} : \mathcal{T} \rightarrow \mathcal{T}$. Also given any nonsingular \mathbf{S}_0 , we can choose \mathbf{D} so that $\mathbf{S}_0\mathbf{D} \in \mathcal{S}$ and $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ will then map this to an element of $\mathcal{K}\mathbf{D}$.

To better understand this property of $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$, consider the operator $\mathbf{T} : \mathcal{H} \rightarrow \mathcal{H}$, associated with the integral kernel $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$, given by

$$\mathbf{T} = (\mathbf{L} - \mathbf{L}_0) - (\mathbf{L} - \mathbf{L}_0)\mathbf{G}(\mathbf{L} - \mathbf{L}_0) = (\mathbf{I} - \mathbf{K}\Psi)^{-1}\mathbf{K}, \tag{5.6}$$

where the last identity follows from (4.12). Define the associated sequence of operators

$$\mathbf{T}_\lambda = \lambda(\mathbf{I} - \lambda\mathbf{K}\Psi)^{-1}\mathbf{K} = \sum_{j=0}^{\infty} \lambda^{j+1}(\mathbf{K}\Psi)^j\mathbf{K}, \tag{5.7}$$

where as \mathbf{K} and Ψ are bounded operators the operator expansion converges for small enough λ . The associated integral kernel $\mathbf{T}_\lambda(\mathbf{x}, \mathbf{x}^0)$ (regarded as a generalized function) can then be written as a series of convolutions, and the first few terms of which are given informally by

$$\begin{aligned} \mathbf{T}_\lambda(\mathbf{x}, \mathbf{x}^0) &= \lambda\delta(\mathbf{x} - \mathbf{x}^0)\mathbf{K}(\mathbf{x}^0) + \lambda^2\mathbf{K}(\mathbf{x})\widehat{\Psi}(\mathbf{x} - \mathbf{x}^0)\mathbf{K}(\mathbf{x}^0) \\ &\quad + \lambda^3 \int_{\mathbb{R}^d} \mathbf{K}(\mathbf{x})\widehat{\Psi}(\mathbf{x} - \mathbf{y}_1)\mathbf{K}(\mathbf{y}_1)\widehat{\Psi}(\mathbf{y}_1 - \mathbf{x}^0)\mathbf{K}(\mathbf{x}^0) \, d\mathbf{y}_1 \\ &\quad + \lambda^4 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbf{K}(\mathbf{x})\widehat{\Psi}(\mathbf{x} - \mathbf{y}_1)\mathbf{K}(\mathbf{y}_1)\widehat{\Psi}(\mathbf{y}_1 - \mathbf{y}_2)\mathbf{K}(\mathbf{y}_2) \\ &\quad \quad \widehat{\Psi}(\mathbf{y}_2 - \mathbf{x}^0)\mathbf{K}(\mathbf{x}^0) \, d\mathbf{y}_1 \, d\mathbf{y}_2 + \dots, \end{aligned} \tag{5.8}$$

in which $\widehat{\Psi}(\mathbf{x})$, is the Fourier transform of the operator $\Psi(\mathbf{k})$ associated with the operator Ψ . Clearly $\widehat{\Psi}(\mathbf{x})$ lies in the space spanned by the $\Psi(\mathbf{k})$, and therefore (3.8) implies that each successive term in the expansion (5.8) lies in \mathcal{K} and hence $\mathbf{T}_\lambda(\mathbf{x}, \mathbf{x}^0) \in \mathcal{K}$ for small enough λ , assuming the convergence of the series is pointwise not just in the L^2 -sense implied by the boundedness of \mathbf{K} and Ψ . As the properties of $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ must be such as to account for the Central Theorem 4.1, there presumably must be some analytic continuation or other arguments which allow us to deduce that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0) \in \mathcal{K}$ for each \mathbf{x} and \mathbf{x}^0 . We make the assumption that such arguments will be found. By definition, \mathcal{S} is a subspace of $L(\mathcal{T})$ such that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$. So as $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ takes values in \mathcal{K} we immediately see that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ maps \mathcal{S} to a subset of \mathcal{K} , as expected. It appears that the converse need not be true as an operator in $L(\mathcal{T})$ that maps \mathcal{S} to a subset of \mathcal{K} , generally need not lie in \mathcal{K} . Thus, the assertion that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0) \in \mathcal{K}$ for each \mathbf{x} and \mathbf{x}^0 appears to contain more information than

that covered by the central theorem 4.1. However, observe that if we consider \mathcal{S} to be the largest set such that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$ then \mathcal{S} becomes a unit algebra. Then, since our theorem implies $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{S} \in \mathcal{K}$ for all $\mathbf{S} \in \mathcal{S}$, by choosing $\mathbf{S} = \mathbf{I}$ we obtain $\mathbf{T}(\mathbf{x}, \mathbf{x}^0) \in \mathcal{K}$.

If \mathbf{L} is a self-adjoint operator then so too is \mathbf{G} . Indeed, supposing that $\mathbf{E}' = \mathbf{G}\mathbf{h}'$ and $\mathbf{E} = \mathbf{G}\mathbf{h}$ then (3.2) implies

$$\mathbf{J} = \mathbf{L}\mathbf{E} - \mathbf{h}, \quad \mathbf{J}' = \mathbf{L}\mathbf{E}' - \mathbf{h}' \text{ with } \mathbf{J}, \mathbf{J}' \in \mathcal{J}, \quad \mathbf{E}, \mathbf{E}' \in \mathcal{E}, \quad (5.9)$$

and using the orthogonality of \mathcal{E} and \mathcal{J} we have

$$\langle \mathbf{G}\mathbf{h}, \mathbf{h}' \rangle_{\mathcal{H}} = \langle \mathbf{E}, \mathbf{L}\mathbf{E}' - \mathbf{J}' \rangle_{\mathcal{H}} = \langle \mathbf{E}, \mathbf{L}\mathbf{E}' \rangle_{\mathcal{H}} = \langle \mathbf{L}\mathbf{E}, \mathbf{E}' \rangle_{\mathcal{H}} = \langle \mathbf{L}\mathbf{E} - \mathbf{J}, \mathbf{E}' \rangle_{\mathcal{H}} = \langle \mathbf{h}, \mathbf{G}\mathbf{h}' \rangle_{\mathcal{H}}. \quad (5.10)$$

which implies \mathbf{G} is self-adjoint. In terms of $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ this says that

$$\mathbf{T}(\mathbf{x}, \mathbf{x}^0) = \mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x}), \quad (5.11)$$

where $\mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})$ is the adjoint of $\mathbf{T}(\mathbf{x}^0, \mathbf{x})$ on the space \mathcal{T} . The extension of $\mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})$ to an operator (going by the same name) acting on $L(\mathcal{T})$ is also the adjoint, with respect to the inner product (4.20), of the extension of $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ that acts on the space $L(\mathcal{T})$. To see this, we have

$$\begin{aligned} \langle \mathbf{A}, \mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})\mathbf{B} \rangle_{L(\mathcal{T})} &= \sum_{i=1}^q \langle \mathbf{A}\mathbf{e}_i, [\mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})\mathbf{B}]\mathbf{e}_i \rangle_{\mathcal{T}} = \sum_{i=1}^q \langle \mathbf{A}\mathbf{e}_i, \mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})(\mathbf{B}\mathbf{e}_i) \rangle_{\mathcal{T}} \\ &= \sum_{i=1}^q \langle \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{A}\mathbf{e}_i, \mathbf{B}\mathbf{e}_i \rangle_{\mathcal{T}} = \langle \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{A}, \mathbf{B} \rangle_{L(\mathcal{T})}. \end{aligned} \quad (5.12)$$

Our theorem then implies $\mathbf{T}^\dagger(\mathbf{x}^0, \mathbf{x})\mathcal{S}\mathbf{D} \subset \mathcal{K}\mathbf{D}$ for all \mathbf{x}, \mathbf{x}^0 , with $\mathbf{x} \neq \mathbf{x}^0$, and for all nonsingular $\mathbf{D} : \mathcal{T} \rightarrow \mathcal{T}$. By swapping \mathbf{x} and \mathbf{x}^0 we see that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathcal{S}\mathbf{D}$ and $\mathbf{T}^\dagger(\mathbf{x}, \mathbf{x}^0)\mathcal{S}\mathbf{D}$ are both subsets of \mathcal{K} . The latter implies that for all $\mathbf{B} \in \mathcal{S}\mathbf{D}$ and $\mathbf{A} \in (\mathcal{K}\mathbf{D})^\perp$ that

$$0 = \langle \mathbf{A}, \mathbf{T}^\dagger(\mathbf{x}, \mathbf{x}^0)\mathbf{B} \rangle_{L(\mathcal{T})} = \langle \mathbf{T}(\mathbf{x}, \mathbf{x}^0)\mathbf{A}, \mathbf{B} \rangle_{L(\mathcal{T})}. \quad (5.13)$$

So we see that $\mathbf{T}(\mathbf{x}, \mathbf{x}^0)$ maps not only $\mathcal{S}\mathbf{D}$ to a subset of $\mathcal{K}\mathbf{D}$, but also $(\mathcal{K}\mathbf{D})^\perp$ to a subset of $(\mathcal{S}\mathbf{D})^\perp$, in which $(\mathcal{S}\mathbf{D})^\perp$ is the orthogonal complement of $\mathcal{S}\mathbf{D}$ in the space $L(\mathcal{T})$ with respect to the inner product (4.20).

Further insight into the relation between the operator \mathbf{T} defined by (5.6) and the Green's operator \mathbf{G} can be gained by applying the operator $(\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}$ to both sides of

$$\mathbf{J} - \mathbf{L}_0\mathbf{E} = (\mathbf{L} - \mathbf{L}_0)\mathbf{E} - \mathbf{h}, \quad \mathbf{J} \in \mathcal{J}, \quad \mathbf{E} \in \mathcal{E}, \quad (5.14)$$

where we now only require that $\mathbf{h} \in \mathcal{H}$. This gives

$$-(\mathbf{L} - \mathbf{L}_0)\mathbf{E} = (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}(\mathbf{L} - \mathbf{L}_0)\mathbf{E} - (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}\mathbf{h}, \quad (5.15)$$

and hence

$$(\mathbf{L} - \mathbf{L}_0)\mathbf{E} = [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}]^{-1}(\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}\mathbf{h}. \quad (5.16)$$

As $\mathbf{E} = \mathbf{G}\mathbf{h}$ we see that the Green's function \mathbf{G} satisfies

$$\begin{aligned} (\mathbf{L} - \mathbf{L}_0)\mathbf{G} &= [\mathbf{I} + (\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma}]^{-1}(\mathbf{L} - \mathbf{L}_0)\mathbf{\Gamma} \\ &= (\mathbf{I} - \mathbf{K}\mathbf{\Psi})^{-1}\mathbf{K}\mathbf{\Gamma} \\ &= \mathbf{T}\mathbf{\Gamma}, \end{aligned} \quad (5.17)$$

where we have made use of (4.21) (with $\lambda = 1$). Instead of (4.14), we may write

$$\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x}) - \mathbb{H}(\mathbf{x}), \text{ with } \mathbb{E} \in \mathfrak{E}, \quad \mathbb{J} \in \mathfrak{J}, \tag{5.18}$$

and we see that if $\Gamma\mathbb{H}$ takes values in \mathcal{SD} , then $(\mathbf{L}-\mathbf{L}_0)\mathbb{E}$ takes values in \mathcal{KD} . The constraint that $\Gamma\mathbb{H}$ takes values in \mathcal{SD} is of course a nonlocal constraint on $\mathbb{H}(\mathbf{x})$ and therefore not as easy to check as the constraint that $\mathbb{S}(\mathbf{x}) \in \mathcal{SD}$ (assuming $\mathbb{H} = -(\mathbf{L} - \mathbf{L}_0)\mathbb{S}$ for some $\mathbb{S} \in \mathfrak{S}$).

6 Links between Green’s functions of different physical problems

In the same way that the theory of exact relations for composites easily provides links between effective tensors, so too does our theory easily provide links between the Green’s functions of different physical problems. The treatment here adapts the theory of links for composites, given in [17] Sect. 4.3, to Green’s functions.

Consider m different physical problems, each described by an equation that can be expressed in the form (1.5):

$$\mathbf{J}^{(i)}(\mathbf{x}) = \mathbf{L}^{(i)}(\mathbf{x})\mathbf{E}^{(i)}(\mathbf{x}) - \mathbf{h}^{(i)}(\mathbf{x}), \text{ with } \mathbf{J}^{(i)} \in \mathcal{J}^{(i)}, \quad \mathbf{E}^{(i)} \in \mathcal{E}^{(i)}, \quad \mathbf{h}^{(i)} \in \mathcal{H}^{(i)}, \tag{6.1}$$

where $i = 1, 2, \dots, m$ indexes each different problem, and each field $\mathbf{J}^{(i)}(\mathbf{x})$, $\mathbf{E}^{(i)}(\mathbf{x})$ and $\mathbf{h}^{(i)}(\mathbf{x})$ takes values in a tensor space $\mathcal{T}^{(i)}$ for every $\mathbf{x} \in \mathbb{R}^d$. The projection $\Gamma_1^{(i)}$ onto $\mathcal{E}^{(i)}$ is assumed to act locally in Fourier space, i.e., if $\mathbf{E}^{(i)} = \Gamma_1^{(i)}\mathbf{A}^{(i)}$ then the Fourier components $\widehat{\mathbf{E}}^{(i)}(\mathbf{k})$ and $\widehat{\mathbf{A}}^{(i)}(\mathbf{k})$ of $\mathbf{E}^{(i)}$ and $\mathbf{A}^{(i)}$ are related via $\widehat{\mathbf{E}}^{(i)}(\mathbf{k}) = \Gamma_1^{(i)}(\mathbf{k})\widehat{\mathbf{A}}^{(i)}(\mathbf{k})$ for some operator $\Gamma_1^{(i)}(\mathbf{k})$ that projects onto a subspace $\mathcal{E}_k^{(i)} \subset \mathcal{T}$.

We can rewrite this set of equations in the equivalent form

$$\underbrace{\begin{pmatrix} \mathbf{J}^{(1)}(\mathbf{x}) \\ \mathbf{J}^{(2)}(\mathbf{x}) \\ \vdots \\ \mathbf{J}^{(m)}(\mathbf{x}) \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} \mathbf{L}^{(1)}(\mathbf{x}) & 0 & \dots & 0 \\ 0 & \mathbf{L}^{(2)}(\mathbf{x}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{L}^{(m)}(\mathbf{x}) \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \mathbf{E}^{(1)}(\mathbf{x}) \\ \mathbf{E}^{(2)}(\mathbf{x}) \\ \vdots \\ \mathbf{E}^{(m)}(\mathbf{x}) \end{pmatrix}}_{\mathbf{E}(\mathbf{x})} - \underbrace{\begin{pmatrix} \mathbf{h}^{(1)}(\mathbf{x}) \\ \mathbf{h}^{(2)}(\mathbf{x}) \\ \vdots \\ \mathbf{h}^{(m)}(\mathbf{x}) \end{pmatrix}}_{\mathbf{h}(\mathbf{x})}, \tag{6.2}$$

where $\mathbf{J}(\mathbf{x})$, $\mathbf{E}(\mathbf{x})$ and $\mathbf{h}(\mathbf{x})$ take values in $\mathcal{T} = \mathcal{T}^{(1)} \oplus \mathcal{T}^{(2)} \oplus \dots \oplus \mathcal{T}^{(m)}$, and satisfy

$$\mathbf{E} \in \mathcal{E}, \quad \mathbf{J} \in \mathcal{J}, \quad \mathbf{h} \in \mathcal{H}, \tag{6.3}$$

in which \mathcal{J} and \mathcal{E} consist of all those fields $\mathbf{J}(\mathbf{x})$ and $\mathbf{E}(\mathbf{x})$, respectively, taking the form indicated in (6.2) with component fields $\mathbf{J}^{(i)} \in \mathcal{J}^{(i)}$ and $\mathbf{E}^{(i)} \in \mathcal{E}^{(i)}$, for $i = 1, 2, \dots, m$, and \mathcal{H} is defined by $\mathcal{H} = \mathcal{E} \oplus \mathcal{J}$. The Green’s function for the system of uncoupled Eqs. (6.2) and (6.3) takes the form

$$\mathbf{G}(\mathbf{x}, \mathbf{x}^0) = \begin{pmatrix} \mathbf{G}^{(1)}(\mathbf{x}, \mathbf{x}^0) & 0 & \dots & 0 \\ 0 & \mathbf{G}^{(2)}(\mathbf{x}, \mathbf{x}^0) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{G}^{(m)}(\mathbf{x}, \mathbf{x}^0) \end{pmatrix}, \tag{6.4}$$

in which $\mathbf{G}^{(i)}(\mathbf{x}, \mathbf{x}^0)$ denotes the Green’s function for the “ i -th” problem. We introduce a constant reference tensor \mathbf{L}_0 and a constant tensor \mathbf{M} that are both assumed to be block diagonal:

$$\mathbf{L}_0 = \begin{pmatrix} \mathbf{L}_0^{(1)} & 0 & \dots & 0 \\ 0 & \mathbf{L}_0^{(2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{L}_0^{(m)} \end{pmatrix}, \quad \mathbf{M} = \begin{pmatrix} \mathbf{M}^{(1)} & 0 & \dots & 0 \\ 0 & \mathbf{M}^{(2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{M}^{(m)} \end{pmatrix}. \tag{6.5}$$

Then associated with the decoupled system (6.2) and (6.4) is an operator

$$\Psi(\mathbf{k}) = \underbrace{\begin{pmatrix} \Psi^{(1)}(\mathbf{k}) & 0 & \dots & 0 \\ 0 & \Psi^{(2)}(\mathbf{k}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \Psi^{(m)}(\mathbf{k}) \end{pmatrix}}_{\text{Block Diagonal}} = \underbrace{\begin{pmatrix} \mathbf{M}^{(1)} - \Gamma^{(1)}(\mathbf{k}) & 0 & \dots & 0 \\ 0 & \mathbf{M}^{(2)} - \Gamma^{(2)}(\mathbf{k}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{M}^{(m)} - \Gamma^{(m)}(\mathbf{k}) \end{pmatrix}}_{\text{Block Diagonal}}, \tag{6.6}$$

in which

$$\Gamma^{(i)}(\mathbf{k}) = [\Gamma_1^{(i)}(\mathbf{k})\mathbf{L}_0\Gamma_1^{(i)}(\mathbf{k})]^{-1}\Gamma_1^{(i)}(\mathbf{k}), \tag{6.7}$$

where $\Gamma_1^{(i)}(\mathbf{k})$ is the projection onto $\mathcal{E}_k^{(i)}$ and the operator inverse is taken on $\mathcal{E}_k^{(i)}$.

As before, we search for subspaces $\mathcal{K} \subset L(T)$ having the property (3.8) and associated subspaces $\mathcal{S} \subset L(T)$ that are subspaces having the property that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$. Due to the special algebraic structure of the problem, the search for such subspaces simplifies: see Sect. 4.3 of [17] and pages 5–3 to 5–11 of [15]. The exact relation then implies that

$$(\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - \mathbf{L}_0) + \mathbf{L}_0 - \mathbf{L} \in \mathcal{K} \quad \text{for all } \mathbf{x}, \mathbf{x}^0, \text{ with } \mathbf{x} \neq \mathbf{x}^0, \tag{6.8}$$

and if $\mathbf{L}^{(i)}(\mathbf{x})$ is self-adjoint for all i we have additionally that for all nonsingular $\mathbf{D} \rightarrow \mathbf{D}$,

$$((\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbf{G}(\mathbf{x}, \mathbf{x}^0)(\mathbf{L}(\mathbf{x}^0) - \mathbf{L}_0) + \mathbf{L}_0 - \mathbf{L})(\mathcal{K}\mathbf{D})^\perp \subset (\mathcal{S}\mathbf{D})^\perp \quad \text{for all } \mathbf{x}, \mathbf{x}^0, \text{ with } \mathbf{x} \neq \mathbf{x}^0, \tag{6.9}$$

in which $(\mathcal{K}\mathbf{D})^\perp$ and $(\mathcal{S}\mathbf{D})^\perp$ are the orthogonal complements of $\mathcal{K}\mathbf{D}$ and $\mathcal{S}\mathbf{D}$ on the space $L(T)$, with respect to the inner product (4.20).

For these exact relations to generate a link between the Green’s functions of the different problems, it is necessary that \mathcal{K} not be separable. It is separable if after some reordering of the indices $i = 1, 2, \dots, m$ labeling the m problems there exists a subdivision of the problems such that the exact relation decouples. In other words, there exists a p , $1 < p < m$, such that there are subspaces $\mathcal{K}_1 \subset L(\mathcal{T}_1)$, and $\mathcal{K}_2 \subset L(\mathcal{T}_2)$, where

$$\mathcal{T}_1 = \mathcal{T}^{(1)} \oplus \mathcal{T}^{(2)} \oplus \dots \oplus \mathcal{T}^{(p)}, \quad \mathcal{T}_2 = \mathcal{T}^{(p+1)} \oplus \mathcal{T}^{(2)} \oplus \dots \oplus \mathcal{T}^{(m)}, \tag{6.10}$$

with \mathcal{K}_1 and \mathcal{K}_2 each having the algebraic properties of an exact relation, and any block diagonal tensor

$$\mathbf{B} = \begin{pmatrix} \mathbf{B}^{(1)} & 0 & \dots & 0 \\ 0 & \mathbf{B}^{(2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{B}^{(m)} \end{pmatrix} \tag{6.11}$$

is in \mathcal{K} if and only if

$$\mathbf{B}_1 = \begin{pmatrix} \mathbf{B}^{(1)} & 0 & \dots & 0 \\ 0 & \mathbf{B}^{(2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{B}^{(p)} \end{pmatrix} \in \mathcal{K}_1 \quad \text{and} \quad \mathbf{B}_2 = \begin{pmatrix} \mathbf{B}^{(p+1)} & 0 & \dots & 0 \\ 0 & \mathbf{B}^{(p+2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{B}^{(m)} \end{pmatrix} \in \mathcal{K}_2. \tag{6.12}$$

When we say \mathcal{K}_1 and \mathcal{K}_2 have the properties of an exact relation, we specifically mean that for all $\mathbf{k} \in \mathbb{R}^d$ one has

$$\mathbf{A}_1 \Psi_j(\mathbf{k}) \mathbf{A}_2 \in \mathcal{K}_j, \quad \text{for all } \mathbf{A}_1, \mathbf{A}_2 \in \mathcal{K}_j, \quad j = 1, 2, \tag{6.13}$$

in which

$$\Psi_1(\mathbf{k}) = \begin{pmatrix} \Psi^{(1)}(\mathbf{k}) & 0 & \dots & 0 \\ 0 & \Psi^{(2)}(\mathbf{k}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \Psi^{(p)}(\mathbf{k}) \end{pmatrix} \text{ and}$$

$$\Psi_2(\mathbf{k}) = \begin{pmatrix} \Psi^{(p+1)}(\mathbf{k}) & 0 & \dots & 0 \\ 0 & \Psi^{(p+2)}(\mathbf{k}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \Psi^{(m)}(\mathbf{k}) \end{pmatrix}. \tag{6.14}$$

7 Exact identities satisfied by the DtN map and boundary field equalities: a generalization of conservation laws

This section generalizes the ideas developed in [48], where it was shown how Hill’s exact relation in the theory of composites, could be used to derive exact identities satisfied by the “Dirichlet-to-Neumann map” of a body Ω containing two elastically isotropic materials with the same shear modulus: in particular, these identities allow one to exactly deduce the volume fractions occupied by the phases from boundary measurements. The key idea was to apply (nonlocal) boundary conditions on the boundary tractions and displacements on the boundary $\partial\Omega$ of Ω in such a way that they mimic the body placed in an appropriate infinite medium with appropriate sources outside. The exact relations satisfied by the fields in the latter problem imply that the fields inside Ω satisfy these exact relations too, and this in turn allows one to obtain additional information about the boundary fields: the boundary field equalities.

We recall the conventions of Remark 1.1, and our assumption made in the beginning of Sect. 5 that $\mathbf{L}(\mathbf{x})$ is smooth enough. We also mention that here and in the next section we assume Ω is sufficiently smooth and we restrict attention to those equations (having the required canonical form) for which the response of a body filled by inhomogeneous material and devoid of sources inside is governed by a “Dirichlet-to-Neumann map” (DtN map) Λ_Ω . Symbolically we may write

$$\partial \mathbf{J} = \Lambda_\Omega(\partial \mathbf{E}), \tag{7.1}$$

where $\partial \mathbf{E}$ informally denotes the boundary information associated with the field \mathbf{E} , and $\partial \mathbf{J}$ informally denotes the boundary information associated with the field \mathbf{J} . In the context of the primary Eq. (1.2), $\partial \mathbf{E}$ represents the value of the potential field $\mathbf{u}(\mathbf{x})$ at the boundary $\partial\Omega$ of Ω and $\partial \mathbf{J}$ represents the value of the flux vector field $\mathbf{n} \cdot \mathbf{J}(\mathbf{x})$ at the boundary $\partial\Omega$, where \mathbf{n} is the outward normal to the surface $\partial\Omega$ (assumed smooth). (Equivalently, for the primary equations, $\partial \mathbf{E}$ can be taken as the tangential values of $\mathbf{E}(\mathbf{x})$ since integrating these over the surface yields the boundary values of $\mathbf{u}(\mathbf{x})$, up to an additive constant vector). The Appendix gives further examples of boundary fields $\partial \mathbf{J}$ and $\partial \mathbf{E}$ that are associated with various physical equations, in particular quasistatic equations. Beyond (1.2) and the equations in the Appendix we avoid giving a precise definition of the fields $\partial \mathbf{E}$ and $\partial \mathbf{J}$.

Specifying $\partial\mathbf{E}$ determines a unique \mathbf{E} (and hence \mathbf{J}) that solve

$$\mathbf{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{E}(\mathbf{x}), \quad \mathbf{J} \in \overline{\mathcal{J}}_\Omega, \quad \mathbf{E} \in \overline{\mathcal{E}}_\Omega. \tag{7.2}$$

where $\overline{\mathcal{J}}_\Omega$ and $\overline{\mathcal{E}}_\Omega$ are the closures of the spaces \mathcal{J}_Ω and \mathcal{E}_Ω comprised of those fields $\mathbf{J}(\mathbf{x})$ and $\mathbf{E}(\mathbf{x})$ defined in Ω which can be extended outside Ω in such a way that with their extensions they lie in \mathcal{J} and \mathcal{E} , respectively. Note that $\overline{\mathcal{J}}_\Omega$ and $\overline{\mathcal{E}}_\Omega$ are not orthogonal, nor even nonintersecting. Associated with $\mathbf{J}(\mathbf{x})$ is the boundary field $\partial\mathbf{J}$. As it depends linearly on $\partial\mathbf{E}$, this linear relation defines the DtN map (7.1).

We make a side remark that, as shown in [35], Chap. 3 (7.2) can be reformulated as an equation in the abstract theory of composites,

$$\underbrace{\mathbf{L}_0^{-1/2}\mathbf{J}(\mathbf{x})}_{\tilde{\mathbf{J}}} = \underbrace{[\mathbf{L}_0^{-1/2}\mathbf{L}(\mathbf{x})\mathbf{L}_0^{-1/2}]}_{\tilde{\mathbf{L}}}\underbrace{\mathbf{L}_0^{1/2}\mathbf{E}(\mathbf{x})}_{\tilde{\mathbf{E}}}, \quad \tilde{\mathbf{J}} \in \tilde{\mathcal{U}} \oplus \tilde{\mathcal{E}} = \mathbf{L}_0^{-1/2}\overline{\mathcal{J}}_\Omega, \quad \tilde{\mathbf{E}} \in \tilde{\mathcal{U}} \oplus \tilde{\mathcal{E}} = \mathbf{L}_0^{1/2}\overline{\mathcal{E}}_\Omega, \tag{7.3}$$

where \mathbf{L}_0 can be any positive definite self-adjoint tensor (the “reference tensor”), and

$$\tilde{\mathcal{U}} \equiv (\mathbf{L}_0^{1/2}\overline{\mathcal{E}}_\Omega) \cap (\mathbf{L}_0^{-1/2}\overline{\mathcal{J}}_\Omega), \quad \tilde{\mathcal{J}} = \mathbf{L}_0^{-1/2}\mathcal{J}_\Omega^0, \quad \tilde{\mathcal{E}} = \mathbf{L}_0^{-1/2}\mathcal{E}_\Omega^0, \tag{7.4}$$

in which \mathcal{J}_Ω^0 and \mathcal{E}_Ω^0 consist of those fields in \mathcal{J} and \mathcal{E} , respectively, that vanish outside Ω . The DtN map is then associated with the “effective operator” \mathbf{L}_* mapping $\tilde{\mathcal{U}}$ to itself. Fields in $\tilde{\mathcal{U}}$ can be uniquely characterized either by the associated value of $\partial\mathbf{E}$ or by the associated value of $\partial\mathbf{J}$: given $\partial\mathbf{E}$ there is a map Ψ to a field $\tilde{\mathbf{U}} = \Psi(\partial\mathbf{E})$ in $\tilde{\mathcal{U}}$, and given $\mathbf{L}_*\tilde{\mathbf{U}} \in \tilde{\mathcal{U}}$ there is map Φ to $\partial\mathbf{J} = \Phi\mathbf{L}_*\tilde{\mathbf{U}} = \Phi\mathbf{L}_*\Psi(\partial\mathbf{E})$. This provides the connection between \mathbf{L}_* and the DtN map: $\mathbf{\Lambda}_\Omega = \Phi\mathbf{L}_*\Psi$. A similar reformulation of (7.2) as a problem in the abstract theory of composites was made independently by Grabovsky (see [15], Chap. 2) who takes $\mathbf{L}_0 = \mathbf{I}$ and refers to fields in \mathcal{U} as “harmonic functions” (as they are indeed harmonic functions in the conductivity problem). We thank Yury Grabovsky for pointing out the need for replacing \mathcal{J}_Ω and \mathcal{E}_Ω by their closures $\overline{\mathcal{J}}_\Omega$ and $\overline{\mathcal{E}}_\Omega$ in (7.2), unless, of course, they are already closed.

Boundary field equalities may be viewed as exact identities satisfied by the DtN map that are independent of the precise microstructure inside the body. For the boundary field equalities derived here we need only assume that $\mathbf{L}(\mathbf{x}) \in \mathcal{M}$ and satisfies the coercivity condition (3.1) for all $\mathbf{x} \in \Omega$.

Specifying $\partial\mathbf{E}$ is one of many possible boundary conditions that uniquely determine the fields inside Ω . Another frequently used one is specifying $\partial\mathbf{J}$. A different sort that we use here is providing some type of mixed nonlocal boundary condition that mimics surrounding the body by infinite homogeneous medium with tensor \mathbf{L}_1 with appropriate sources placed outside the body. Sources outside the body can be considered as a superposition of localized delta function sources. So let us define $\Omega^C \equiv \mathbb{R}^d \setminus \Omega$ and consider a single delta function source at $\mathbf{x}_0 \in \Omega^C$.

Then in Ω^C the equations informally take the form

$$\mathbf{J}(\mathbf{x}) = \mathbf{L}_1\mathbf{E}(\mathbf{x}) + (\mathbf{L}_1 - \mathbf{L}_0)\mathbf{s}^0\delta(\mathbf{x} - \mathbf{x}_0), \quad \mathbf{J} \in \mathcal{J}_{\Omega^C}, \quad \mathbf{E} \in \mathcal{E}_{\Omega^C}, \tag{7.5}$$

where \mathcal{J}_{Ω^C} and \mathcal{E}_{Ω^C} are comprised of those fields $\mathbf{J}(\mathbf{x})$ and $\mathbf{E}(\mathbf{x})$ defined outside Ω which can be extended inside Ω in such a way that with their extensions they lie in \mathcal{J} and \mathcal{E} , respectively. Now one can easily (numerically if not analytically) solve for the problem of a point source in an infinite homogeneous medium having moduli \mathbf{L}_1 :

$$\mathbf{J}^0(\mathbf{x}) = \mathbf{L}_1\mathbf{E}^0(\mathbf{x}) + (\mathbf{L}_1 - \mathbf{L}_0)\mathbf{s}^0\delta(\mathbf{x} - \mathbf{x}_0), \quad \mathbf{J}^0 \in \mathcal{J}, \quad \mathbf{E}^0 \in \mathcal{E}. \tag{7.6}$$

Alternatively, the analysis of this section will go through if we replace the delta function with any square-integrable source that is compactly supported outside Ω , say with $\mathbf{s}^0\delta(\mathbf{x} - \mathbf{x}_0)$ replaced by $\mathbf{s}^0r(\mathbf{x})$ where the scalar-valued function $r(\mathbf{x})$ is square integrable and $r(\mathbf{x}) = 0$ inside Ω .

Subtracting (7.6) from (7.5) gives informally

$$\tilde{\mathbf{J}}(\mathbf{x}) = \mathbf{L}_1\tilde{\mathbf{E}}(\mathbf{x}), \quad \tilde{\mathbf{J}} \in \mathcal{J}_{\Omega^c}, \quad \tilde{\mathbf{E}} \in \mathcal{E}_{\Omega^c}, \quad \tilde{\mathbf{J}}(\mathbf{x}) = \mathbf{J}(\mathbf{x}) - \mathbf{J}^0(\mathbf{x}), \quad \tilde{\mathbf{E}}(\mathbf{x}) = \mathbf{E}(\mathbf{x}) - \mathbf{E}^0(\mathbf{x}). \tag{7.7}$$

The boundary information $\partial\tilde{\mathbf{J}}$ and $\partial\tilde{\mathbf{E}}$ associated with the fields $\tilde{\mathbf{J}}$ and $\tilde{\mathbf{E}}$ are linked by the exterior DtN map $\Lambda_{\Omega^c}: \Lambda_{\Omega^c}(\partial\tilde{\mathbf{E}}) = \partial\tilde{\mathbf{J}}$. As the material outside Ω is homogeneous, Λ_{Ω^c} is in principle computable and will be assumed known. Since the fields outside and inside the body must be compatible, i.e., share the same boundary information $\partial\mathbf{J}$ and $\partial\mathbf{E}$, we see that $\partial\mathbf{E}$ and $\partial\mathbf{J}$ must be such that informally

$$\partial\mathbf{J} - \partial\mathbf{J}^0 = \Lambda_{\Omega^c}(\partial\mathbf{E} - \partial\mathbf{E}^0), \quad \text{i.e., } \partial\mathbf{J} - \Lambda_{\Omega^c}(\partial\mathbf{E}) = \partial\mathbf{J}^0 - \Lambda_{\Omega^c}(\partial\mathbf{E}^0), \tag{7.8}$$

in which $\partial\mathbf{J}^0$ and $\partial\mathbf{E}^0$ denote the boundary information associated with \mathbf{J}^0 and \mathbf{E}^0 . Some care needs to be taken. For example in the conductivity problem in which $\mathbf{J}(\mathbf{x})$ is the current, $\partial\mathbf{J}$ represents the flux $\mathbf{J} \cdot \mathbf{n}$ where \mathbf{n} is the outward normal to Ω . So in defining Λ_{Ω^c} it is important that it maps to the flux $\mathbf{J} \cdot \mathbf{n}$ where again \mathbf{n} is the outward normal to Ω , not the outward normal to Ω^c .

Equation (7.8) provides the needed boundary conditions that constrain $\partial\mathbf{J}$ and $\partial\mathbf{E}$. They are not so pleasant as they are nonlocal and involve Λ_{Ω^c} . Instead of specifying $\partial\mathbf{E}$ or $\partial\mathbf{J}$, one specifies $\partial\mathbf{J} - \Lambda_{\Omega^c}(\partial\mathbf{E})$. Supposing that the DtN map Λ_{Ω} entering (7.1) has been experimentally measured, or numerically calculated, then (7.8) provides the explicit equation

$$\Lambda_{\Omega}(\partial\mathbf{E}) - \Lambda_{\Omega^c}(\partial\mathbf{E}) = \partial\mathbf{J}^0 - \Lambda_{\Omega^c}(\partial\mathbf{E}^0) \tag{7.9}$$

that can be solved for $\partial\mathbf{E}$. The question arises as to whether a solution exists, and if so, is it unique? However, it is exactly the same as solving the governing equations (1.5) for the body Ω surrounded by homogeneous medium with tensor \mathbf{L}_1 and with the source term $\mathbf{h}(\mathbf{x}) = (\mathbf{L}_1 - \mathbf{L}_0)\mathbf{s}^0\delta(\mathbf{x} - \mathbf{x}_0)$, so uniqueness of \mathbf{E} (and hence $\partial\mathbf{E}$ and $\partial\mathbf{J}$) is assured.

Now instead of considering a single experiment, one can consider q experiments with \mathbf{s}^0 replaced by $\mathbf{s}_1^0, \mathbf{s}_2^0, \dots, \mathbf{s}_q^0$, each source remaining at \mathbf{x}_0 . The fields $\mathbf{E}(\mathbf{x})$ and $\mathbf{J}(\mathbf{x})$ are then replaced by $\mathbf{E}_i(\mathbf{x})$ and $\mathbf{J}_i(\mathbf{x})$. Let us define, as previously,

$$\mathbf{S}^0\mathbf{e}_i = \mathbf{s}_i^0, \quad \mathbb{E}(\mathbf{x})\mathbf{e}_i = \mathbf{E}_i(\mathbf{x}), \quad \mathbb{J}(\mathbf{x})\mathbf{e}_i = \mathbf{J}_i(\mathbf{x}), \quad \mathbb{P}(\mathbf{x}) = \mathbb{J}(\mathbf{x}) - \mathbf{L}_0\mathbb{E}(\mathbf{x}), \tag{7.10}$$

and let $\partial\mathbb{E}$ and $\partial\mathbb{J}$ represent the boundary information associated with $\mathbb{E}(\mathbf{x})$ and $\mathbb{J}(\mathbf{x})$, respectively. If $\mathbf{s}_1^0, \mathbf{s}_2^0, \dots, \mathbf{s}_q^0$ are chosen so that $\mathbf{S}^0 \in \mathcal{SD}$ for some nonsingular $\mathbf{D}: \mathcal{T} \rightarrow \mathcal{T}$ then our theorem implies that $\mathbb{P}(\mathbf{x}) \in \mathcal{KD}$ for all $\mathbf{x} \in \Omega$. If \mathcal{S} contains a nonsingular element \mathbf{S}^1 then there is no restriction on \mathbf{S}^0 as we can choose $\mathbf{D} = (\mathbf{S}^1)^{-1}\mathbf{S}^0$. This then constrains $\mathbb{J}(\mathbf{x}) - \mathbf{L}_0\mathbb{E}(\mathbf{x})$ to lie in \mathcal{KD} for all $\mathbf{x} \in \Omega$. This should then naturally provide constraints on the boundary information $\partial\mathbb{E}$ and $\partial\mathbb{J}$, thus yielding exact identities satisfied by the DtN map. The examples in the next section demonstrate this explicitly. These exact identities must be satisfied for every choice of nonsingular $\mathbf{D}: \mathcal{T} \rightarrow \mathcal{T}$, every choice of \mathbf{x}_0 outside Ω , and for every choice of $\mathbf{S}^0 \in \mathcal{SD}$. It seems unlikely that the set of exact identities produced as \mathbf{x}_0 varies outside Ω will all be independent. Rather it is probably the case that a source at \mathbf{x}_0 outside Ω has the same effect as a set of sources around the boundary of Ω ,

implying that it suffices to take sources restricted to the boundary of Ω . (Technically, one should take sources just outside Ω and let them approach the boundary).

If $\mathbf{L}(\mathbf{x})$ is self-adjoint, one has the additional constraint that if $\mathbf{S}^0 \in (\mathcal{KD})^\perp$ then $\mathbb{J}(\mathbf{x}) - \mathbf{L}_0\mathbb{E}(\mathbf{x})$ must lie in $(\mathcal{SD})^\perp$ for all $\mathbf{x} \in \Omega$, and this then implies additional constraints on the boundary information $\partial\mathbb{E}$ and $\partial\mathbb{J}$.

Depending on the nature of the exact relations, boundary field equalities may involve the volume fraction of one phase in a body containing two phases and thus allow the volume fraction to be exactly determined (e.g., see [48]).

Note that the constitutive law inside the body Ω that $\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x})$ for some $\mathbf{L}(\mathbf{x})$ such that $W_{\mathbf{M}}(\mathbf{L}(\mathbf{x})) \in \mathcal{KD}$ for all \mathbf{x} [or more generally that $\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x}) + (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbb{S}(\mathbf{x})$ inside Ω for some $\mathbf{L}(\mathbf{x})$ with $W_{\mathbf{M}}(\mathbf{L}(\mathbf{x})) \in \mathcal{K}$ and for some $\mathbb{S}(\mathbf{x}) \in \mathcal{S}$] can just be viewed as a nonlinear constraint on the fields \mathbb{J} and \mathbb{E} inside Ω . This constraint, coupled with the appropriate boundary conditions, allows one to deduce the boundary field equalities that constrain the boundary information $\partial\mathbb{E}$ and $\partial\mathbb{J}$. From this perspective our analysis immediately applies to a large range of nonlinear problems. This is developed further in the next section.

8 A alternative view of some boundary field equalities

As an interesting illustration of boundary field inequalities, expressed in a different form, let us focus on Eq. (1.2) with initially $h_{i\alpha}(\mathbf{x}) = 0$ within Ω . We will concentrate on cases $d = 2$ and $d = 3$ since these are of greatest practical importance. First, starting in the case $d = 2$, and defining new fields, $\tilde{J}_{j\beta}(\mathbf{x})$, with components given by

$$\tilde{J}_{1\beta}(\mathbf{x}) = -E_{2\beta}(\mathbf{x}), \quad \tilde{J}_{2\beta}(\mathbf{x}) = E_{1\beta}(\mathbf{x}), \quad (8.1)$$

Eq. (1.2) can be rewritten as

$$\mathbf{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathcal{R}^\perp\tilde{\mathbf{J}}(\mathbf{x}), \quad \nabla \cdot \mathbf{J} = 0, \quad \nabla \cdot \tilde{\mathbf{J}} = 0, \quad (8.2)$$

with \mathcal{R}^\perp being a 4-index tensor with elements

$$\{\mathcal{R}^\perp\}_{i\beta j\gamma} = \delta_{\beta\gamma}\{\mathbf{R}^\perp\}_{ij}, \quad \mathbf{R}^\perp = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (8.3)$$

where \mathbf{R}^\perp is the matrix for a 90° rotation. Thus, the action of \mathcal{R}^\perp on $\tilde{\mathbf{J}}(\mathbf{x})$ is to rotate its columns by 90° , each column being a divergence-free field, to give $\mathbf{E}(\mathbf{x})$, each column then being a curl-free field. We can also consider $2m$ solutions to (8.2) labeled by an index pair $k\nu$, $k = 1, 2$, $\nu = 1, 2, \dots, m$, with fields $\mathbf{J}^{k\nu}(\mathbf{x})$ and $\tilde{\mathbf{J}}^{k\nu}(\mathbf{x})$ each satisfying (8.2), so that the resulting set of equations can then be rewritten as

$$\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathcal{R}^\perp\tilde{\mathbb{J}}(\mathbf{x}), \quad \nabla \cdot \mathbb{J} = 0, \quad \nabla \cdot \tilde{\mathbb{J}}(\mathbf{x}) = 0, \quad (8.4)$$

where $\mathbb{J}(\mathbf{x})$ and $\tilde{\mathbb{J}}(\mathbf{x})$ are four-index fields with elements

$$\mathbb{J}_{i\alpha k\nu}(\mathbf{x}) = J_{i\alpha}^{k\nu}(\mathbf{x}), \quad \tilde{\mathbb{J}}_{i\alpha k\nu}(\mathbf{x}) = \tilde{J}_{i\alpha}^{k\nu}(\mathbf{x}), \quad (8.5)$$

and the divergence acts on the first index of these fields. The constitutive law in (8.4), the boundedness and coercivity of \mathbf{L} , and the fact that $\mathbf{L}(\mathbf{x})$ is constrained to take values in \mathcal{M} can be replaced by the constraint that $\mathbf{Q}(\mathbf{x})$, defined to have elements

$$Q_{i\alpha k\nu 1}(\mathbf{x}) = \mathbb{J}_{i\alpha k\nu}(\mathbf{x}), \quad Q_{i\alpha k\nu 2}(\mathbf{x}) = \tilde{\mathbb{J}}_{i\alpha k\nu}(\mathbf{x}), \quad (8.6)$$

takes values in a set \mathcal{B} , where a 5-index tensor \mathbf{Q} is defined to lie in \mathcal{B} if and only if there exists a four-index tensor $\mathbf{L} \in \mathcal{M}$ satisfying the boundedness and coercivity constraint that $\beta_0\mathbf{I} \geq \mathbf{L} \geq \alpha_0\mathbf{I} > 0$, and

$$Q_{i\alpha kv1} = L_{i\alpha j\beta} R_{j\beta\ell\gamma}^\perp Q_{\ell\gamma kv2}, \tag{8.7}$$

where (following the Einstein summation convention) sums over repeated indices are assumed. If the manifold \mathcal{M} has dimension m_0 , then \mathcal{B} will be a r -dimensional object (in a $8m^2$ -dimensional space) with $r = 4m^2 + m_0$.

At points \mathbf{x} where $\mathbb{J}(\mathbf{x})$ is nonsingular this constraint that $\mathbf{Q}(\mathbf{x}) \in \mathcal{B}$ is equivalent to requiring that the fields satisfy the clearly nonlinear constraint,

$$\mathbf{L}(\mathbf{x}) \equiv -\mathbb{J}(\mathbf{x})[\tilde{\mathbb{J}}(\mathbf{x})]^{-1}\mathcal{R}^\perp \in \mathcal{M}, \tag{8.8}$$

and that $\mathbf{L}(\mathbf{x})$ thus defined satisfies the boundedness and coercivity condition (3.1). We can of course lump all the last four indices of $\mathbf{Q}(\mathbf{x})$ into a single index γ taking values from 1 to $s = 4m^2$, and then regard $\mathbf{Q}(\mathbf{x})$ as a $2 \times s$ matrix-valued field satisfying $\nabla \cdot \mathbf{Q} = 0$, i.e., the columns of \mathbf{Q} are then divergence-free vector fields. The mapping of the last four indices onto γ can be chosen so the first $2m^2$ columns of \mathbf{Q} (are associated with \mathbb{J} while the last $2m^2$ columns of \mathbf{Q} are associated with $\tilde{\mathbb{J}}$. We let \mathbf{Q}_1 and \mathbf{Q}_2 denote the $2 \times 2m^2$ -matrix-valued fields formed by the first and last set of $2m^2$ columns, so these are associated with \mathbb{J} and $\tilde{\mathbb{J}}$, respectively. Associated with $\mathbf{Q}(\mathbf{x})$, $\mathbf{Q}_1(\mathbf{x})$, and $\mathbf{Q}_2(\mathbf{x})$ are then also the fluxes $\mathbf{q}(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}(\mathbf{x})$, $\mathbf{q}_1(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}_1(\mathbf{x})$, $\mathbf{q}_2(\mathbf{x}) = \mathbf{n}(\mathbf{x}) \cdot \mathbf{Q}_2(\mathbf{x})$ at the boundary $\partial\Omega$, where \mathbf{n} is the outward normal. Thus, the first and last $2m^2$ elements of $\mathbf{q}(\mathbf{x})$ give $\mathbf{q}_1(\mathbf{x})$ and $\mathbf{q}_2(\mathbf{x})$, respectively.

The boundary information for Eq. (1.2) is usually taken as the values of the potential \mathbf{u} and flux $\mathbf{n} \cdot \mathbf{J}$, where \mathbf{n} is the surface normal. However, as noted in the previous section, specifying the tangential value of $\nabla \mathbf{u}$ at the surface allows one to recover \mathbf{u} (up to a trivial constant) and thus is an equivalent condition. So we see that instead of using the potential \mathbf{u} as our boundary information one can use $\mathbf{n} \cdot \tilde{\mathbf{J}}$. Then (7.8) can be interpreted as a nonlocal, linear constraint on the boundary flux \mathbf{q} . It can be rewritten as

$$\mathbf{q}_1 - \Lambda_{\Omega^c}(\mathbf{q}_2) = \mathbf{q}_1^0 - \Lambda_{\Omega^c}(\mathbf{q}_2^0), \tag{8.9}$$

where \mathbf{q}_1^0 and \mathbf{q}_2^0 are the fluxes across $\partial\Omega$ associated with solving the equations in an infinite homogeneous medium with constant tensor $\mathbf{L}_1 \in \mathcal{M}$ with a source $\mathbb{S}(\mathbf{x})$ with support outside Ω taking values in \mathcal{SD} , and with Λ_{Ω^c} being the exterior DtN map associated with this medium. Our theorem then forces $\mathbb{J}(\mathbf{x}) - \mathbf{L}_0\mathbb{E}(\mathbf{x})$ within Ω to lie in \mathcal{KD} and this constraint can alternatively be written as $\mathbf{Q}(\mathbf{x}) \in \mathcal{D}$ implying $\mathbf{q}(\mathbf{x}) \in \mathbf{n}(\mathbf{x})\mathcal{D}$. To precisely define \mathcal{D} one can return to the representation where \mathbf{Q} is a 5-index tensor. Then $\mathbf{Q} \in \mathcal{D}$ if and only if $\mathbf{Z} \in \mathcal{KD}$ where \mathbf{Z} is the four-index tensor with elements

$$Z_{i\alpha sv} = Q_{i\alpha sv1}(\mathbf{x}) - \{\mathbf{L}_0\}_{i\alpha j\beta} R_{j\beta k\gamma}^\perp Q_{k\gamma sv2}. \tag{8.10}$$

The constraint that $\mathbf{q}(\mathbf{x}) \in \mathbf{n}(\mathbf{x})\mathcal{D}$ for all $\mathbf{x} \in \partial\Omega$ is a boundary field equality.

We can relax the constraint that $\mathbb{S}(\mathbf{x})$ is zero inside Ω , and given \mathbf{D} , \mathbf{L}_0 , α_0 , β_0 redefine \mathcal{B} so that a 5-index tensor \mathbf{Q} is in \mathcal{B} if and only if there exists an $\mathbf{S} \in \mathcal{SD}$ and a $\mathbf{L} \in \mathcal{M}$ with $\beta_0\mathbf{I} \geq \mathbf{L} \geq \alpha_0\mathbf{I} > 0$, such that

$$Q_{i\alpha sv1} + \{\mathbf{L}_0\}_{i\alpha j\beta} S_{j\beta sv} = L_{i\alpha j\beta} [R_{j\beta k\gamma}^\perp Q_{k\gamma sv2} + S_{j\beta sv}], \tag{8.11}$$

where sums over repeated indices are assumed. Again our theorem implies that with the boundary conditions (8.9) we have the boundary field equality that $\mathbf{q}(\mathbf{x}) \in \mathbf{n}(\mathbf{x})\mathcal{D}$ for all $\mathbf{x} \in \partial\Omega$, where \mathcal{D} is defined as before.

Of course, $\mathbf{Q}(\mathbf{x})$ being divergence-free field in a simply connected two-dimensional region Ω is equivalent to the existence of a s -component potential $\mathbf{w}(\mathbf{x})$ such that

$$\mathbf{Q}(\mathbf{x}) = \mathbf{R}^\perp \nabla \mathbf{w}. \tag{8.12}$$

The nonlocal linear constraints on the boundary flux \mathbf{q} can then be rephrased as nonlocal linear constraints on $\mathbf{w}(\mathbf{x})$ at the boundary $\mathbf{x} \in \partial\Omega$. The result that $\mathbf{Q}(\mathbf{x}) \in \mathcal{D}$ then implies $\nabla \mathbf{w}(\mathbf{x}) \in \mathcal{C} \equiv \mathbf{R}^\perp \mathcal{D}$, and this not only constrains the tangential derivatives of the surface potential, but also implies the additional boundary field equalities (1.10).

Now consider the three-dimensional case. Then a vector field $\mathbf{e}(\mathbf{x})$ is the gradient of a potential $u(\mathbf{x})$ in a simply connected domain Ω if and only if the 3×3 antisymmetric matrix-valued field

$$\mathbf{A}(\mathbf{x}) = \begin{pmatrix} 0 & -e_3(\mathbf{x}) & e_2(\mathbf{x}) \\ e_3(\mathbf{x}) & 0 & -e_1(\mathbf{x}) \\ -e_2(\mathbf{x}) & e_1(\mathbf{x}) & 0 \end{pmatrix}, \quad (8.13)$$

is divergence free. The requirement that $\mathbf{A}(\mathbf{x})$ be antisymmetric can be viewed as constraining it to lie in a three-dimensional subspace. Knowing the boundary value of the flux $\mathbf{n}(\mathbf{x}) \cdot \mathbf{A}(\mathbf{x})$ (where $\mathbf{n}(\mathbf{x})$ is the outward normal to $\partial\Omega$) gives the tangential components of $\mathbf{e}(\mathbf{x})$, which can be integrated to give $\mathbf{u}(\mathbf{x})$ for $\mathbf{x} \in \partial\Omega$. Thus, say, with a constitutive law $\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x})$ with $\mathbb{E} = \nabla \mathbb{U}$ where $\mathbb{U}(\mathbf{x})$ is represented as a $3m^2$ -component potential, and $\mathbb{J}(\mathbf{x})$ is represented as a divergence-free $3 \times 3m^2$ matrix-valued field, we can replace the $3 \times 3m^2$ matrix-valued \mathbb{E} by a $3 \times 3 \times 3m^2$ three-index field \mathbb{A} that is antisymmetric in the first pair of indices. Then the constitutive relation with the restriction that $\mathbf{L}(\mathbf{x}) \in \mathcal{M}$, and that $\mathbf{L}(\mathbf{x})$ is bounded and coercive, can be replaced by a nonlinear restriction $\mathbf{Q}(\mathbf{x}) \in \mathcal{B}$ involving a subset \mathcal{B} (independent of \mathbf{x}) of a nonlinear manifold, and a divergence-free $3 \times s$ matrix-valued field $\mathbf{Q}(\mathbf{x})$, where $s = 12m^2$ with components comprised of the $3 \times 3m^2$ -components of $\mathbb{J}(\mathbf{x})$ and the $3 \times 9m^2$ -components of $\mathbb{A}(\mathbf{x})$. (\mathcal{B} is defined so that it ensures \mathbb{A} is antisymmetric in the first pair of indices.) The boundary information is contained in $\mathbf{q}(\mathbf{x}) = \mathbf{n} \cdot \mathbf{Q}(\mathbf{x})$ that is restricted to satisfy a nonlocal, linear, relation of the form (8.9) where the components of \mathbf{q}_1 represent the components of $\mathbf{n} \cdot \mathbb{J}$ and the components of \mathbf{q}_2 represent the components of $\mathbf{n} \cdot \mathbb{A}$, and when appropriately defined the first $3m^2$ elements of \mathbf{q} give \mathbf{q}_1 , while the remaining last $9m^2$ elements of \mathbf{q} give \mathbf{q}_2 . Our result that $\mathbb{J}(\mathbf{x}) - \mathbf{L}_0 \mathbb{E}(\mathbf{x}) \in \mathcal{SD}$ for all $\mathbf{x} \in \Omega$ then again implies $\mathbf{Q}(\mathbf{x}) \in \mathcal{D}$, for some appropriately defined subspace \mathcal{D} . We thus obtain the boundary field equality that $\mathbf{q}(\mathbf{x}) \in \mathbf{n}(\mathbf{x}) \cdot \mathcal{D}$.

Similarly, in three dimensions, a current field $\mathbf{j}(\mathbf{x})$ satisfying $\nabla \cdot \mathbf{j}(\mathbf{x}) = 0$ can be expressed in terms of the curl of some vector potential $\mathbf{g}(\mathbf{x})$, or equivalently in terms of the antisymmetric part of $\nabla \mathbf{g}$. The values of $\mathbf{n}(\mathbf{x}) \cdot \mathbf{j}(\mathbf{x})$ only provide partial information about the tangential values of $\nabla \mathbf{g}(\mathbf{x})$ at the boundary of $\partial\Omega$ and these are insufficient to determine, by integration, $\mathbf{g}(\mathbf{x})$ for $\mathbf{x} \in \partial\Omega$. However, we can think of prescribing the boundary value of $\mathbf{g}(\mathbf{x})$ for $\mathbf{x} \in \partial\Omega$, which then allows one to determine $\mathbf{n}(\mathbf{x}) \cdot \mathbf{j}(\mathbf{x})$. More generally, with say $\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x})$ where $\mathbb{E} = \nabla \mathbb{U}$ and $\mathbb{U}(\mathbf{x})$ is a $3m^2$ -component potential, the $3 \times 3m^2$ matrix-valued field $\mathbb{J}(\mathbf{x})$ satisfying $\nabla \cdot \mathbb{J} = 0$ can be replaced by the appropriate components of $\nabla \mathbb{G}(\mathbf{x})$, for some $3 \times 3m^2$ matrix-valued potential $\mathbb{G}(\mathbf{x})$. Instead of a relation like (8.9), we obtain a restriction on the boundary potentials like

$$\mathbb{U} - \mathbf{A}_{\Omega^c}(\mathbb{G}) = \mathbb{U}^0 - \mathbf{A}_{\Omega^c}(\mathbb{G}^0), \quad (8.14)$$

for some appropriately defined exterior DtN map \mathbf{A}_{Ω^c} . Thus, instead of relations involving the components of a divergence-free $3 \times 12m^2$ matrix-valued field $\mathbf{Q}(\mathbf{x})$, one can express everything in terms of $\nabla \mathbf{w}$ where $\mathbf{w}(\mathbf{x})$ is a t -component potential, where $t = 12m^2$, comprised of the $3m^2$ elements of \mathbb{U} and the $9m^2$ elements of $\mathbb{G}(\mathbf{x})$. The restriction that $\mathbf{Q}(\mathbf{x}) \in \mathcal{B}$ is then equivalent to a nonlinear local restriction $\nabla \mathbf{w} \in \mathcal{A}$. The result that

$\mathbf{Q}(\mathbf{x}) \in \mathcal{D}$, can be rewritten as $\nabla \mathbf{w} \in \mathcal{C}$ for some appropriately defined subspace \mathcal{C} , and the boundary field equality that $\mathbf{q}(\mathbf{x}) \in \mathbf{n}(\mathbf{x}) \cdot \mathcal{D}$ implies restrictions on the tangential derivatives of \mathbf{w} at the surface $\partial\Omega$, and also implies the additional boundary field equality (1.10).

Just as in two dimensions, one can redefine \mathcal{B} (and hence \mathcal{A}) to allow for a weaker restriction of the form (8.11).

9 Conjectured extension of exact relations to some nonlinear minimization problems

The canonical equations (1.5) are the Euler–Lagrange equations for the minimizer $\underline{\mathbb{E}} = \mathbb{E}$ of the following minimization problem

$$\inf_{\underline{\mathbb{E}} \in \mathcal{E}} \langle \mathbf{L}\underline{\mathbb{E}} - 2\mathbf{h}, \underline{\mathbb{E}} \rangle_{\mathcal{H}}. \tag{9.1}$$

Similarly, the extended equations (4.14) are the Euler–Lagrange equations for the minimizer $\underline{\mathbb{E}} = \mathbb{E}$ of the following minimization problem

$$\inf_{\underline{\mathbb{E}} \in \mathcal{E}} \langle \mathbf{L}\underline{\mathbb{E}} + 2(\mathbf{L} - \mathbf{L}_0)\mathbb{S}, \underline{\mathbb{E}} \rangle_{\mathfrak{H}}, \tag{9.2}$$

where the inner product on $\mathfrak{H} = \mathcal{E} \oplus \mathfrak{J}$ is defined by (3.11). Now for each point \mathbf{x} let $\mathcal{G}(\mathbf{x})$ denote a subset of the manifold \mathcal{M} that is closed under homogenization. This is guaranteed if given any \mathbf{y} -periodic function $\mathbf{L}(\mathbf{y}) \in \mathcal{G}(\mathbf{x})$ for all $\mathbf{y} \in \mathbb{R}^d$, then the associated effective tensor \mathbf{L}^* also lies in $\mathcal{G}(\mathbf{x})$. (That it suffices to consider periodic functions was established in [1,43]). We also assume that $\mathcal{G}(\mathbf{x})$ varies smoothly with \mathbf{x} . Then consider the following double minimization problem

$$\inf_{\mathbf{L}} \inf_{\underline{\mathbb{E}} \in \mathcal{E}} \langle \mathbf{L}\underline{\mathbb{E}} + 2(\mathbf{L} - \mathbf{L}_0)\mathbb{S}, \underline{\mathbb{E}} \rangle_{\mathfrak{H}}, \tag{9.3}$$

$$\mathbf{L}(\mathbf{x}) \in \mathcal{G}(\mathbf{x}), \forall \mathbf{x}$$

Following the ideas of Kohn [20], we switch the order of taking infimums to get the nonlinear minimization problem:

$$\inf_{\underline{\mathbb{E}} \in \mathcal{E}} \int_{\mathbb{R}^d} W(\mathbf{x}, \underline{\mathbb{E}}(\mathbf{x})) \, d\mathbf{x}, \tag{9.4}$$

where for all $\mathbf{x} \in \mathbb{R}^d$ and all $\mathbf{B} \in L(T)$,

$$W(\mathbf{x}, \mathbf{B}) = \inf_{\mathbf{A} \in \mathcal{G}(\mathbf{x})} \langle \mathbf{A}\mathbf{B} + 2(\mathbf{A} - \mathbf{L}_0)\mathbb{S}(\mathbf{x}), \mathbf{B} \rangle_{L(T)}. \tag{9.5}$$

One can view the nonlinear “energy function” $W(\mathbf{x}, \mathbf{B})$ as being the infimum of a continuous set of quadratic functions. If $\mathbb{S}(\mathbf{x})$ is smooth, one expects the problem (9.4) to have a smooth minimizer $\underline{\mathbb{E}} = \mathbb{E}$. Otherwise, if a minimizing sequence $\underline{\mathbb{E}}_j, j = 1, 2, \dots$ develops fine-scale oscillations, it would be indicative that any associated sequence of tensor fields $\mathbf{L}(\mathbf{x}) = \mathbf{L}_j(\mathbf{x})$ that are a minimizing sequence for (9.3) also develops fine-scale microstructure as j increases. The development of such fine-scale microstructure indicates that composite materials are needed in the construction, having some homogenized effective tensor $\mathbf{L}^*(\mathbf{x})$. However, because $\mathcal{G}(\mathbf{x})$ is closed under homogenization we should be able to just directly use materials in $\mathcal{G}(\mathbf{x})$ whose tensor matches that of the desired effective tensor $\mathbf{L}^*(\mathbf{x})$. In this way fine-scale oscillations in $\mathbf{L}_j(\mathbf{x})$ and hence $\underline{\mathbb{E}}_j(\mathbf{x})$ (with j large) can be removed.

This argument strongly suggests that there should be a minimizer of (9.4) and that minimizer $\underline{\mathbb{E}}(\mathbf{x}) = \mathbb{E}(\mathbf{x})$ should satisfy the nonlinear Euler–Lagrange equations:

$$\mathbb{J}(\mathbf{x}) = \frac{\partial W(\mathbf{x}, \mathbf{B})}{\partial \mathbf{B}} \Big|_{\mathbf{B}=\mathbb{E}(\mathbf{x})}, \quad \mathbb{J} \in \mathfrak{J}, \quad \mathbb{E} \in \mathfrak{E}. \quad (9.6)$$

Then from (9.5) it follows that there exists some $\mathbf{L}(\mathbf{x}) \in \mathcal{G}(\mathbf{x}) \subset \mathcal{M}$ such that

$$\mathbb{J}(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbb{E}(\mathbf{x}) + (\mathbf{L}(\mathbf{x}) - \mathbf{L}_0)\mathbb{S}(\mathbf{x}). \quad (9.7)$$

Indeed, a tangent plane to a minimum of a set of quadratic functions must be tangent to at least one of them. If $\mathbb{S}(\mathbf{x})$ entering (9.5) takes values in \mathcal{SD} for some nonsingular $\mathbf{D} : \mathcal{T} \rightarrow \mathcal{T}$ we conclude that

$$\mathbb{P}(\mathbf{x}) = \mathbb{J}(\mathbf{x}) - \mathbf{L}_0\mathbb{E}(\mathbf{x}) \quad (9.8)$$

necessarily takes values in \mathcal{KD} . Also if all tensors in $\mathcal{G}(\mathbf{x})$ are self-adjoint for all \mathbf{x} , then if $\mathbb{S}(\mathbf{x})$ takes values in $(\mathcal{KD})^\perp$, $\mathbb{P}(\mathbf{x})$ necessarily takes values in $(\mathcal{SD})^\perp$.

10 Conclusions

We have laid the foundations of the theory of exact relations for wide classes of linear partial differential equations, where the coefficients are position dependent and satisfy appropriate nonlinear constraints, and have derived exact relations satisfied by their Green's functions. Similar to the theory of exact relations in composites, it all boils down to finding subspaces $\mathcal{K} \subset L(\mathcal{T})$ satisfying (2.9) and associated subspaces $\mathcal{S} \subset L(\mathcal{T})$ such that $\mathbf{A}\mathcal{S} \subset \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$. The algebraic search for such subspaces \mathcal{K} or, more precisely the subspaces \mathcal{K}_0 consisting of symmetric tensors in \mathcal{K} , has been intensively studied by Grabovsky, Sage, and subsequent coworkers in the case when $\Psi(\mathbf{k})$ is only a function of $\mathbf{k}/|\mathbf{k}|$. Their progress is summarized in [14, 15]. To apply their results to our setting a relatively easy task remains, namely to identify the subspaces \mathcal{K} and \mathcal{S} associated with each \mathcal{K}_0 . We caution, though, that some of these results are for equations such as thermoelasticity where the field $\mathbf{E}(\mathbf{x})$ has constant components, independent of \mathbf{x} , such as the temperature increment θ . Then $\mathbf{E}(\mathbf{x})$ is not square integrable, and our analysis does not apply in its current form. It may apply if we expand the constitutive law and treat those terms involving constant fields as source terms. Thus, for example, consider the thermoelastic equation

$$\boldsymbol{\epsilon}(\mathbf{x}) = \mathcal{S}(\mathbf{x})\boldsymbol{\sigma}(\mathbf{x}) + \boldsymbol{\alpha}(\mathbf{x})\theta, \quad \boldsymbol{\epsilon} = [\nabla \mathbf{u} + (\nabla \mathbf{u})^T]/2, \quad \nabla \cdot \boldsymbol{\sigma} = 0, \quad (10.1)$$

where $\boldsymbol{\epsilon}(\mathbf{x})$ is the strain, $\mathbf{u}(\mathbf{x})$ is the displacement field, $\boldsymbol{\sigma}(\mathbf{x})$ is the stress, $\theta = T - T_0$ is the change in temperature T measured from some base temperature T_0 , $\mathcal{S}(\mathbf{x})$ is the compliance tensor (inverse elasticity tensor), and $\boldsymbol{\alpha}(\mathbf{x})$ is the tensor of thermal expansion. Provided $\boldsymbol{\alpha}(\mathbf{x})$ is square integrable, we can treat $\mathbf{h}(\mathbf{x}) = -\boldsymbol{\alpha}(\mathbf{x})\theta$ as our source term, $\boldsymbol{\sigma}(\mathbf{x})$ as the field $\mathbf{E}(\mathbf{x})$, and $\boldsymbol{\epsilon}(\mathbf{x})$ as the field $\mathbf{J}(\mathbf{x})$. Of course, there could be additional source terms, not just those arising from the thermal expansion.

For many other equations of interest, such as wave equations in lossy media, $\Psi(\mathbf{k})$ for $\mathbf{k} \neq 0$ has the form

$$\Psi(\mathbf{k}) = \frac{1}{q(\mathbf{k})} \sum_{j=0}^p |k|^j \Psi_j(\mathbf{k}/|\mathbf{k}|), \quad (10.2)$$

where $q(\mathbf{k})$ is scalar-valued, and so (2.9) holds for all $\mathbf{k} \neq 0$ if and only if for all unit vectors \mathbf{n} , and for $j = 0, 1, \dots, p$,

$$\mathbf{A}_1 \Psi_j(\mathbf{n}) \mathbf{A}_2 \in \mathcal{K}, \quad \text{for all } \mathbf{A}_1, \mathbf{A}_2 \in \mathcal{K}. \quad (10.3)$$

Thus, we are confronted with algebraic questions that are essentially the same as those investigated by Grabovsky, Sage, and subsequent coworkers. The next step will be to search for specific examples of physical relevance, beyond those encountered in the theory of exact relations for composites. Irrespective of whether such exact relations exist for wave equations in lossy media, it is to be emphasized that the current theory already applies to many of the examples studied by Grabovsky, Sage, and coworkers as summarized in the book [15].

We also made large strides in developing the theory of “boundary field equalities” and provided for the first time a general theory for exact relations satisfied by the DtN map for bodies containing appropriate inhomogeneous media. Again, examples are needed to illuminate the theory and bring our results from an abstract setting to practice. Our boundary field equalities relied heavily on the existence of an appropriate constitutive law inside the body, or at least by imposing nonlinear constraints on fields so they could be related by an appropriate constitutive law, where the tensor $\mathbf{L}(\mathbf{x})$ entering the constitutive law depends upon the local fields. An interesting question is: Are there boundary field equalities, beyond the standard conservation laws, that do not require this of the interior fields? For example, in the setting of Sect. 8 one may ask about the existence of sets \mathcal{B} and appropriate boundary conditions in the two-dimensional case where \mathbf{Q} is a divergence-free $2 \times s$ matrix-valued field and s is not the square of an even integer, and in the three-dimensional case where \mathbf{Q} is a divergence-free $2 \times s$ matrix-valued field and $s/3$ is not the square of an even integer. (Of course, one may add to \mathbf{Q} an arbitrary number of “dummy” divergence-free columns that do not participate in the analysis, so one would want to exclude such trivial manipulations).

In our analysis we made heavy use of the tensor \mathbf{L}_0 , but the Green’s function $\mathbf{G}(\mathbf{x}, \mathbf{x}^0)$ is independent of \mathbf{L}_0 . We could have replaced \mathbf{L}_0 by any other tensor \mathbf{L}'_0 on the manifold $\mathcal{M} = W_{\mathbf{M}}^{-1}(\mathcal{K}_0)$ (where \mathcal{K}_0 consists of all self-adjoint maps in \mathcal{K}) and the analysis would have carried through. This raises the question, which we have not explored, as to whether this different choice of \mathbf{L}_0 would yield new constraints on $\mathbf{G}(\mathbf{x}, \mathbf{x}^0)$, or just recover those obtained with the original choice of \mathbf{L}_0 .

Author details

¹Department of Mathematics, University of Utah, Salt Lake City, UT, USA,

²Department of Mathematics, University of Houston, Houston, TX, USA.

Acknowledgements

GWM is grateful to the National Science Foundation for support through the Research Grants DMS-1211359 and DMS-1814854. Both authors thank the Institute for Mathematics and its Applications at the University of Minnesota for hosting their visit there during the Fall 2016 where this work was initiated as part of the program on Mathematics and Optics. Yury Grabovsky and the referees are thanked for their comments which led to significant improvements of the manuscript.

Appendix: Some physical equations that can be expressed in the required canonical form

Here we give examples of some physical equations that can be expressed in the required canonical form. The examples are by no means comprehensive: for further examples, see [26] Chap. 2, [35] Chap. 1, [27, 28] and the appendix of [29]. In the equations that follow we omit the source terms. We emphasize that if one is interested in exact relations satisfied by the DtN map and the associated boundary field equalities, then it is not necessary that the source terms have a physical significance provided they are localized outside the body Ω .

The simplest canonical equations are those of electrical conductivity

$$\underbrace{\mathbf{j}(\mathbf{x})}_{\mathbf{J}(\mathbf{x})} = \underbrace{\boldsymbol{\sigma}(\mathbf{x})}_{\mathbf{L}(\mathbf{x})} \underbrace{\mathbf{e}(\mathbf{x})}_{\mathbf{E}(\mathbf{x})}, \quad \nabla \cdot \mathbf{j} = 0, \quad \mathbf{e} = -\nabla V = 0, \quad (10.4)$$

where $\mathbf{j}(\mathbf{x})$ and $\mathbf{e}(\mathbf{x})$ are the electrical current and electric field and $V(\mathbf{x})$ is the electrical potential. The boundary fields $\partial\mathbf{J}$ and $\partial\mathbf{E}$ are the flux $\mathbf{n} \cdot \mathbf{j}(\mathbf{x})$ and boundary voltage $V(\mathbf{x})$, respectively, with $\mathbf{x} \in \partial\Omega$ and \mathbf{n} being the outward normal to Ω . As displayed in the table at the beginning of Sect. 2.1 in [26] (adapted from one of Batchelor [4]), the equations for dielectrics, magnetostatics, heat conduction, particle diffusion, flow in porous media, and antiplane elasticity all take the same form as (10.4) and so any analysis applicable to (10.4) applies to them as well.

Another important example is that of linear elasticity,

$$\underbrace{\boldsymbol{\sigma}(\mathbf{x})}_{\mathbf{J}(\mathbf{x})} = \underbrace{\mathbf{C}(\mathbf{x})}_{\mathbf{L}(\mathbf{x})} \underbrace{\boldsymbol{\epsilon}(\mathbf{x})}_{\mathbf{E}(\mathbf{x})}, \quad \nabla \cdot \boldsymbol{\sigma} = 0, \quad \boldsymbol{\epsilon} = [\nabla \mathbf{u} + (\nabla \mathbf{u})^T]/2, \quad (10.5)$$

where $\boldsymbol{\sigma}(\mathbf{x})$ (not to be confused for the conductivity tensor field) is the stress, $\boldsymbol{\epsilon}(\mathbf{x})$ is the strain, $\mathbf{u}(\mathbf{x})$ is the displacement, and $\mathbf{C}(\mathbf{x})$ is the elasticity tensor field. The boundary fields $\partial\mathbf{J}(\mathbf{x})$ and $\partial\mathbf{E}(\mathbf{x})$ are the traction $\mathbf{n} \cdot \boldsymbol{\sigma}(\mathbf{x})$ and boundary displacement field $\mathbf{u}(\mathbf{x})$, respectively.

It is also possible to have equations that couple fields together, such as the magneto-electric equations,

$$\underbrace{\begin{pmatrix} \mathbf{d} \\ \mathbf{b} \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} \boldsymbol{\epsilon} & \boldsymbol{\beta} \\ \boldsymbol{\beta}^T & \boldsymbol{\mu} \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \mathbf{e} \\ \mathbf{h} \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \quad \nabla \cdot \mathbf{d} = \nabla \cdot \mathbf{b} = 0, \quad \mathbf{e} = -\nabla V, \quad \mathbf{h} = -\nabla \psi, \quad (10.6)$$

where \mathbf{d} and \mathbf{b} are the electric displacement field and magnetic induction, \mathbf{e} and \mathbf{h} are the electric and magnetic fields, V and ψ are the electric potential and magnetic scalar potential (assuming there are no free currents), $\boldsymbol{\epsilon}(\mathbf{x})$ is the free-body electrical permittivity (with $\mathbf{h} = 0$), $\boldsymbol{\beta}(\mathbf{x})$ is the second-order magnetoelectric coupling tensor, $\boldsymbol{\mu}(\mathbf{x})$ is the free-body magnetic permeability (with $\mathbf{e} = 0$). The boundary fields $\partial\mathbf{J}(\mathbf{x})$ and $\partial\mathbf{E}(\mathbf{x})$ are then the flux pair $(\mathbf{n} \cdot \mathbf{d}(\mathbf{x}), \mathbf{n} \cdot \mathbf{b}(\mathbf{x}))$ and the potential pair $(V(\mathbf{x}), \psi(\mathbf{x}))$, respectively, with $\mathbf{x} \in \partial\Omega$.

Thermoelectricity also takes this form, but one has to be careful in defining the fields to ensure that the associated tensor $\mathbf{L}(\mathbf{x})$ is symmetric (see, e.g., Sect. 2.4 in [26]).

Fields that are coupled together need not have the same tensorial rank, an example being the equations of piezoelectricity,

$$\underbrace{\begin{pmatrix} \boldsymbol{\epsilon} \\ \mathbf{d} \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} \boldsymbol{\mathcal{S}} & \boldsymbol{\mathcal{D}} \\ \boldsymbol{\mathcal{D}}^T & \boldsymbol{\epsilon} \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \boldsymbol{\sigma} \\ \mathbf{e} \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \quad (10.7)$$

where $\boldsymbol{\mathcal{S}}(\mathbf{x})$ is the compliance tensor under short-circuit boundary conditions (i.e., with $\mathbf{e} = 0$), $\boldsymbol{\mathcal{D}}(\mathbf{x})$ is the piezoelectric stress coupling tensor, and $\boldsymbol{\epsilon}(\mathbf{x})$ is the free-body dielectric tensor (i.e., with $\boldsymbol{\tau} = 0$). The strain field $\boldsymbol{\epsilon}$, electric displacement field \mathbf{d} , stress field $\boldsymbol{\sigma}$, and electric field \mathbf{e} satisfy the usual differential constraints:

$$\boldsymbol{\epsilon} = [\nabla \mathbf{u} + (\nabla \mathbf{u})^T]/2, \quad \nabla \cdot \mathbf{d} = 0, \quad \nabla \cdot \boldsymbol{\sigma} = 0, \quad \mathbf{e} = -\nabla V. \quad (10.8)$$

Since the stresses and strains are symmetric matrices, $\boldsymbol{\mathcal{D}}$ is a third-order tensor that maps vectors to symmetric matrices. The boundary fields $\partial\mathbf{J}(\mathbf{x})$ and $\partial\mathbf{E}(\mathbf{x})$ are the displacement, flux pair $(\mathbf{u}(\mathbf{x}), \mathbf{n} \cdot \mathbf{d}(\mathbf{x}))$ and the traction, voltage pair $(\mathbf{n} \cdot \boldsymbol{\sigma}, V)$, respectively.

Of course, more than two fields can be coupled together. Thus, by combining a piezo-electric material and a magnetostrictive material in a composite, we can obtain a material where there is coupling between electric fields, elastic fields, and magnetic fields.

For fields varying in time at constant frequency ω , with wavelengths and attenuation lengths much bigger than the size of the body under consideration, the quasistatic equations are applicable. For dielectrics these take the same form as (10.4):

$$\mathbf{d}(\mathbf{x}) = \boldsymbol{\varepsilon}(\mathbf{x})\mathbf{e}(\mathbf{x}), \quad \nabla \cdot \mathbf{j} = 0, \quad \mathbf{e} = -\nabla V = 0, \tag{10.9}$$

where everything is now complex-valued: $\mathbf{d}(\mathbf{x})$ and $\mathbf{e}(\mathbf{x})$ are the complex-valued electrical displacement field and electric field, $V(\mathbf{x})$ is the complex-valued electrical potential, and $\boldsymbol{\varepsilon}(\mathbf{x})$ is the complex-valued electrical permittivity. (The physical displacement field, electric field, and potential are the real parts of $\mathbf{d}(\mathbf{x})e^{-i\omega t}$, $\mathbf{e}(\mathbf{x})e^{-i\omega t}$, and $Ve^{-i\omega t}$, respectively). Let us set

$$\mathbf{d} = \mathbf{d}' + i\mathbf{d}'', \quad \mathbf{e} = \mathbf{e}' + i\mathbf{e}'', \quad V = V' + iV'', \quad \boldsymbol{\varepsilon} = \boldsymbol{\varepsilon}' + \boldsymbol{\varepsilon}'', \tag{10.10}$$

where the primed fields denote the real parts, while the doubled primed fields denote the imaginary parts. Physically, $\boldsymbol{\varepsilon}''(\mathbf{x})$ is associated with electrical energy loss into heat and is positive semidefinite. Assuming it is positive definite and that an inverse $[\boldsymbol{\varepsilon}'']^{-1}$ exists, substitution of (10.10) in (10.9), followed by suitable manipulation, gives the equivalent coupled field equations of Gibiansky and Cherkaev [7]:

$$\underbrace{\begin{pmatrix} \mathbf{e}'' \\ \mathbf{d}'' \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} [\boldsymbol{\varepsilon}'']^{-1} & [\boldsymbol{\varepsilon}'']^{-1}\boldsymbol{\varepsilon}' \\ \boldsymbol{\varepsilon}'[\boldsymbol{\varepsilon}'']^{-1} & \boldsymbol{\varepsilon}'' + \boldsymbol{\varepsilon}'[\boldsymbol{\varepsilon}'']^{-1}\boldsymbol{\varepsilon}' \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} -\mathbf{d}' \\ \mathbf{e}' \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \tag{10.11}$$

Clearly \mathbf{L} is real and symmetric and by inspection of the quadratic form associated with \mathbf{L} one sees that it is positive definite. Now $\partial\mathbf{J}$ consists of the voltage, flux pair $(V'', \mathbf{n} \cdot \mathbf{d}'')$ while $\partial\mathbf{E}$ consists of the flux, voltage pair $(-\mathbf{n} \cdot \mathbf{d}', V')$. As Gibiansky and Cherkaev show, similar manipulations can be done for viscoelasticity in the quasistatic limit where the equations have the form (10.5), but with all fields being complex. More generally, the Gibiansky–Cherkaev approach can be applied to equations where the tensor entering the constitutive law is not self-adjoint, but its self-adjoint part is positive definite, to an equivalent form where the tensor $\mathbf{L}(\mathbf{x})$ entering the constitutive law is self-adjoint and positive definite [25] (see also Sect. 13.4 of [26]): such manipulations can be applied, for example, to electrical conduction in the presence of a magnetic field where, due to the Hall effect, the conductivity tensor $\boldsymbol{\sigma}(\mathbf{x})$ entering (10.4) is not symmetric.

Wave equations, can be expressed in the form (4.12) with an identity like (4.15) holding. For example, at fixed frequency ω with a $e^{-i\omega t}$ time dependence, as recognized in [32] the acoustic equations, with $P(\mathbf{x})$ the (complex) pressure, $\mathbf{v}(\mathbf{x})$ the (complex) velocity, $\boldsymbol{\rho}(\mathbf{x}, \omega)$ the effective mass density matrix, and $\kappa(\mathbf{x}, \omega)$ the bulk modulus, take the form

$$\underbrace{\begin{pmatrix} -i\mathbf{v} \\ -i\nabla \cdot \mathbf{v} \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} -(\omega\rho)^{-1} & 0 \\ 0 & \omega/\kappa \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \nabla P \\ P \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \tag{10.12}$$

(and $\partial\mathbf{E}$ and $\partial\mathbf{J}$ can be identified with the boundary values of $P(\mathbf{x})$ and $\mathbf{n} \cdot \mathbf{v}(\mathbf{x})$ at $\partial\Omega$, respectively). Here we allow for effective mass density matrices that, at a given frequency, can be anisotropic and complex-valued as may be the case in metamaterials [30,33,44,50].

Maxwell's equations, with $\mathbf{e}(\mathbf{x})$ the (complex) electric field, $\mathbf{h}(\mathbf{x})$ the (complex) magnetizing field, $\boldsymbol{\mu}(\mathbf{x}, \omega)$ the magnetic permeability, $\boldsymbol{\varepsilon}(\mathbf{x})$ the electric permittivity, take the form

$$\underbrace{\begin{pmatrix} -i\mathbf{h} \\ i\nabla \times \mathbf{h} \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} -[\omega\boldsymbol{\mu}]^{-1} & 0 \\ 0 & \omega\boldsymbol{\varepsilon} \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \nabla \times \mathbf{e} \\ \mathbf{e} \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \quad (10.13)$$

(and $\partial\mathbf{E}$ and $\partial\mathbf{J}$ can be identified with the tangential values of $\mathbf{e}(\mathbf{x})$ and $\mathbf{h}(\mathbf{x})$ at $\partial\Omega$, respectively). The linear elastodynamic equations, with $\mathbf{u}(\mathbf{x})$ the (complex) displacement, $\boldsymbol{\sigma}(\mathbf{x})$ the (complex) stress, $\mathcal{C}(\mathbf{x}, \omega)$ the elasticity tensor, $\boldsymbol{\rho}(\mathbf{x}, \omega)$ the effective mass density matrix, take the form

$$\underbrace{\begin{pmatrix} -\boldsymbol{\sigma}/\omega \\ -\nabla \cdot \boldsymbol{\sigma}/\omega \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} -\mathcal{C}/\omega & 0 \\ 0 & \omega\boldsymbol{\rho} \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} [\nabla\mathbf{u} + (\nabla\mathbf{u})^T]/2 \\ \mathbf{u} \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}, \quad (10.14)$$

(and $\partial\mathbf{E}$ and $\partial\mathbf{J}$ can be identified with the values of $\mathbf{u}(\mathbf{x})$ and the traction $\mathbf{n} \cdot \boldsymbol{\sigma}(\mathbf{x})$ at $\partial\Omega$, respectively). The preceding three equations have been written in this form so $\text{Im}\mathbf{L}(\mathbf{x}) \geq 0$ when $\text{Im}\omega \geq 0$, where complex frequencies have the physical meaning of the solution increasing exponentially in time. Under assumptions that the material moduli are lossy, or that the frequency ω is complex with positive imaginary part, one can easily manipulate them into equivalent forms similar to the Gibiansky–Cherkaev form in (10.11) with a positive semidefinite tensor entering the constitutive law [32,34]. Of course, the boundary fields $\partial\mathbf{E}$ and $\partial\mathbf{J}$ then need to be appropriately redefined.

For thin plates, the dynamic plate equations at constant frequency can be written in the form

$$\underbrace{\begin{pmatrix} i\mathbf{M} \\ \nabla \cdot (\nabla \cdot \mathbf{M}) \end{pmatrix}}_{\mathbf{J}(\mathbf{x})} = \underbrace{\begin{pmatrix} -\mathcal{D}(\mathbf{x})/\omega & 0 \\ 0 & h(\mathbf{x})\omega\rho(\mathbf{x}) \end{pmatrix}}_{\mathbf{L}(\mathbf{x})} \underbrace{\begin{pmatrix} \nabla\nabla v \\ iv \end{pmatrix}}_{\mathbf{E}(\mathbf{x})}. \quad (10.15)$$

Here $\mathbf{M}(\mathbf{x}, t)$ is the (complex) bending moment tensor, $\mathcal{D}(\mathbf{x})$ is the fourth-order tensor of plate rigidity coefficients, $h(\mathbf{x})$ is the plate thickness, $\rho(\mathbf{x})$ is the density, and $v = \partial w/\partial t$ is the velocity of the (complex) vertical deflection $w(\mathbf{x}, t)$ of the plate. Note that the matrix $\mathbf{L}(\mathbf{x})$ has positive definite imaginary part when ω has positive imaginary part. $\partial\mathbf{E}$ can be identified with the boundary values of the pair $(\nabla v, v)$ while $\partial\mathbf{J}$ can be identified with the boundary values of the pair $(\mathbf{M}\mathbf{n}, (\nabla \cdot \mathbf{M}) \cdot \mathbf{n})$, in which \mathbf{n} is the outward normal to $\partial\Omega$. Again, when the material moduli are lossy, or the frequency ω is complex with positive imaginary part, this can be manipulated into the Gibiansky–Cherkaev form in (10.11) with a positive semidefinite tensor entering the constitutive law, and with appropriately redefined boundary fields.

Further examples of wave equations at constant frequency that can be represented in the required form are given in the appendix of [29].

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 3 March 2018 Accepted: 23 January 2019

Published online: 13 February 2019

References

1. Allaire, G.: *Shape Optimization by the Homogenization Method*. Applied Mathematical Sciences, vol. 146. Springer, Berlin (2002)
2. Backus, G.E.: Long-wave elastic anisotropy produced by horizontal layering. *J. Geophys. Res.* **67**(11), 4427–4440 (1962)
3. Ball, J.M.: Convexity conditions and existence theorems in non-linear elasticity. *Arch. Ration. Mech. Anal.* **63**(4), 337–403 (1977)
4. Batchelor, G.K.: Transport properties of two-phase materials with random structure. *Annu. Rev. Fluid Mech.* **6**, 227–255 (1974)
5. Beněšová, B., Kružík, Martin: weak lower semicontinuity of integral functionals and applications. *SIAM Rev.* **59**(4), 703–766 (2017)
6. Cherkav, A.V.: *Variational Methods for Structural Optimization*. Applied Mathematical Sciences. Springer, Berlin (2000)
7. Cherkav, A.V., Gibiansky, L.V.: Variational principles for complex conductivity, viscoelasticity, and similar problems in media with complex moduli. *J. Math. Phys.* **35**(1), 127–145 (1994)
8. Clark, K.E., Milton, G.W.: Modeling the effective conductivity function of an arbitrary two-dimensional polycrystal using sequential laminates. *Proc. R. Soc. Edinb.* **124A**(4), 757–783 (1994)
9. Dell'Antonio, G.F., Figari, R., Orlandi, E.: An approach through orthogonal projections to the study of inhomogeneous or random media with linear response. *Ann. Inst. Henri Poincaré (A) Phys. théor.* **44**(1), 1–28 (1986)
10. Dykhne, A. M.: Conductivity of a two-dimensional two-phase system. *Zhurnal eksperimental'noi i teoreticheskoi fiziki / Akademiia Nauk SSSR*, 59:110–115, July 1970. English translation in *Soviet Physics JETP* 32(1):63–65 (1971)
11. Eyre, D.J., Milton, G.W.: A fast numerical scheme for computing the response of composites using grid refinement. *Eur. Phys. J. Appl. Phys.* **6**(1), 41–47 (1999)
12. Faraco, D., Székelyhidi, L.: Tartars conjecture and localization of the quasiconvex hull in $\mathbb{R}^{2 \times 2}$. *Acta Math.* **200**(2), 279–305 (2008)
13. Grabovsky, Y.: Exact relations for effective tensors of polycrystals. I: necessary conditions. *Arch. Ration. Mech. Anal.* **143**(4), 309–329 (1998)
14. Grabovsky, Y.: Algebra, geometry and computations of exact relations for effective moduli of composites. In: Gianfranco Capriz and Paolo Maria Mariano, editors, *Advances in Multifield Theories of Continua with Substructure, Modelling and Simulation in Science, Engineering and Technology*, pp. 167–197. Birkhäuser Verlag, Boston (2004)
15. Grabovsky, Y.: *Composite Materials: Mathematical Theory and Exact Relations*. IOP Publishing, Bristol (2016)
16. Grabovsky, Y.: From microstructure-independent formulas for composite materials to rank-one convex, non-quasiconvex functions. *Arch. Ration. Mech. Anal.* **227**(2), 607–636 (2018)
17. Grabovsky, Y., Milton, G.W., Sage, D.S.: Exact relations for effective tensors of composites: Necessary conditions and sufficient conditions. *Commun. Pure Appl. Math. (New York)* **53**(3), 300–353 (2000)
18. Grabovsky, Y., Sage, D.S.: Exact relations for effective tensors of polycrystals. II: applications to elasticity and piezoelectricity. *Arch. Ration. Mech. Anal.* **143**(4), 331–356 (1998)
19. Kohler, W., Papanicolaou, G. C.: Bounds for the effective conductivity of random media. In Burridge, R., Childress, S., Papanicolaou, G.C. (eds.) *Macroscopic Properties of Disordered Media: Proceedings of a Conference Held at the Courant Institute, June 1–3, 1981, Lecture Notes in Physics*, vol. 154, pp. 111–130. Springer, Berlin (1982)
20. Kohn, R.V.: The relaxation of a double-well energy. *Contin. Mech. Thermodyn.* **3**(3), 193–236 (1991)
21. Lax, P.D.: *Functional Analysis*. Wiley, New York (2002)
22. Levin, V. M.: Thermal expansion coefficients of heterogeneous materials. *Inzhenernyi Zhurnal. Mekhanika Tverdogo Tela: MTT*, 2(1):88–94 (1967). English translation in *Mechanics of Solids* 2(1):58–61 (1967)
23. Milgrom, M.: Linear response of general composite systems to many coupled fields. *Phys. Rev. B: Condens. Matter Mater. Phys.* **41**(18), 12484–12494 (1990)
24. Milton, G.W.: Multicomponent composites, electrical networks and new types of continued fraction. I. *Commun. Math. Phys.* **111**(2), 281–327 (1987)
25. Milton, G.W.: On characterizing the set of possible effective tensors of composites: the variational method and the translation method. *Commun. Pure Appl. Math. (New York)* **43**(1), 63–125 (1990)
26. Milton G.W.: *The Theory of Composites*. Cambridge Monographs on Applied and Computational Mathematics, vol. 6. In: Ciarlet, P.G., Iserles, A., Kohn, R.V., Wright, M.H. (eds.) Cambridge University Press, Cambridge (2002)
27. Milton, G.W.: Sharp inequalities that generalize the divergence theorem: an extension of the notion of quasi-convexity. *Proc. R. Soc. A: Math. Phys. Eng. Sci.* **469**(2157), 20130075 (2013). See addendum [29]
28. Milton, G.W.: Addendum to “Sharp inequalities that generalize the divergence theorem: an extension of the notion of quasi-convexity. *Proc. R. Soc. A: Math. Phys. Eng. Sci.* **471**(2176), 20140886 (2015). See [28]
29. Milton, G.W.: A new route to finding bounds on the generalized spectrum of many physical operators. *J. Math. Phys.* **59**, 061508 (2018)
30. Milton, G.W., Briane, M., Willis, J.R.: On cloaking for elasticity and physical equations with a transformation invariant form. *New J. Phys.* **8**(10), 248 (2006)
31. Milton, G.W., Golden, K.M.: Representations for the conductivity functions of multicomponent composites. *Commun. Pure Appl. Math. (New York)* **43**(5), 647–671 (1990)
32. Milton, G.W., Seppecher, P., Guy, B.: Minimization variational principles for acoustics, elastodynamics and electromagnetism in lossy inhomogeneous bodies at fixed frequency. *Proc. R. Soc. A: Math. Phys. Eng. Sci.* **465**(2102), 367–396 (2009)
33. Milton, G.W., Willis, J.R.: On modifications of Newton's second law and linear continuum elastodynamics. *Proc. R. Soc. A: Math. Phys. Eng. Sci.* **463**(2079), 855–880 (2007)
34. Milton, G.W., Willis, J.R.: Minimum variational principles for time-harmonic waves in a dissipative medium and associated variational principles of Hashin–Shtrikman type. *Proc. R. Soc. A: Math. Phys. Eng. Sci.* **466**(2122), 3013–3032 (2010)

35. Milton, G.W. (ed.): *Extending the Theory of Composites to Other Areas of Science*. Milton–Patton Publishers, Salt Lake City (2016)
36. Moulinec, H., Suquet, P.M.: A fast numerical method for computing the linear and non-linear properties of composites. *C. r. Séances l'Acad. Sci. Sér. II* **318**, 1417–1423 (1994)
37. Müller, S., Šverák, V.: Convex integration for Lipschitz mappings and counterexamples to regularity. *Ann. Math.* **157**(3), 715–742 (2003)
38. Murat, F.: Compacité par compensation. *Ann. Sc. Norm. Super. Pisa, Classe Sci. Ser. IV* **5**(3), 489–507 (1978). (French) [Compactness through compensation]
39. Murat, F.: Compacité par compensation: condition nécessaire et suffisante de continuité faible sous une hypothèse de rang constant. (French) [Compensated compactness: Necessary and sufficient conditions for weak continuity under a constant-rank hypothesis]. *Ann. Sc. Norm. Super. Pisa, Classe Sci. Ser. IV*
40. Murat, F.: A survey on compensated compactness. In: Cesari, L. (ed.) *Contributions to Modern Calculus of Variations*. Pitman Research Notes in Mathematics Series, vol. 148, pp. 145–183, Harlow, Essex (1987). Longman Scientific and Technical. Papers from the symposium marking the centenary of the birth of Leonida Tonelli held in Bologna, May 13–14, (1985)
41. Olver, P.J., Sivaloganathan, J.: The structure of null Lagrangians. *Nonlinearity (Bristol)* **1**(2), 389–398 (1988)
42. Pablo, P.: Weak continuity and weak lower semicontinuity for some compensation operators. *Proc. R. Soc. Edinb. Sect. A, Math. Phys. Sci.* **113**(3–4), 267–279 (1989)
43. Raitums, U.E.: On the local representation of G -closure. *Arch. Ration. Mech. Anal.* **158**(3), 213–234 (2001)
44. Schoenberg, M., Sen, P.N.: Properties of a periodically stratified acoustic half-space and its relation to a Biot fluid. *J. Acoust. Soc. Am.* **73**(1), 61–67 (1983)
45. Tartar, L.: Compensated compactness and applications to partial differential equations. In: Knops, R.J. (ed.) *Nonlinear Analysis and Mechanics, Heriot–Watt Symposium, Volume IV*. Research Notes in Mathematics, vol. 39, pp. 136–212. Pitman Publishing Ltd, London (1979)
46. Tartar, L.: Estimation de coefficients homogénéisés. (French) [Estimation of homogenization coefficients]. In: Glowinski, R., Lions, J.L. (eds.) *Computing Methods in Applied Sciences and Engineering: Third International Symposium, Versailles, France, December 5–9, 1977*. Lecture Notes in Mathematics, vol. 704, pp. 364–373. Springer, Berlin (1979). English translation in *Topics in the Mathematical Modelling of Composite Materials*, pp. 9–20, Cherkaev, A., Kohn, R. (eds.). ISBN 0-8176-3662-5
47. Tartar, L.: *The General Theory of Homogenization: A Personalized Introduction*. Lecture Notes of the Unione Matematica Italiana. Springer, Berlin (2009)
48. Thaler, A.E., Milton, G.W.: Exact determination of the volume of an inclusion in a body having constant shear modulus. *Inverse Probl.* **30**(12), 125008 (2014)
49. Torquato, S.: *Random Heterogeneous Materials: Microstructure and Macroscopic Properties*. Interdisciplinary Applied Mathematics, vol. 16. Springer, Berlin (2002)
50. Willis, J.R.: The non-local influence of density variations in a composite. *Int. J. Solids Struct.* **21**(7), 805–817 (1985)
51. Zhang, K.: On the structure of quasiconvex hulls. *Ann. l'Inst. Henri Poincaré. Anal. Linéaire* **15**(6), 663–686 (1998)
52. Zhikov, V. V.: Estimates for the homogenized matrix and the homogenized tensor. *Uspekhi Matematicheskikh Nauk = Russ. Math. Surv.* 46:49–109 (1991). English translation in *Russ. Math. Surv.* 46(3):65–136 (1991)