

Answers to Odd-Numbered Problems

CHAPTER 1

Exercises 1.1

- (a) ordinary, first order
(c) partial, second order
(e) ordinary, third order
(g) ordinary, second order
- Both y and z are solutions.
- Both y and z are solutions.
- Both u_1 and u_2 are solutions.
- u_1 is a solution; u_2 is not a solution.
- $y = 16x^2 + C_1x + C_2$
- $y = Ce^{3x}$.
- $r = 2, -2$; $y_1(x) = e^{2x}$ and $y_2(x) = e^{-2x}$ are solutions.
- $r = 3$; $y(x) = e^{3x}$ is a solution.
- $r = 1/2, r = 4$; $y_1(x) = e^{x/2}$ and $y_2(x) = e^{4x}$ are solutions.
- No real values of r ; $r = 2 \pm 4i$ are complex values.
- $r = 2, -2$; $y_1(x) = e^{2x}$ and $y_2(x) = e^{-2x}$ are solutions.
- $r = 3, -3$; $y_1(x) = x^3$ and $y_2(x) = x^{-3}$ are solutions.
- $r = 2$; $y = x^2$ is a solution.
- $r = 5, -1$; $y_1(x) = x^5$ and $y_2(x) = x^{-1}$ are solutions.

Exercises 1.3

- (b) $y = 2e^{5x}$.
- (b) $y = \frac{e}{e - 2e^x}$.
- (b) $y = -\sin 3x + \frac{1}{3} \cos 3x$.
- (b) $y = -\frac{17}{4}x + 9\sqrt{x}$.
(c) y' and y'' are not defined at $x = 0$; there is no solution to $y'(0) = 2$.
- $xy' - 3y + 3 = 0$.
- $y' = \frac{2y^3 - 6}{3xy^2}$
- $y' - 2y = -4e^{-2x}$.
- $y'' = 0$.
- $y'' - 4y' + 4y = 0$.
- $x^2y'' + xy' - y = 0$.
- $y'' + 9y = 0$.

25. $y''' = 0$.

CHAPTER 2

Exercises 2.1

1. $y = -\frac{1}{2} + Ce^{2x}$.

3. $y = 1 + Ce^{-x^2}$.

5. $y = e^{-x} + Ce^x$.

7. $y = x^{-2} \sin x + Cx^{-2}$.

9. $y = \frac{2}{9}(x+1)^{5/2} + C(x+1)^{-2}$.

11. $y = \sin x \cos x + C \cos x = \frac{1}{2} \sin 2x + C \cos x$.

13. $y = e^x + \frac{C}{x}$.

15. $y = x(\ln x)^2 + Cx$.

17. $y = 1 + Ce^{-e^x}$.

19. $y = x - 1 + 2e^{-x}$.

21. $y = \frac{\ln(1+e^x)}{e^x} + (e - \ln 2)e^{-x}$.

23. $y = \frac{5 - \cos 2x}{2 \sin x}$.

25. (a) $y = \frac{c}{b} + Ce^{-bx/a}$ (c) $y = \frac{c}{b} + \left(\alpha - \frac{c}{b}e^{-bx/a}\right)$

Exercises 2.2

1. $y = \left(\frac{x^2}{4} + C\right)^2$.

3. $\tan^{-1} y = x^3 + C$ or $y = \tan(x^3 + C)$.

5. $\cot y = \ln \sqrt{\frac{1-x}{1+x}} + C$.

7. $e^{-y} = e^x - xe^x + C$.

9. $\sqrt{y^2 - 1} = \frac{x}{1 + Cx}$.

11. $y^2 = C(\ln x)^2 - 1$.

13. $\ln |y| = -\ln |x| - \frac{1}{x} - 1$.

15. $y = xe^{x^2-1}$.

17. $y + \ln |y| = \frac{1}{3}x^3 - x - 5$.

19. $y = \frac{x + C}{1 - Cx}$

Exercises 2.3

1. $y = \frac{2}{Cx - 3x^3}$.

3. $y = (Ce^{2x} - e^x)^2.$
5. $y = \frac{1}{\sqrt[3]{Cx^3 - 2x^3 \ln x}}.$
7. $y^2 = Cx + x^2.$
9. $x \ln x + \frac{x+y}{e^{y/x}} = Cx.$
11. $\csc(y/x) - \cot(y/x) = Cx.$
13. $y^2 = \frac{C}{1+x^2} - 1.$
15. $y = \frac{\ln |\sec x + \tan x| + C}{x}.$
17. $y + \ln |1 - y| = C - x - \ln |1 + x|.$
19. $y = -x \ln(C - \ln x).$
21. $y = C(3x^2 + 1)^{1/3} - 3.$
23. $y = \frac{1}{Cx + \ln x + 1}.$
25. $2y^3 = x^3 - Cx.$
27. (a) $u = \sin y$ (b) $\sin y = e^{-x^2}(4x + C).$

EXERCISES 2.4

Exercises 2.4.1

1. $x^2 + 3y^2 = C.$
3. $\frac{x^2}{2} + y^2 - 4y = C.$
5. $y^2 = \ln(\sin^2 x) + C.$
7. $y = -\frac{1}{2}x^2 + C.$
9. $x^2 + \frac{1}{2}y^2 = C;$ ellipses, center at the origin, major axis horizontal.
11. $x^2 + y^2 - Cy = 0.$

Exercises 2.4.2

1. (a) $A(t) = 50 \left(\frac{9}{10}\right)^{t/2} \approx 50e^{-0.05268t}.$ (b) $A(4) = 50 \left(\frac{9}{10}\right)^2 = 40.5$ grams.
(c) $T \approx 13.16$ hours.
3. $t = \frac{2 \ln 10}{\ln 2} \approx 6.64$ hours.
5. (a) $P(t) \approx 0.25e^{0.0421t}$ (b) ≈ 1.6573 square centimeters (c) ≈ 16.464 hours
7. (a) $P(t) \approx 4.5e^{0.01438t}.$ (b) 48.19 years (c) ≈ 6.93 billion.

Exercises 2.4.3

1. (a) $40.1^\circ.$ (b) 1.62 minutes.
3. (a) $u(t) = 150 - 100e^{\frac{t}{10} \ln(3/4)} = 150 - 100 \left(\frac{3}{4}\right)^{t/10}$

(b) $t = \frac{10 \ln(1/2)}{\ln(3/4)} \approx 24.09$ minutes

(c) The temperature will never reach 200° ; $\lim_{t \rightarrow \infty} u(t) = 150$

5. (a) Approximately 12:12 (b) Approximately 12:48

Exercises 2.4.4

1. (a) $v = \left(v_0 + \frac{g}{r}\right) e^{-rt} - \frac{g}{r}$ (b) $\lim_{t \rightarrow \infty} v = -\frac{g}{r}$.

(c) $y = y_0 + \frac{1}{r} \left(v_0 + \frac{g}{r}\right) (1 - e^{-rt}) - \frac{g}{r}t$

3. $k \approx 17.8$

Exercises 2.4.5

1. (a) $A(t) = 10,000 (1 - e^{-t/200})$ (b) $t = 200 \ln 5 \approx 322$ minutes

3. (a) $A(t) = \frac{9}{2} (1 - e^{-t/150})$ (b) $t = 150 \ln 3 \approx 165$ minutes

5. (a) $A(t) = \frac{3}{20} t(100 - t)$ (b) $\max = A(50) = 375$

Exercises 2.4.6

1. (a) 3259 people. (b) ≈ 6.89 days.

3. (a) $\frac{d^2y}{dt^2} = k \frac{dy}{dt} (M - 2y)$; $\frac{dy}{dt} > 0$ for $0 < y < M/2$, $\frac{dy}{dt} < 0$ for $y > M/2$.
 dy/dt has a maximum when $y = M/2$

5. $k \approx 0.0006$

Exercises 2.4.7

1. (a) $\frac{dP}{dt} = kt(1000 - R)$ (b) $P(t) = 1000 - Ce^{-kt^2/2}$ (c) $P(t) = 1000 - 950e^{-kt^2/2}$


3. $A(t) = 1000 \left(\frac{4}{5}\right)^{t^2/100}$

5. $A(t) = \frac{140,000}{140 + 3t}$

7. $a = \frac{\ln 2}{24}$, $b = \frac{\ln 2}{6}$


Exercises 2.5

1. (a) and (b)




(c) $y = e^x$

3. (a) and (b)



(c) $y = e^{x^2}$

5. Initial conditions: $y(0) = 0$ $y'(0) = 1$ $y''(0) = 2$



CHAPTER 3

Exercises 3.2

1. Yes
3. Yes
5. Yes
7. (a) $r = -1, r = 4$.
- (b) Fundamental set: $y_1(x) = x^{-1}, y_2(x) = x^4$; general solution: $y = C_1x^{-1} + C_2x^4$.
- (c) $y = \frac{9}{5}x^{-1} + \frac{1}{5}x^4$.
- (d) The trivial solution: $y \equiv 0$.
9. $y'' - 2y' - 3y = 0$.
11. $y'' = 0$.
13. $x^2y'' - 2xy' + 2y = 0$.
15. $W[y_1, y_2](x) = e^{-\int_a^x p(t) dt} \neq 0$ for all x .
17. $\{y_1(x) = x, y_2(x) = x^2\}$.
19. $\{y_1(x) = e^{x^2}, y_2(x) = e^{-x^2}\}$.
21. $\alpha\delta - \beta\gamma \neq 0$.
23. $W[y_1 + y_2, y_1 - y_2] = -2W[y_1, y_2]$.
25. Set $u(x) = \frac{y_2(x)}{y_1(x)}$. Then

$$u'(x) = \frac{y_1y_2' - y_2y_1'}{y_1^2} = \frac{W[y_1, y_2]}{y_1^2} \equiv 0.$$

Therefore, $u \equiv \lambda$ constant, which implies that $y_2 = \lambda y_1$.

Exercises 3.3

1. $y = C_1 e^{2x} + C_2 e^{-4x}$.
 3. $y = C_1 e^{5x} + C_2 x e^{5x}$.
 5. $y = e^{-2x} [C_1 \cos 3x + C_2 \sin 3x]$.
 7. $y = C_1 + C_2 e^{-2x}$.
 9. $y = C_1 e^{2\sqrt{3}x} + C_2 e^{-2\sqrt{3}x}$.
 11. $y = e^x [C_1 \cos x + C_2 \sin x]$.
 13. $y = C_1 e^{6x} + C_2 e^{-5x}$.
 15. $y = e^{-x/2} [C_1 \cos x/2 + C_2 \sin x/2]$.
 17. $y = C_1 e^{4x} + C_2 x e^{4x}$.
 19. $y = 2e^{2x} - e^{3x}$.
 21. $y = -3e^{-x} - 2xe^{-x}$.
 23. $y = -e^x \cos x$.
 25. $y'' + 3y' - 10y = 0$.
 27. $y'' + 4y = 0$.
 29. $y'' - \frac{5}{2}y' + y = 0$.
 31. $y'' + 2y' + 10y = 0$.
 33. $y'' + 16y = 0$.
 35. $y = (1 + \beta)e^{x/2} + (1 - \beta)e^{-x/2}$; $\beta = -1$.
 37. (a) $a^2 - 4b > 0$ (b) $a^2 - 4b = 0$ (c) $a^2 - 4b < 0$
 39. (a) $y'' + by = 0$, $b > 0$; general solution: $y = C_1 \cos x\sqrt{b} + C_2 \sin x\sqrt{b}$, all solutions are bounded.
 (b) $y'' + ay' = 0$; general solution: $y = C_1 + C_2 e^{-ax}$ and $\lim_{x \rightarrow \infty} y = C_1$.
- The solution that satisfies the initial conditions is: $y = \left(\alpha + \frac{\beta}{a}\right) - \frac{\beta}{a} e^{-ax}$; $k = \alpha + \frac{\beta}{a}$.
41. $r_1, r_2 = \frac{-a \pm \sqrt{a^2 - 4b}}{2} = \frac{-a}{2} \pm \frac{\sqrt{a^2 - 4b}}{2} = \alpha \pm \beta$.
- General solution:
- $$\begin{aligned}
 y = C_1 e^{(\alpha+\beta)x} + C_2 e^{(\alpha-\beta)x} &= C_1 e^{\alpha x} e^{\beta x} + C_2 e^{\alpha x} e^{-\beta x} \\
 &= e^{\alpha x} \left[(C_1 + C_2) \frac{e^{\beta x} + e^{-\beta x}}{2} + (C_1 - C_2) \frac{e^{\beta x} - e^{-\beta x}}{2} \right] \\
 &= e^{\alpha x} (K_1 \cosh \beta x + K_2 \sinh \beta x).
 \end{aligned}$$
43. $y = C_1 x^{-2} + C_2 x^4$.
 45. $y = C_1 x^2 + C_2 x^2 \ln x$.

Exercises 3.4

1. $z(x) = x^2 \ln x + \frac{1}{2}$; $y = C_1 x^2 + C_2 x^{-1} + x^2 \ln x + \frac{1}{2}$.
3. $z(x) = -x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2$; $y = C_1 x + C_2 x^2 - x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2$.
5. $z(x) = -(1 + x^2)$; $y = C_1 x + C_2 e^x - (1 + x^2)$.
7. $y = C_1 e^{-x} + C_2 e^{2x} - \frac{2}{3} x e^{-x}$.
9. $y = C_1 \cos 2x + C_2 \sin 2x + \frac{1}{4} \cos 2x \ln(\cos 2x) + \frac{1}{2} x \sin 2x$.
11. $y = C_1 e^x + C_2 x e^x - e^x \cos x$.
13. $y = C_1 e^{-2x} + C_2 x e^{-2x} - e^{-2x} \ln x$.
15. $y = C_1 \cos 3x + C_2 \sin 3x + \sin 3x \ln(\sec 3x + \tan 3x) - 1$.
17. $y = C_1 x + C_2 x^{-1} + x \ln x$.
19. $y = C_1 x + C_2 x \ln x + x^2$.

Exercises 3.5

1. $y = C_1 e^{-x} + C_2 e^{3x} - e^{2x}$.
3. $y = C_1 e^{-3x} + C_2 x e^{-3x} + \frac{1}{4} e^{3x}$.
5. $y = C_1 e^{-2x} + C_2 - \frac{1}{2} \cos 2x - \frac{1}{2} \sin 2x$.
7. $y = C_1 e^{-x/2} + C_2 e^{-x} + x^2 - 6x + 14 - \frac{9}{10} \cos x - \frac{3}{10} \sin x$.
9. $y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{1}{2} x + \frac{1}{4}$.
11. $y = C_1 e^{-2x} + C_2 e^{-4x} + \frac{3}{2} x e^{-2x}$.
13. $y = C_1 \cos 3x + C_2 \sin 3x + \frac{2}{3} + \frac{1}{162} (9x^2 - 6x + 1) e^{3x}$.
15. $y = e^x (C_1 \cos 2x + C_2 \sin 2x) + \frac{1}{10} e^{-x} \cos 2x + \frac{1}{20} e^{-x} \sin 2x$.
17. $y = e^x - \frac{1}{2} e^{-2x} - x - \frac{1}{2}$.
19. $y = \frac{13}{15} e^{-x} + \frac{1}{12} e^{2x} + \frac{1}{20} \cos 2x - \frac{3}{20} \sin 2x$.
21. $z = A + (Bx^2 + Cx)e^{-x} + D \cos 3x + E \sin 3x$.
23. $z = Ax^2 + Bx + C + Dx \cos x + Ex \sin x$.
25. $z = (Ax^3 + Bx^2)e^{2x} + Cx^2 + Dx + E + (Fx + G) \cos 2x + (Hx + I) \sin 2x$.
27. $z = Ae^{-x} + Bxe^{-x} \cos x + Cxe^{-x} \sin x + D$.
29. $y = C_1 e^{2x} + C_2 x e^{2x} + \frac{8}{25} \cos x + \frac{6}{25} \sin x + 3x e^{2x} \ln x$.
31. $y = C_1 \cos 3x + C_2 \sin 3x + \frac{3}{8} \cos x - \sin 3x \ln(\sec 3x + \tan 3x) + 1$.
33. $y_1 - y_2$ is a solution of the reduced equation $y'' + ay' + by = 0$ with $a, b > 0$. As shown in Exercises 3.3, Problem 38, $y_1 - y_2 \rightarrow 0$ as $x \rightarrow \infty$. If $a = 0, b > 0$, then all solutions of the reduced equation are bounded (Problem 39 (a), Exercises 3.3).

Exercises 3.6

1. The equation of motion is $y(t) = \sin(8t + \frac{1}{2}\pi)$. The amplitude is 1 and the frequency is $8/2\pi = 4/\pi$.
3. $\pm 2\pi A/T$.

5. $y = C_1 \cos \omega t + C_2 \sin \omega t =$

$$\sqrt{C_1^2 + C_2^2} \left(\frac{C_1}{\sqrt{C_1^2 + C_2^2}} \cos \omega t + \frac{C_2}{\sqrt{C_1^2 + C_2^2}} \sin \omega t \right) \quad (*)$$

Let $A = \sqrt{C_1^2 + C_2^2}$. Since $\left(\frac{C_1}{\sqrt{C_1^2 + C_2^2}}\right)^2 + \left(\frac{C_2}{\sqrt{C_1^2 + C_2^2}}\right)^2 = 1$ there is an angle ϕ_0 such that

$$\sin \phi_0 = \frac{C_1}{\sqrt{C_1^2 + C_2^2}} \quad \text{and} \quad \cos \phi_0 = \frac{C_2}{\sqrt{C_1^2 + C_2^2}}$$

(a) Substituting into (*), we get

$$y = C_1 \cos \omega t + C_2 \sin \omega t = A (\cos \omega t \sin \phi_0 + \sin \omega t \cos \phi_0) = A \sin(\omega t + \phi_0)$$

(b) $A \sin(\omega t + \phi_0) = A \cos(\omega t + \phi_0 - \frac{\pi}{2}) = A \cos(\omega t + \psi_0)$ where $\psi_0 = \phi_0 - \frac{1}{2}\pi$.

7. Assume that $r_1 > r_2$. If $C_1 = 0$ or $C_2 = 0$, then $y = C_1 e^{r_1 t} + C_2 e^{r_2 t}$ can never be zero. If both C_1 and C_2 are nonzero, then $C_1 e^{r_1 t} + C_2 e^{r_2 t} = 0$ implies $e^{(r_1 - r_2)t} = -\frac{C_2}{C_1}$. Since $e^{(r_1 - r_2)t}$ is an increasing function ($r_1 > r_2$), it can take the value $\frac{C_2}{C_1}$ at most once. By the same reasoning, $y'(t) = C_1 r_1 e^{r_1 t} + C_2 r_2 e^{r_2 t}$ can be zero at most once. Therefore the motion can change direction at most once.

9. Suppose $\gamma \neq \omega$. Set $z = A \cos \gamma t + B \sin \gamma t$ and use undetermined coefficients. The result is

$$z = \frac{F_0/m}{\omega^2 - \gamma^2} \cos \gamma t.$$

11. Suppose $\gamma = \omega$. Then a particular solution z has the form $z = At \cos \omega t + Bt \sin \omega t$. Substituting z into the equation, we get

$$A = 0, \quad B = \frac{F_0}{2\omega m}, \quad \text{and so} \quad z = \frac{F_0}{2\omega m} t \sin \omega t.$$

Exercises 3.7

1. $y = C_1 e^x + C_2 e^{2x} + C_3 e^{3x}$.

3. $y = C_1 e^{2x} + C_2 e^{-2x} + e^x [C_3 \cos 2x + C_4 \sin 2x]$.

5. $y = C_1 \cos x + C_2 \sin x + e^{2x} [C_3 \cos 3x + C_4 \sin 3x]$.

7. $y = C_1 + C_2 x + C_3 e^x + C_4 e^{-x} + C_5 \cos x + C_6 \sin x$.

9. $y = 2x - 1$.

11. $y = \frac{1}{5} e^x - \frac{1}{5} \cos 3x - \frac{1}{15} \sin x$

13. $y^{(4)} + 4y''' + 7y'' + 42y' + 90y = 0$.

15. $y^{(5)} - 2y^{(4)} - 2y''' - 2y'' - 3y' = 0$.

17. $y^{(5)} - 2y^{(4)} + y''' - 2y'' = 0$.

19. $y^{(4)} - y'' = 0$.

21. $y = C_1 e^{-x} + C_2 \cos x + C_3 \sin x + \frac{1}{4} e^x + 4.$
 23. $y = C_1 \cos x + C_2 \sin x + C_3 x \cos x + C_4 x \sin x + 6 + \frac{1}{9} \cos 2x.$
 25. $y = -\frac{9}{16} e^{2x} + \frac{9}{16} e^{-x} \cos x \sqrt{3} + \frac{77\sqrt{3}}{144} e^{-x} \sin x \sqrt{3} + \frac{1}{12} x e^{2x}$

CHAPTER 4

Exercises 4.1

1. $\frac{1}{s^2}.$
 3. $\frac{1}{s^2 + 1}.$
 5. $\frac{1}{2(s-1)} - \frac{1}{2(s+1)}.$
 7. $\frac{s-a}{(s-a)^2 + b^2}.$

Exercises 4.2

1. $\frac{3}{s} - \frac{2}{s^2} + \frac{2}{s^3}.$
 3. $\frac{3}{s} + \frac{4}{s-3} - \frac{2s}{s^2+4}.$
 5. $\frac{10}{s^3} - \frac{4}{(s+3)^2+4}.$
 7. $\frac{2s}{(s^2+1)^2} + \frac{2(s^2-4)}{(s^2+4)^2}.$
 9. $\sinh \beta x = \frac{e^{\beta x} - e^{-\beta x}}{2}; \quad \frac{1}{2} \left[\frac{1}{s-\beta} - \frac{1}{s+\beta} \right] = \frac{\beta}{s^2 - \beta^2}.$
 11. $\frac{1}{2} \left[\frac{1}{s-3} + \frac{1}{s-2} - \frac{1}{s-1} + \frac{1}{s-4} \right].$
 15. $Y(s) = \frac{1}{s-2}.$
 17. $Y(s) = \frac{2}{(s-2)(s+4)} - \frac{9}{(s^2+9)(s+4)} - \frac{3}{s+4}.$
 19. $Y(s) = \frac{2}{(s+3)^2}.$
 21. $Y(s) = \frac{3}{s(s-5)(s+3)} + \frac{4}{(s-5)(s+3)^2} + \frac{s-5}{(s-5)(s+3)}.$
 23. Set $g(x) = \int_0^x f(t) dt.$ Then $g'(x) = f(x)$ and $g(0) = 0.$

$$F(s) = \mathcal{L}[f(x)] = \mathcal{L}[g'(x)] = s\mathcal{L}[g(x)] - g(0) = s\mathcal{L}[g(x)].$$

$$\text{Therefore, } \mathcal{L}[g(x)] = \frac{1}{s} F(s).$$

Exercises 4.3

1. $f(x) = 6e^{-7x}.$

3. $f(x) = 2 \cos 5x + \frac{1}{5} \sin 5x.$
5. $f(x) = e^{-4x} \cos x.$
7. $f(x) = e^{-2x} \cos 2x + e^{-2x} \sin 2x.$
9. $f(x) = 2xe^{-2x} - e^x \cos x - e^x \sin x.$
11. $f(x) = \frac{1}{2} e^{-x} - \frac{1}{2} \cos x + \frac{1}{2} \sin x.$
13. $f(x) = \frac{1}{4} - \frac{1}{4} \cos 2x.$
15. $f(x) = \frac{1}{2} - e^x + \frac{3}{2} e^{-2x}.$
17. $f(x) = e^{2x} - 4e^x + 2x + 3.$
19. $y = \frac{2}{3} e^{-2x} + \frac{1}{3} e^x.$
21. $y = \frac{3}{2} e^{-x} - \frac{1}{2} \cos x + \frac{1}{2} \sin x.$
23. $y = e^x \sin x.$
25. $y = \frac{3}{4} e^{-x} + \frac{1}{4} e^x + \frac{1}{2} xe^x.$
27. $y = \frac{1}{4} e^x + xe^{-x} + x - 2.$
29. $y = e^{-2x} + e^x.$
31. $y = -\frac{1}{5} e^{-2x} \cos 2x - \frac{1}{10} e^{-2x} \sin 2x + \frac{1}{5} e^{-x}$
33. $\alpha = \frac{1}{4}.$
35. $\beta = -\frac{26}{5}.$
37. $y = \frac{7}{4} e^{2(x-1)} - 3e^{x-1} + \frac{1}{2} x + \frac{3}{4}.$

Exercises 4.4

1. $F(s) = 2e^{-5s} \frac{1}{s}.$
3. $F(s) = \frac{2}{s^2} - 2e^{-3s} \frac{1}{s^2} - 5e^{-3s} \frac{1}{s}.$
5. $f(x) = 0 + 5u(x-4); \quad F(s) = 5e^{-4s} \frac{1}{s}.$
7. $f(x) = 0 + (x-2)u(x-2) + 2u(x-2); \quad F(s) = e^{-2s} \frac{1}{s^2} + 2e^{-2s} \frac{1}{s}.$
9. $f(x) = 8 + 2(x-5)u(x-5) + 2u(x-5); \quad F(s) = \frac{8}{s} + 2e^{-5s} \frac{1}{s^2} + 2e^{-5s} \frac{1}{s}.$
11. $f(x) = x^2 - (x-3)^2 u(x-3) - 3(x-3)u(x-3); \quad F(s) = \frac{2}{s^3} - e^{-3s} \frac{2}{s^3} - 3e^{-3s} \frac{1}{s^2}.$
13. $f(x) = x - 1 - (x-2)u(x-2) - u(x-2) + e^{-2} e^{-(x-2)} u(x-2);$
 $F(s) = \frac{1}{s^2} - \frac{1}{s} - e^{-2s} \frac{1}{s^2} - e^{-2s} \frac{1}{s} + e^{-2} e^{-2s} \frac{1}{s+1}.$
15. $f(x) = \sin 2x - \sin 2(x-\pi)u(x-\pi) + (x-\pi)u(x-\pi) + \pi u(x-\pi);$
 $F(s) = \frac{2}{s^2+4} - e^{-\pi s} \frac{2}{s^2+4} + e^{-\pi s} \frac{1}{s^2} + \pi e^{-\pi s} \frac{1}{s}.$
17. $f(x) = x - (x-2)u(x-2) - 2u(x-2) + (x-4)^2 u(x-4);$
 $F(s) = \frac{1}{s^2} - e^{-2s} \frac{1}{s^2} - 2e^{-2s} \frac{1}{s} + e^{-4s} \frac{2}{s^3}.$
19. $f(x) = 1 - u(x-2) + (x-2)u(x-2) - (x-4)u(x-4) - 2u(x-4) + e^{-(x-4)} u(x-4);$
 $F(s) = \frac{1}{s} - e^{-2s} \frac{1}{s} + e^{-2s} \frac{1}{s^2} - e^{-4s} \frac{1}{s^2} - 2e^{-4s} \frac{1}{s} + e^{-4s} \frac{1}{s+1}.$

Exercises 4.5

1. $f(x) = 2 + 2u(x-3) = \begin{cases} 2, & 0 \leq x < 3 \\ 4, & x \geq 3 \end{cases}$.
3. $f(x) = \sin x - \sin x u(x-\pi) = \begin{cases} \sin x, & 0 \leq x < \pi \\ 0, & x \geq \pi. \end{cases}$
5. $f(x) = \cos x - \cos x u(x-\pi) + \sin x u(x-\pi) = \begin{cases} \cos x, & 0 \leq x < \pi \\ \sin x, & x \geq \pi. \end{cases}$
7. $f(x) = \cos \pi x - \frac{1}{\pi} \sin \pi x u(x-2) = \begin{cases} \cos \pi x, & 0 \leq x < 2 \\ \cos \pi x - \frac{1}{\pi} \sin \pi x & x \geq 2. \end{cases}$
9. $f(x) = 3e^{3(x-3)}u(x-3) - 2e^{2(x-3)}u(x-3) = \begin{cases} 0, & 0 \leq x < 3 \\ 3e^{3(x-3)} - 2e^{2(x-3)}, & x \geq 3. \end{cases}$
11. $f(x) = 2 + e^{(x-1)}u(x-1) - e^2 e^{(x-2)}u(x-2) = \begin{cases} 2, & 0 \leq x < 1 \\ 2 + e^{(x-1)}, & 1 \leq x < 2 \\ 2 + e^{(x-1)} - e^x, & x \geq 2. \end{cases}$
13. $f(x) = \cos 2x - 1 + u(x-2) - \cos 2(x-2)u(x-2) = \begin{cases} \cos 2x - 1, & 0 \leq x < 2 \\ \cos 2x - \cos 2(x-2) & x \geq 2. \end{cases}$
15. $f(x) = 2e^\pi e^{-2x} \cos 3x u(x-\pi/2) - e^\pi e^{-2x} \sin 3x u(x-\pi/2)$
 $= \begin{cases} 0, & 0 \leq x < \pi/2 \\ 2e^\pi e^{-2x} \cos 3x - e^\pi e^{-2x} \sin 3x & x \geq \pi/2. \end{cases}$

Exercises 4.6

1. $y = -\frac{1}{2} + \frac{5}{2}e^{2x} + u(x-1) \left[-\frac{1}{2} + \frac{1}{2}e^{2(x-1)}\right]$
 $= \begin{cases} -\frac{1}{2} + \frac{5}{2}e^{2x}, & 0 \leq x < 1 \\ -1 + \frac{5}{2}e^{2x} + \frac{1}{2}e^{2(x-1)}, & x \geq 1 \end{cases}$
3. $y = 1 - \cos x + \sin x - u(x-1)[\cos(x-1) - 1]$
 $= \begin{cases} 1 - \cos x + \sin x, & 0 \leq x < \pi \\ 2 \cos x, & x \geq \pi \end{cases}$
5. $y = 1 - e^{-x} - xe^{-x} + u(x-2) \left[x - 4 + xe^{-(x-2)}\right]$
 $= \begin{cases} 1 - e^{-x} - xe^{-x}, & 0 \leq x < 2 \\ -3 - e^{-x} - xe^{-x} + x + xe^{-(x-2)}, & x \geq 2 \end{cases}$
7. $y = -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x + u(x-1) \left[\frac{1}{3} + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{x-1}\right]$
 $= \begin{cases} -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x, & 0 \leq x < 1 \\ -\frac{1}{6}e^{3x} + \frac{1}{2}e^x + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{(x-1)}, & x \geq 1 \end{cases}$
9. $y = \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + u(x-1) \left[xe^{-(x-1)} - 1\right]$.

$$= \begin{cases} \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x}, & 0 \leq x < 1 \\ \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + xe^{-(x-1)} - 1, & x \geq 1 \end{cases}$$

CHAPTER 5

Exercises 5.2

1. $x = 4, y = 1$.
3. $x = 4 - 2a, y = a, a$ any real number.
5. $x = -3, y = 1$.
7. No solution.
9. $x = 2a - 3, y = a, a$ any real number.
11. $x = \frac{3}{7}a + 1, y = \frac{5}{7}a - 1, z = a, a$ any real number; a line in 3-space.
13. No solution.
15. $x = 2, y = 1, z = 1$; a point in 3-space.

Exercises 5.3

1. matrix of coefficients: 3; augmented matrix: 3; $x = 5, y = 3, z = -1$.
3. matrix of coefficients: 2; augmented matrix: 2;

$$x = 4 - 2a, y = a, z = -2, \quad a \text{ any real number.}$$

5. matrix of coefficients: 3; augmented matrix: 3;

$$x_1 = -1, x_2 = -1 - 2a, x_3 = 3 + a, x_4 = a, \quad a \text{ any real number.}$$

7. matrix of coefficients: 3; augmented matrix: 3;

$$x_1 = 8 + 2a - 3b, x_2 = a, x_3 = 3 - 1 - 2b, x_4 = b, x_5 = -3,$$

a, b any real numbers.

9. $x = 2, y = 5$.
11. $x = -3 - a, y = 2 + 2a, z = a, a$ any real number.
13. $x = \frac{10}{7}, y = \frac{2}{7}, z = \frac{3}{2}$.
15. $x_1 = 11 - 2a + b, x_2 = a, x_3 = 3 - b, x_4 = b, a, b$ any real numbers.
17. $x_1 = -2, x_2 = -5, x_3 = -1, x_4 = 5$.
19. $x_1 = 3 - 2a, x_2 = a, x_3 = 2, x_4 = 1$.
21. (i) $k \neq -3, 2$ (ii) $k = -3$ (iii) $k = 2$.
23. (i) $a \neq -3, 3$ (ii) $a = -3$ (iii) $a = 3$.
25. The system has at least one solution if $b = a, a$ any real number.
27. (a) No (b) No (c) Yes
29. $y = \frac{4}{3}e^x - \frac{1}{3}e^{-2x}$.

31. $y = -\frac{7}{2} + \frac{1}{2}e^{2x} + 4e^{-x/2}$.

Exercises 5.4

1. Yes.
3. No. The leading 1 in the third row is not to the right of the leading 1 in the second row.
5. No. The leading 1 in the last column is not the only nonzero in its column.
7. Yes
9. $x = 10, y = -9, z = -7$
11. $x = -3 - a, y = 2 + 2a, z = a, a$ any real number.
13. $x_1 = 11 - 2a + b, x_2 = a, x_3 = 3 - b, x_4 = b, a, b$ any real numbers.
15. $x_1 = 7 - 2a - b, x_2 = 1 + 3a - 4b, x_3 = a, x_4 = b, a, b$ any real numbers.
17. $x = y = 0$.
19. $x = y = z = 0$.
21. $x_1 = 2a - b, x_2 = -a + 4b, x_3 = a, x_4 = b, a, b$ any real numbers.
23. $x_1 = x_2 = x_3 = x_4 = 0$.
25. Consider the system

$$\begin{aligned} x + y &= 0 \\ 2x + 2y &= 0 \\ 3x + 3y &= 0 \end{aligned}$$

This system has the solutions $x = -a, y = a, a$ any real number.

27. $a = 4$.
29. (a) Solution set $\mathcal{S} : x = 1 + a, y = -1 - a, z = a, a$ any real number.

(b) $\mathcal{S} : \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} + a \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$.

Exercises 5.5

1. (a) $\begin{pmatrix} 0 & 4 \\ 3 & 5 \\ 1 & -1 \end{pmatrix}$. (c) $\begin{pmatrix} 8 & -4 \\ 1 & 4 \end{pmatrix}$. (e) $\begin{pmatrix} 2 & 8 \\ 2 & 8 \\ 5 & -3 \end{pmatrix}$.
3. (a) $\begin{pmatrix} -4 & -3 \\ 28 & -6 \\ -20 & 24 \end{pmatrix}$. (c) Not defined. (e) $\begin{pmatrix} 1 & 3 \\ -3 & -12 \\ -41 & 21 \end{pmatrix}$.
5. (a) $c_{32} = 2$ (b) $c_{13} = 34$ (c) $d_{21} = 5$ (d) $d_{22} = 1$.
7. (a) $d_{22} = 6$ (b) $d_{12} = -4$ (c) $d_{23} = -18$.
11. (a) $AB = \begin{pmatrix} 4 & 7 & 10 \\ 0 & -5 & -14 \end{pmatrix}$, BA not defined.

$$(b) AC = \begin{pmatrix} 14 & 5 \\ -2 & -3 \end{pmatrix}, \quad CA = \begin{pmatrix} -1 & 14 \\ 5 & 12 \end{pmatrix}.$$

$$(c) AD = DA = \begin{pmatrix} 4 & 4 \\ -2 & 2 \end{pmatrix}.$$

$$13. A(BD) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}$$

$$(AB)D = \begin{pmatrix} 0 & -2 \\ 3 & 5 \end{pmatrix} \begin{pmatrix} -2 & 3 & -2 \\ 3 & 0 & -4 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}.$$

$$15. (a) 3 \times 3 \quad (c) \text{ Does not exist} \quad (e) 2 \times 3.$$

Exercises 5.6

$$1. A^{-1} = \begin{pmatrix} 1/2 & 0 \\ -3/2 & 1 \end{pmatrix}.$$

$$3. A^{-1} = \begin{pmatrix} -2 & 1 \\ 3/2 & -1/2 \end{pmatrix}.$$

$$5. A^{-1} = \begin{pmatrix} -11 & 2 & 2 \\ -4 & 0 & 1 \\ 6 & -1 & -1 \end{pmatrix}.$$

7. No inverse.

9. No inverse.

11. $\det A = \pm 1$.

13. $x = 5, y = 0$.

15. $x = \frac{9}{2}, y = -5$.

17. $x = \frac{7}{9}, y = \frac{1}{3}, z = -\frac{5}{9}$.

19. -31 .

21. -45 .

23. 30 .

25. -21 .

27. -18 .

29. 26 .

31. $x = 0, 1, -3$.

33. $y = -\frac{25}{37}$.

35. Cramer's rule does not apply.

37. $x = 0$.

39. $\lambda = -4, 7$.

Exercises 5.7

3. Dependent; $(-4, 8, 9) = 2(1, -2, 3) + 3(-2, 4, 1)$.

5. Dependent; $(-2, 6, 3) = (1, -1, 3) + 2(0, 2, 3) - 3(1, -1, 2)$.
7. Dependent; $(7, -4, 1) = 3(1, -2, 1) + 2(2, 1, -1)$.
9. Dependent; $(4, -2, 0, 2) = 2(2, -1, 0, 1)$.
11. $b \neq -\frac{1}{3}$.
13. $b = 0, -7$.
17. No; a linearly dependent set can have linearly independent subsets. For example, $\{(1, -2, 3), (-2, 4, 1)\}$ is a linearly independent subset of $\{(1, -2, 3), (-2, 4, 1)\}, (-4, 8, 9)$.
19. $W(x) = -a$; linearly independent.
21. $W(x) = -2x^{-6}$; linearly independent.
23. $W(x) = e^{2x}(x - 2)$; linearly independent.
25. (a) False (b) True (c) True.

Exercises 5.8

1. $2, \begin{pmatrix} 1 \\ 0 \end{pmatrix}; 3, \begin{pmatrix} 1 \\ -1 \end{pmatrix}$.
3. $-1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}; 4, \begin{pmatrix} 2 \\ 3 \end{pmatrix}$.
5. $1, 1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}$.
7. $2, 2, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$.
9. $2 + i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} + i \begin{pmatrix} 0 \\ -1 \end{pmatrix}; 2 - i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} - i \begin{pmatrix} 0 \\ -1 \end{pmatrix}$.
11. $8, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}; 1, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}; 2, \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$.
13. $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$.
15. $1 + i, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + i \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}; 1 - i, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - i \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}; 0, \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$.
17. $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$.
19. $7, \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}; -2, \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix}; -2, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$.

$$21. \quad 2, \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}; \quad 2, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}; \quad 6, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}; \quad 4, \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}.$$

CHAPTER 6

Exercises 6.1

$$1. \quad \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -3 & t \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 0 \\ \sin 2t \end{pmatrix}.$$

$$3. \quad \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ e^t \end{pmatrix}.$$

$$5. \quad \begin{aligned} x'_1 &= 2x_1 - x_2 + e^{2t} \\ x'_2 &= 3x_1 + 2e^{-t} \end{aligned}.$$

$$7. \quad \begin{aligned} x'_1 &= 2x_1 + 3x_2 - x_3 + e^t \\ x'_2 &= -2x_1 + x_3 + 2e^{-t} \\ x'_3 &= 2x_1 + 3x_2 + e^{2t} \end{aligned}.$$

$$9. \quad \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} \sin t \\ -2 \cos t \end{pmatrix}.$$

$$11. \quad \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 2 & 1 & 3 \\ 1 & -3 & 0 \\ 2 & -1 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 3e^{2t} \\ -2 \cos t \\ t \end{pmatrix}.$$

Exercises 6.2

1. Independent

3. Independent

5. Dependent

7. Dependent

9. Dependent

11. (c) $\mathbf{x}(t) = c_1 \mathbf{u} + c_2 \mathbf{v}$, where c_1, c_2 are arbitrary constants.

$$(d) \quad \mathbf{x}(t) = -2 \begin{pmatrix} e^{2t} \\ e^{2t} \end{pmatrix} + \begin{pmatrix} 3e^{3t} \\ 2e^{3t} \end{pmatrix}.$$

$$13. \quad (b) \quad \mathbf{x}(t) = \begin{pmatrix} -4te^{-t} \\ 2 - e^{-t} \\ 2 \end{pmatrix}.$$

15. The system is equivalent to $y''' + y'' - 4y' - 4y = 0$. A fundamental set of solutions of this equation is $\{y_1 = e^{-2t}, y_2 = e^{-t}, y_3 = e^{2t}\}$. Therefore, linearly independent solutions of the given system are:

$$\mathbf{x}_1 = \begin{pmatrix} e^{-2t} \\ -2e^{-2t} \\ 4e^{-2t} \end{pmatrix} = e^{-2t} \begin{pmatrix} 1 \\ -2 \\ 4 \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} e^{-t} \\ -e^{-t} \\ e^{-t} \end{pmatrix} = e^{-t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} e^{2t} \\ 2e^{2t} \\ 4e^{2t} \end{pmatrix} = e^{2t} \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix}.$$

17. The system is equivalent to $y'' + \frac{6}{t}y' + \frac{6}{t^2}y = 0$. A fundamental set of solutions of this equation is $\{y_1 = t^{-3}, y_2 = t^{-2}\}$. Therefore, linearly independent solutions of the given system are:

$$\mathbf{x}_1 = \begin{pmatrix} t^{-3} \\ -3t^{-4} \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} t^{-2} \\ -2t^{-3} \end{pmatrix}.$$

Exercises 6.3

1. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-3t} \begin{pmatrix} 4 \\ -1 \end{pmatrix}.$
3. $\mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + C_2 e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}; \quad \mathbf{x}(t) = -3e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + 4e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$
5. $\mathbf{x}(t) = C_1 e^{5t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 2 \\ -1 \end{pmatrix}.$
7. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$
9. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}; \quad \mathbf{x}(t) = -e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + 2e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}.$
11. $\mathbf{x}(t) = C_1 e^{10t} \begin{pmatrix} 9 \\ 7 \\ 4 \end{pmatrix} + C_2 e^{6t} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + C_3 e^{5t} \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}.$
13. (a) $\mathbf{x}' = \begin{pmatrix} 0 & 1 \\ -b & -a \end{pmatrix} \mathbf{x};$ (b) $\lambda^2 + a\lambda + b = 0;$ (c) They are the same.

Exercises 6.4

1. $\mathbf{x}(t) = C_1 \left[\cos 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right] + C_2 \left[\cos 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} \right].$
3. $\mathbf{x}(t) = C_1 e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + C_2 \left[e^t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right].$

$$5. \quad \mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} + C_2 \left[e^{-t} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} \right].$$

$$7. \quad \mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + C_3 e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix};$$

$$\mathbf{x}(t) = e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$$

$$9. \quad \mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} + C_2 \left[\cos 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] + C_3 \left[\cos 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} \right].$$

$$11. \quad \mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + C_3 e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix};$$

$$\mathbf{x}(t) = \frac{7}{2} e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \frac{9}{2} e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \frac{3}{2} e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}$$

$$13. \quad \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_3 \left[e^t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$$

$$15. \quad \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \left[\cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + C_3 e^t \left[\cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right];$$

$$\mathbf{x}(t) = e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + e^t \left[\cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + 3e^t \left[\cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$$

$$17. \quad \mathbf{x}(t) = C_1 e^{6t} \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} + C_2 e^{-2t} \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} + C_3 e^{-2t} \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix}.$$