

TEST #2 - version 2

Please, show your work, justify every step and write legibly. When you are done, scan, save the file as LASTNAME_FIRSTNAME_T2.pdf and email to dlabate@math.uh.edu or dlabate@uh.edu.

NOTE: You need to send your email before 11:30AM on March 25 to receive credit.

• (1) [8 Pts]

- (a) Compute the sine series of $f(x) = \sin x$, on the interval $0 \leq x \leq \pi$.
- (b) Does the sine series of f converge uniformly to f in $[0, \pi]$? If not, where does it fail to converge? Justify.
- (c) Without computing the Fourier coefficients, set up the computation of the cosine series of $f(x) = \sin x$, on the interval $0 \leq x \leq \pi$.
- (d) Does the cosine series of f converge uniformly to f in $[0, \pi]$? If not, where does it fail to converge? Justify.

• (2) [8 Pts] Consider the function

$$f(x) = \begin{cases} 1 & \text{if } -\pi \leq x \leq -\frac{\pi}{4} \\ 0 & \text{if } -\frac{\pi}{4} < x < \frac{\pi}{4} \\ -1 & \text{if } \frac{\pi}{4} \leq x \leq \pi \end{cases}$$

- (a) Sketch the function.
- (b) Compute the Fourier series of f , valid in the interval $[-\pi, \pi]$. (Hint: Take advantage of even/odd properties to reduce calculations.)
- (c) Does the Fourier series converge to f uniformly on $[-\pi \leq x \leq \pi]$? If not, where does it fail to converge? Justify.

• (3) [4 Pts]

- (a) Compute the Fourier series of $f(x) = \sin^2 x$ valid in the interval $[-\pi, \pi]$
- (b) Compute the Complex Fourier series of $f(x) = \sin^2 x$ valid in the interval $[-\pi, \pi]$ [Hint: there is no need of integration. You can derive it using the result of part (a).]