

TEST #3 - version 2

Please, show your work, justify every step and write legibly. When you are done, scan, save the file as LASTNAME.FIRSTNAME\_T3.pdf and email to [dlabate@math.uh.edu](mailto:dlabate@math.uh.edu) or [dlabate@uh.edu](mailto:dlabate@uh.edu). NOTE: You need to send your email before 11:30AM on April 20 to receive credit.

• (1) [3 Pts] Let  $A > 0$  be a fixed number. Does the following functions define a causal filter? Justify your answer.

$$(a) h_1(t) = \begin{cases} 0 & \text{if } t < 0; \\ e^{-At} & \text{if } t \geq 0. \end{cases}$$

$$(b) h_2(t) = \begin{cases} e^{At} & \text{if } t < 0; \\ e^{-At} & \text{if } t \geq 0. \end{cases}$$

• (2) [9 Pts] Let  $f(t) = \begin{cases} 1 & \text{if } -\pi \leq t \leq \pi; \\ 0 & \text{otherwise.} \end{cases}$

We found that its Fourier transform is  $\hat{f}(\omega) = \sqrt{\frac{2}{\pi}} \frac{\sin(\pi\omega)}{\omega}$ .

(a) Use the Fourier transform of  $f$  and the properties of the Fourier transform to compute the Fourier transform  $\hat{c}(\omega)$  of the function  $c(t) = (f * f)(t)$

(b) Use the Fourier transform of  $f$  and the properties of the Fourier transform to compute the Fourier transform of

$$h(t) = \begin{cases} 1 & \text{if } -1 \leq t \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

(c) Use the Fourier transform of  $h$  and the properties of the Fourier transform to compute the Fourier transform of

$$g(t) = \begin{cases} t^2 & \text{if } -1 \leq t \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

• (3) [6 Pts] Let

$$f(t) = \begin{cases} 1 & 0 \leq t \leq 2 \\ 0 & \text{otherwise;} \end{cases} \quad g(t) = \begin{cases} t & 0 \leq t \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

(a) Compute  $h(t) = (f * g)(t)$ .

(b) Sketch the graph of  $f, g, h$  over the interval  $[-1, 4]$ .