

# Lecture 2


## Section 7.2 The Logarithm Function, Part I

Jiwen He

Department of Mathematics, University of Houston

[jiwenhe@math.uh.edu](mailto:jiwenhe@math.uh.edu)

<http://math.uh.edu/~jiwenhe/Math1432>



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{rs} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .  
To calculate  $2^6$ , we do in our head or on a paper

$$2 \times 2 \times 2 \times 2 \times 2 \times 2,$$

but what does the computer actually do when we type

$$2^6$$

- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r-s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r-s} = (x^r)^s,$$

$$\frac{d}{dx}x^r = rx^{r-1}, \quad \int x^r dx = \frac{1}{r+1}x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .

$f(x) = x^{\frac{1}{n}}$  is the inverse function of  $g(x) = x^n$  for  $x > 0$ .

$$\Rightarrow g \circ f(x) = \left(x^{\frac{1}{n}}\right)^n = x.$$

- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ?$$

and

$$x^r = ? \text{ for } r \text{ real.}$$





# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$

$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know yet that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What We Do/Don't Know About $f(x) = x^r$ ?

We know that:

- For  $r = n$  positive integer,  $f(x) = x^n = \overbrace{x \cdot x \cdots x}^{n \text{ times}}$ .
- For  $r = 0$ ,  $f(x) = x^0 = 1$ .
- For  $r = -n$ ,  $f(x) = \left(\frac{1}{x}\right)^n$ ,  $x \neq 0$ .  $\Rightarrow x^{-1} = \frac{1}{x}$ .
- For  $r = \frac{p}{q}$  rational,  $f(x) = y$ ,  $x > 0$ , where  $y^q = x^p$ .
- Properties ( $r$  and  $s$  rational)

$$x^{r+s} = x^r \cdot x^s, \quad x^{r \cdot s} = (x^r)^s,$$


$$\frac{d}{dx} x^r = r x^{r-1}, \quad \int x^r dx = \frac{1}{r+1} x^{r+1} + C, \quad r \neq -1.$$

We DO NOT know **yet** that:

$$\int x^{-1} dx = \int \frac{1}{x} dx = ? \quad \text{and} \quad x^r = ? \text{ for } r \text{ real.}$$



# What is the Natural Log Function?



## Definition

The function


$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0,$$

is called the **natural logarithm function**.

- $\ln 1 = 0$ .
- $\ln x < 0$  for  $0 < x < 1$ ,  $\ln x > 0$  for  $x > 1$ .
- $\frac{d}{dx}(\ln x) = \frac{1}{x} > 0 \Rightarrow \ln x$  is increasing.
- $\frac{d^2}{dx^2}(\ln x) = -\frac{1}{x^2} < 0 \Rightarrow \ln x$  is concave down.



# What is the Natural Log Function?



## Definition

The function


$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0,$$

is called the **natural logarithm function**.

- $\ln 1 = 0$ .
- $\ln x < 0$  for  $0 < x < 1$ ,  $\ln x > 0$  for  $x > 1$ .
- $\frac{d}{dx}(\ln x) = \frac{1}{x} > 0 \Rightarrow \ln x$  is increasing.
- $\frac{d^2}{dx^2}(\ln x) = -\frac{1}{x^2} < 0 \Rightarrow \ln x$  is concave down.



# What is the Natural Log Function?



## Definition

The function


$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0,$$

is called the **natural logarithm function**.

- $\ln 1 = 0$ .
- $\ln x < 0$  for  $0 < x < 1$ ,  $\ln x > 0$  for  $x > 1$ .
- $\frac{d}{dx}(\ln x) = \frac{1}{x} > 0 \Rightarrow \ln x$  is increasing.
- $\frac{d^2}{dx^2}(\ln x) = -\frac{1}{x^2} < 0 \Rightarrow \ln x$  is concave down.



# What is the Natural Log Function?



## Definition

The function


$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0,$$

is called the **natural logarithm function**.

- $\ln 1 = 0$ .
- $\ln x < 0$  for  $0 < x < 1$ ,  $\ln x > 0$  for  $x > 1$ .
- $\frac{d}{dx}(\ln x) = \frac{1}{x} > 0 \Rightarrow \ln x$  is increasing.
- $\frac{d^2}{dx^2}(\ln x) = -\frac{1}{x^2} < 0 \Rightarrow \ln x$  is concave down.



# What is the Natural Log Function?



## Definition

The function


$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0,$$

is called the **natural logarithm function**.

- $\ln 1 = 0$ .
- $\ln x < 0$  for  $0 < x < 1$ ,  $\ln x > 0$  for  $x > 1$ .
- $\frac{d}{dx}(\ln x) = \frac{1}{x} > 0 \Rightarrow \ln x$  is increasing.
- $\frac{d^2}{dx^2}(\ln x) = -\frac{1}{x^2} < 0 \Rightarrow \ln x$  is concave down.



# Example 1: $\ln x = 0$ and $(\ln x)' = 1$ at $x = 1$



## Exercise 7.2.23

Show that

$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = 1.$$

Proof.

$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = \lim_{x \rightarrow 1} \frac{\ln x - \ln 1}{x - 1} = \frac{d}{dx}(\ln x) \Big|_{x=1} = \frac{1}{x} \Big|_{x=1} = 1.$$




The limit has the **indeterminate form**  $\left(\frac{0}{0}\right)$  and is interpreted here in terms of the **derivative** of  $\ln x$ .





# Example 1: $\ln x = 0$ and $(\ln x)' = 1$ at $x = 1$



## Exercise 7.2.23

Show that

$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = 1.$$

Proof.


$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = \lim_{x \rightarrow 1} \frac{\ln x - \ln 1}{x - 1} = \frac{d}{dx}(\ln x) \Big|_{x=1} = \frac{1}{x} \Big|_{x=1} = 1.$$



The limit has the **indeterminate form**  $\left(\frac{0}{0}\right)$  and is interpreted here in terms of the **derivative** of  $\ln x$ .



# Example 1: $\ln x = 0$ and $(\ln x)' = 1$ at $x = 1$



## Exercise 7.2.23

Show that

$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = 1.$$

Proof.


$$\lim_{x \rightarrow 1} \frac{\ln x}{x - 1} = \lim_{x \rightarrow 1} \frac{\ln x - \ln 1}{x - 1} = \frac{d}{dx}(\ln x) \Big|_{x=1} = \frac{1}{x} \Big|_{x=1} = 1.$$



The limit has the **indeterminate form**  $\left(\frac{0}{0}\right)$  and is interpreted here in terms of the **derivative** of  $\ln x$ .



# Example 2: $\ln x$ and $x - 1$



## Exercise 7.2.24(a)

Show that

$$\frac{x-1}{x} \leq \ln x \leq x-1, \quad \forall x > 0. \quad (1)$$

### Proof.


- By the mean-value theorem,  $\exists c$  between 1 and  $x$  s.t.

$$\ln x = \int_1^x \frac{1}{t} dt = \frac{1}{c}(x-1).$$

- If  $x > 1$ , then  $\frac{1}{x} < \frac{1}{c} < 1$  and  $x-1 > 0$  so (1) holds.
- If  $0 < x < 1$ , then  $1 < \frac{1}{c} < \frac{1}{x}$  and  $x-1 < 0$  so (1) holds.  $\square$



# Example 2: $\ln x$ and $x - 1$



## Exercise 7.2.24(a)

Show that

$$\frac{x-1}{x} \leq \ln x \leq x-1, \quad \forall x > 0. \quad (1)$$

**Proof.**


- By the mean-value theorem,  $\exists c$  between 1 and  $x$  s.t.

$$\ln x = \int_1^x \frac{1}{t} dt = \frac{1}{c}(x-1).$$

- If  $x > 1$ , then  $\frac{1}{x} < \frac{1}{c} < 1$  and  $x-1 > 0$  so (1) holds.
- If  $0 < x < 1$ , then  $1 < \frac{1}{c} < \frac{1}{x}$  and  $x-1 < 0$  so (1) holds.  $\square$



# Example 2: $\ln x$ and $x - 1$



## Exercise 7.2.24(a)

Show that

$$\frac{x-1}{x} \leq \ln x \leq x-1, \quad \forall x > 0. \quad (1)$$

**Proof.**


- By the mean-value theorem,  $\exists c$  between 1 and  $x$  s.t.

$$\ln x = \int_1^x \frac{1}{t} dt = \frac{1}{c}(x-1).$$

- If  $x > 1$ , then  $\frac{1}{x} < \frac{1}{c} < 1$  and  $x-1 > 0$  so (1) holds.
- If  $0 < x < 1$ , then  $1 < \frac{1}{c} < \frac{1}{x}$  and  $x-1 < 0$  so (1) holds.  $\square$



# Example 2: $\ln x$ and $x - 1$



## Exercise 7.2.24(a)

Show that

$$\frac{x-1}{x} \leq \ln x \leq x-1, \quad \forall x > 0. \quad (1)$$

### Proof.


- By the mean-value theorem,  $\exists c$  between 1 and  $x$  s.t.

$$\ln x = \int_1^x \frac{1}{t} dt = \frac{1}{c}(x-1).$$

- If  $x > 1$ , then  $\frac{1}{x} < \frac{1}{c} < 1$  and  $x-1 > 0$  so (1) holds.
- If  $0 < x < 1$ , then  $1 < \frac{1}{c} < \frac{1}{x}$  and  $x-1 < 0$  so (1) holds.  $\square$



# Example 3: $\ln n$ and Harmonic Number




## Exercise 7.2.25(a)

Show that for  $n \geq 2$

$$\frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} < \ln n < 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n-1}.$$



# Example 3: In $n$ and Harmonic Number




## Proof.

Let  $P = \{1, 2, \dots, n\}$  be a partition of  $[1, n]$ . Then

$$L_f(P) = \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} < \int_1^n \frac{1}{t} dt < 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-1} = U_f(P).$$






Example 4: Euler's Constant  $\gamma$ 

## Exercise 7.2.25(c)

Show that

$$\frac{1}{2} < \gamma = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n-1} - \ln n \right) < 1.$$



Example 4: Euler's Constant  $\gamma$ 


## Proof.

- The sum of the shaded areas is given by

$$S_n = U_f(P) - \int_1^n \frac{1}{t} dt = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n-1} - \ln n.$$



# Example 4: Euler's Constant $\gamma$




## Proof. (cont.)

- The sum of the areas of the triangles formed by connecting the points  $(1, 1), \dots, (n, \frac{1}{n})$  is

$$T_n = \frac{1}{2} \cdot 1 \left[ \left(1 - \frac{1}{2}\right) + \dots + \left(\frac{1}{n-1} - \frac{1}{n}\right) \right] = \frac{1}{2} \left(1 - \frac{1}{n}\right).$$



# Example 4: Euler's Constant $\gamma$




Proof. (cont.)

- The sum of the areas of the indicated rectangles is

$$R_n = 1 \left[ \left( 1 - \frac{1}{2} \right) + \cdots + \left( \frac{1}{n-1} - \frac{1}{n} \right) \right] = 1 - \frac{1}{n}.$$



# Example 4: Euler's Constant $\gamma$



Proof. (cont.)

- Since  $T_n < S_n < R_n$ ,

$$\frac{1}{2} \left( 1 - \frac{1}{n} \right) < 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n-1} - \ln n < 1 - \frac{1}{n}.$$

Letting  $n \rightarrow \infty$  we have  $\frac{1}{2} < \gamma < 1$ .



# Basic Property: $\ln(xy) = \ln x + \ln y$

## Lemma

$$\ln(xy) = \ln x + \ln y, \quad x > 0, y > 0.$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln(xy) = \frac{1}{xy} y = \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (\ln x + \ln y) = \frac{1}{x}.$$

- Then

$$\ln(xy) = \ln x + \ln y + C$$

for some constant  $C$ . At  $x = 1$ , both sides take the same value of  $\ln y$ , thus  $C = 0$ . □



# Basic Property: $\ln(xy) = \ln x + \ln y$

## Lemma

$$\ln(xy) = \ln x + \ln y, \quad x > 0, y > 0.$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln(xy) = \frac{1}{xy} y = \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (\ln x + \ln y) = \frac{1}{x}.$$

- Then

$$\ln(xy) = \ln x + \ln y + C$$

for some constant  $C$ . At  $x = 1$ , both sides take the same value of  $\ln y$ , thus  $C = 0$ . □



# Basic Property: $\ln(xy) = \ln x + \ln y$

## Lemma

$$\ln(xy) = \ln x + \ln y, \quad x > 0, y > 0.$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln(xy) = \frac{1}{xy} y = \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (\ln x + \ln y) = \frac{1}{x}.$$

- Then

$$\ln(xy) = \ln x + \ln y + C$$

for some constant  $C$ . At  $x = 1$ , both sides take the same value of  $\ln y$ , thus  $C = 0$ . □





# Basic Property: $\ln(xy) = \ln x + \ln y$

## Lemma

$$\ln(xy) = \ln x + \ln y, \quad x > 0, y > 0.$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln(xy) = \frac{1}{xy} y = \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (\ln x + \ln y) = \frac{1}{x}.$$

- Then

$$\ln(xy) = \ln x + \ln y + C$$

for some constant  $C$ . At  $x = 1$ , both sides take the same value of  $\ln y$ , thus  $C = 0$ . □



# Basic Property: $\ln x^r = r \ln x$

## Lemma

$$\ln x^r = r \ln x \quad (r \text{ rational}).$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln x^r = \frac{1}{x^r} \frac{d}{dx} x^r = \frac{1}{x^r} r x^{r-1} = r \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (r \ln x) = r \frac{1}{x}.$$

- Then

$$\ln x^r = r \ln x + C$$

for some constant  $C$ . At  $x = 1$ , both sides are zero, thus  $C = 0$ .



# Basic Property: $\ln x^r = r \ln x$

## Lemma

$$\ln x^r = r \ln x \quad (r \text{ rational}).$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln x^r = \frac{1}{x^r} \frac{d}{dx} x^r = \frac{1}{x^r} r x^{r-1} = r \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (r \ln x) = r \frac{1}{x}.$$

- Then

$$\ln x^r = r \ln x + C$$

for some constant  $C$ . At  $x = 1$ , both sides are zero, thus  $C = 0$ . □



# Basic Property: $\ln x^r = r \ln x$

## Lemma

$$\ln x^r = r \ln x \quad (r \text{ rational}).$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln x^r = \frac{1}{x^r} \frac{d}{dx} x^r = \frac{1}{x^r} r x^{r-1} = r \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (r \ln x) = r \frac{1}{x}.$$

- Then

$$\ln x^r = r \ln x + C$$

for some constant  $C$ . At  $x = 1$ , both sides are zero, thus  $C = 0$ . □



# Basic Property: $\ln x^r = r \ln x$

## Lemma

$$\ln x^r = r \ln x \quad (r \text{ rational}).$$

## Proof.

- Left side:

$$\frac{d}{dx} \ln x^r = \frac{1}{x^r} \frac{d}{dx} x^r = \frac{1}{x^r} r x^{r-1} = r \frac{1}{x}.$$

- Right side:

$$\frac{d}{dx} (r \ln x) = r \frac{1}{x}.$$


- Then

$$\ln x^r = r \ln x + C$$

for some constant  $C$ . At  $x = 1$ , both sides are zero, thus  $C = 0$ . □



$$\text{Range} = (-\infty, \infty)$$



### Theorem

The log function  $\ln x$  has range  $(-\infty, \infty)$  and

$$\lim_{x \rightarrow 0^+} \ln x = -\infty, \quad \lim_{x \rightarrow \infty} \ln x = \infty.$$

### Proof.

- Let  $M > 0$  arbitrary in  $\mathbb{R}$ . Since  $\ln 2 > 0$ ,  $\exists n \in \mathbb{N}$  s.t.


$$n \ln 2 > M, \quad -n \ln 2 < -M.$$

- Since  $n \ln 2 = \ln(2^n)$  and  $-n \ln 2 = \ln(2^{-n})$ ,

$$\ln(2^n) > M, \quad \ln(2^{-n}) < -M. \quad \square$$



$$\text{Range} = (-\infty, \infty)$$



### Theorem

The log function  $\ln x$  has range  $(-\infty, \infty)$  and

$$\lim_{x \rightarrow 0^+} \ln x = -\infty, \quad \lim_{x \rightarrow \infty} \ln x = \infty.$$

### Proof.

- Let  $M > 0$  arbitrary in  $\mathbb{R}$ . Since  $\ln 2 > 0$ ,  $\exists n \in \mathbb{N}$  s.t.


$$n \ln 2 > M, \quad -n \ln 2 < -M.$$

- Since  $n \ln 2 = \ln(2^n)$  and  $-n \ln 2 = \ln(2^{-n})$ ,

$$\ln(2^n) > M, \quad \ln(2^{-n}) < -M. \quad \square$$



$$\text{Range} = (-\infty, \infty)$$



### Theorem

The log function  $\ln x$  has range  $(-\infty, \infty)$  and

$$\lim_{x \rightarrow 0^+} \ln x = -\infty, \quad \lim_{x \rightarrow \infty} \ln x = \infty.$$

### Proof.

- Let  $M > 0$  arbitrary in  $\mathbb{R}$ . Since  $\ln 2 > 0$ ,  $\exists n \in \mathbb{N}$  s.t.


$$n \ln 2 > M, \quad -n \ln 2 < -M.$$

- Since  $n \ln 2 = \ln(2^n)$  and  $-n \ln 2 = \ln(2^{-n})$ ,

$$\ln(2^n) > M, \quad \ln(2^{-n}) < -M. \quad \square$$





Limit:  $\lim_{x \rightarrow \infty} \frac{\ln x}{x^r}$ 

## Theorem

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x^r} = 0 \quad \text{for any } r > 0.$$

$\ln x$  grows slower than any positive power as  $x \rightarrow \infty$ .

## Proof.

- Choose a rational number  $p$  s.t.  $1 - r < p < 1$ . For  $x > 1$ ,

$$\ln x = \int_1^x \frac{1}{t} dt < \int_1^x \frac{1}{t^p} dt = \frac{1}{1-p} t^{1-p} \Big|_1^x = \frac{1}{1-p} (x^{1-p} - 1)$$


- Then

$$0 < \frac{\ln x}{x^r} < \frac{1}{1-p} \frac{x^{1-p} - 1}{x^r} = \frac{1}{1-p} (x^{1-p-r} - x^{-r})$$

Use the pinching theorem to take the limit as  $x \rightarrow \infty$ . □



Limit:  $\lim_{x \rightarrow \infty} \frac{\ln x}{x^r}$



### Theorem

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x^r} = 0 \quad \text{for any } r > 0.$$

$\ln x$  grows slower than any positive power as  $x \rightarrow \infty$ .

### Proof.

- Choose a rational number  $p$  s.t.  $1 - r < p < 1$ . For  $x > 1$ ,

$$\ln x = \int_1^x \frac{1}{t} dt < \int_1^x \frac{1}{t^p} dt = \frac{1}{1-p} t^{1-p} \Big|_1^x = \frac{1}{1-p} (x^{1-p} - 1).$$


- Then

$$0 < \frac{\ln x}{x^r} < \frac{1}{1-p} \frac{x^{1-p} - 1}{x^r} = \frac{1}{1-p} (x^{1-p-r} - x^{-r})$$

Use the pinching theorem to take the limit as  $x \rightarrow \infty$ . □



Limit:  $\lim_{x \rightarrow \infty} \frac{\ln x}{x^r}$



## Theorem

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x^r} = 0 \quad \text{for any } r > 0.$$

$\ln x$  grows slower than any positive power as  $x \rightarrow \infty$ .

## Proof.

- Choose a rational number  $p$  s.t.  $1 - r < p < 1$ . For  $x > 1$ ,


$$\ln x = \int_1^x \frac{1}{t} dt < \int_1^x \frac{1}{t^p} dt = \frac{1}{1-p} t^{1-p} \Big|_1^x = \frac{1}{1-p} (x^{1-p} - 1).$$

- Then

$$0 < \frac{\ln x}{x^r} < \frac{1}{1-p} \frac{x^{1-p} - 1}{x^r} = \frac{1}{1-p} (x^{1-p-r} - x^{-r})$$

Use the pinching theorem to take the limit as  $x \rightarrow \infty$ . □



Limit:  $\lim_{x \rightarrow 0^+} x^r \ln x$ 

## Corollary


$$\lim_{x \rightarrow 0^+} x^r \ln x = 0 \quad \text{for any } r > 0.$$

## Proof.

Let  $y = x^{-1}$ . Then

$$\lim_{x \rightarrow 0^+} x^r \ln x = \lim_{y \rightarrow \infty} y^{-r} \ln y^{-1} = - \lim_{y \rightarrow \infty} \frac{\ln y}{y^r} = 0. \quad \square$$



Limit:  $\lim_{x \rightarrow 0^+} x^r \ln x$ 

## Corollary

$$\lim_{x \rightarrow 0^+} x^r \ln x = 0 \quad \text{for any } r > 0.$$


## Proof.

Let  $y = x^{-1}$ . Then

$$\lim_{x \rightarrow 0^+} x^r \ln x = \lim_{y \rightarrow \infty} y^{-r} \ln y^{-1} = - \lim_{y \rightarrow \infty} \frac{\ln y}{y^r} = 0. \quad \square$$



# Number e



## Definition

The number  $e$  is defined by

$$\ln e = 1$$

i.e., the unique number at which  
 $\ln x = 1$ .

## Theorem


$$\ln e^r = r \quad \text{for any rational number } r.$$

## Proof.

$$\ln e^r = r \ln e = r$$



# Number e



## Definition

The number  $e$  is defined by

$$\ln e = 1$$

i.e., the unique number at which  
 $\ln x = 1$ .

## Theorem


$$\ln e^r = r \quad \text{for any rational number } r.$$

Proof.

$$\ln e^r = r \ln e = r$$



# Number e



## Definition

The number  $e$  is defined by

$$\ln e = 1$$

i.e., the unique number at which  
 $\ln x = 1$ .

## Theorem

$$\ln e^r = r \quad \text{for any rational number } r.$$

## Proof.

$$\ln e^r = r \ln e = r$$





## Quiz

## Quiz

1.  $\ln 1 = ?$  : (a)  $-1$ , (b)  $0$ , (c)  $1$ .
2.  $\ln e = ?$  : (a)  $0$ , (b)  $1$ , (c)  $e$ .
3.  $\lim_{x \rightarrow 0^+} \ln x = ?$  : (a)  $-\infty$ , (b)  $0$ , (c)  $\infty$ .
4.  $\lim_{x \rightarrow \infty} \ln x = ?$  : (a)  $-\infty$ , (b)  $0$ , (c)  $\infty$ .



# Outline

- Definition and Properties of the Natural Log Function
  - Definition of the Natural Log Function
  - Examples
  - Algebraic Properties of the Natural Log Function
- Range and Limits of the Natural Log Function
  - Range of the Natural Log Function
  - Limits of the Natural Log Function
- Number  $e$

