

Math 3331 Differential Equations

5.1 Definition of the Laplace Transform

Jiwen He

Department of Mathematics, University of Houston

`jiwenhe@math.uh.edu`
`math.uh.edu/~jiwenhe/math3331`




5.1 Definition of the Laplace Transform

- Use of the Laplace Transform
 - DEs with constant coefficients
 - Discontinuous forcings
- Definition of the Laplace Transform
- Examples
 - $\mathcal{L}(1)(s) = 1/s, s > 0$
 - $\mathcal{L}(e^{at})(s) = 1/s - a, s > a$
 - $\mathcal{L}(t)(s) = 1/s^2$
 - $\mathcal{L}(t^n)(s) = n!/s^{n+1}$
 - $\mathcal{L}(\sin at)(s)$ and $\mathcal{L}(\cos at)(s)$
 - $\mathcal{L}(f)(s)$, f being discontinuous
- Piecewise Continuous Functions
- Functions of Exponential Order
- Existence of the Laplace Transform



Use of the Laplace Transform

- Technique for solving linear DEs with constant coefficients
- Useful for discontinuous forcings



Definition of the Laplace Transform

Def.: Given a real or complex function $f(t)$, the Laplace (\mathcal{L}) transform of f is the following function of s :

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt \\ &\equiv \lim_{T \rightarrow \infty} \int_0^T e^{-st} f(t) dt \end{aligned}$$

Notation:

$$F(s) = \mathcal{L}(f)(s) = \mathcal{L}\{f(t)\}(s)$$



Example 1: $\mathcal{L}(1)(s) = 1/s, s > 0$

$$\begin{aligned}\mathcal{L}(1)(s) &= \int_0^{\infty} 1 e^{-st} dt = \lim_{T \rightarrow \infty} \int_0^T e^{-st} dt = \lim_{T \rightarrow \infty} \left[-\frac{1}{s} e^{-st} \Big|_{t=0}^T \right] \\ &= \lim_{T \rightarrow \infty} \left[-\frac{1}{s} e^{-sT} + \frac{1}{s} \right] = \frac{1}{s} \quad \text{for } s > 0\end{aligned}$$



Example 2: $\mathcal{L}(e^{at})(s) = 1/s - a, s > a$

$$\begin{aligned}\mathcal{L}(e^{at})(s) &= \int_0^{\infty} e^{at} e^{-st} dt = \lim_{T \rightarrow \infty} \int_0^T e^{-(s-a)t} dt \\ &= \lim_{T \rightarrow \infty} \left[-\frac{1}{s-a} e^{-(s-a)t} \Big|_{t=0}^T \right] \\ &= \lim_{T \rightarrow \infty} \left[-\frac{1}{s-a} e^{-(s-a)T} + \frac{1}{s-a} \right] = \frac{1}{s-a} \quad \text{for } s > a\end{aligned}$$



Example 3: $\mathcal{L}(t)(s) = 1/s^2$

$$\mathcal{L}(t)(s) = \int_0^{\infty} t e^{-st} dt$$

$$= \lim_{T \rightarrow \infty} \int_0^T t e^{-st} dt$$

$$= \lim_{T \rightarrow \infty} \left(-\frac{t}{s} e^{-st} - \frac{1}{s^2} e^{-st} \right) \Big|_{t=0}^T$$

$$= \lim_{T \rightarrow \infty} \left(-\frac{T}{s} e^{-sT} - \frac{1}{s^2} e^{-sT} + \frac{1}{s^2} \right) = \frac{1}{s^2}$$

Integration by parts

$$\int t e^{-st} dt$$

$$= -\frac{1}{s} \left(t e^{-st} - \int e^{-st} dt \right)$$

$$= -\frac{t}{s} e^{-st} - \frac{1}{s^2} e^{-st}$$



Example 4: $\mathcal{L}(t^n)(s) = n!/s^{n+1}$

$$\mathcal{L}(t^n)(s) = \int_0^{\infty} t^n e^{-st} dt$$

$$= \lim_{T \rightarrow \infty} \int_0^T t^n e^{-st} dt$$

$$= \lim_{T \rightarrow \infty} \left(-\frac{t^n}{s} e^{-st} - \dots - \frac{n!}{s^{n+1}} e^{-st} \right) \Big|_{t=0}^T$$

$$= \lim_{T \rightarrow \infty} \left(-\frac{T^n}{s} e^{-sT} - \dots - \frac{n!}{s^{n+1}} e^{-sT} + \frac{n!}{s^{n+1}} \right) = \dots$$

$$= \frac{n!}{s^{n+1}}$$

Integration by parts

$$\int t^n e^{-st} dt$$

$$= -\frac{1}{s} t^n e^{-st}$$

$$+ \frac{n}{s} \int t^{n-1} e^{-st} dt$$

$$= -\frac{t^n}{s} e^{-st} - \dots - \frac{n!}{s^{n+1}} e^{-st}$$



Example 5: $\mathcal{L}(\sin at)(s)$ and $\mathcal{L}(\cos at)(s)$

$$\mathcal{L}(\sin at)(s) = \int_0^{\infty} \sin at e^{-st} dt = \cdots = \frac{a}{s^2 + a^2}$$

$$\mathcal{L}(\cos at)(s) = \int_0^{\infty} \cos at e^{-st} dt = \cdots = \frac{s}{s^2 + a^2}$$



Example 6: $\mathcal{L}(f)(s)$, f being discontinuous

Compute the Laplace transform of the step function

$$f(t) = \begin{cases} 1, & 0 \leq t < 1 \\ 0, & t \geq 1 \end{cases}$$

$$\begin{aligned} \mathcal{L}(f)(s) &= \int_0^1 1 e^{-st} dt = -\frac{1}{s} e^{-st} \Big|_{t=0}^1 \\ &= -\frac{1}{s} e^{-s} + \frac{1}{s}. \end{aligned}$$



Piecewise Continuous Functions

Def.: $f(t)$ is piecewise continuous if

- in any finite interval $0 < t < T$ there are at most finitely many discontinuities
- at any point of discontinuity t_d the left and right limits f_{\mp} exist:

$$f_{-}(t_d) = \lim_{t \rightarrow t_d^{-}} f(t), \quad f_{+}(t_d) = \lim_{t \rightarrow t_d^{+}} f(t)$$

Ex.: $f(t) = \begin{cases} 0 & \text{if } 0 \leq t < 1 \\ e^{t-1} & \text{if } t \geq 1 \end{cases}$ has

a discontinuity at $t_d = 1$:

$$f_{-}(1) = 0, \quad f_{+}(1) = 1$$



Functions of Exponential Order

Def.: $f(t)$ is of exponential order if there are constants C, a s.t.

$$|f(t)| \leq Ce^{at} \text{ for all } t$$

Meaning: $f(t)$ grows at most exponentially if $t \rightarrow \infty$

Ex.: e^{t^2} is *not* of exponential order

Ex.: $e^{10,000t}$ is of exponential order



Existence of the Laplace Transform

Thm.: If $f(t)$ is piecewise continuous in $0 \leq t < \infty$ and of exponential order, then $\mathcal{L}(f)(s)$ exists for $s > a$.

