

$$f'(x) = r(x)$$

Math 1431

Test 4 - Practice Problems

$$g'(x) = s(x)$$

1. Suppose $f(x)$ is an anti-derivative of $r(x)$, and $g(x)$ is an anti-derivative of $s(x)$. We are given the data in the table about the functions f , g , r and s .

| | | | | |
|--------|---|---|---|---|
| x | 1 | 2 | 3 | 4 |
| $f(x)$ | 3 | 2 | 1 | 4 |
| $r(x)$ | 1 | 4 | 2 | 3 |
| $g(x)$ | 2 | 1 | 4 | 3 |
| $s(x)$ | 4 | 2 | 3 | 1 |

$$f(3) = 1$$

$$g(3) = 4$$

$$f(1) = 3$$

$$g(1) = 2$$

$$a) \int_1^3 (3r(x) - 2s(x)) dx = \int_1^3 (3f'(x) - 2g'(x)) dx$$

$$= (3f(x) - 2g(x)) \Big|_1^3$$
$$= (3 \cdot f(3) - 2 \cdot g(3)) - (3f(1) - 2g(1))$$
$$= (3 - 8) - (9 - 4) = -10.$$

$$g(4) = 3$$

$$g(1) = 2$$

- b) Give the average value of $s(x)$ on the interval $[1, 4]$.

$$\rightarrow = \frac{1}{4-1} \int_1^4 s(x) dx = \frac{1}{3} g(x) \Big|_1^4$$

recall: $g'(x) = s(x)$

$$= \frac{1}{3} (g(4) - g(1)) = \frac{1}{3} (3 - 2)$$

$$= \frac{1}{3}.$$

Average value of $s(x)$ on $[1, 4]$.

2. Suppose $F''(x) = x^2 - \frac{2}{\sqrt{x}} + 1$, $F'(1) = -3$ and $F(1) = 2$. Find $F(x)$.

$$\frac{2}{\sqrt{x}} = 2x^{-\frac{1}{2}}$$

$$F'(x) = \frac{1}{3}x^3 - 2 \cdot 2x^{\frac{1}{2}} + x + C_1 \Rightarrow -3 = \frac{1}{3} - 4 + 1 + C_1$$

$$\Rightarrow C_1 = -\frac{1}{3} \Rightarrow F'(x) = \frac{1}{3}x^3 - 4x^{\frac{1}{2}} + x - \frac{1}{3}$$

$$\Rightarrow F(x) = \frac{1}{12}x^4 - 4 \cdot \frac{2}{3}x^{\frac{3}{2}} + \frac{1}{2}x^2 - \frac{1}{3}x + C_2$$

$$2 = \frac{1}{12} - \frac{8}{3} + \frac{1}{2} - \frac{1}{3} + C_2$$

Find C_2 .

Rewrite $F(x)$.

3. Evaluate: $\frac{d}{dx} \int_{-1}^{2x^3} \sin(t^2) dt$ and $\frac{d}{dx} \int_{1-2x}^{2x^3} \sin(t^2) dt$.

$$\rightarrow = \sin((2x^3)^2) \cdot 6x^2 = 6x^2 \sin(4x^6)$$

$$\rightarrow = \frac{d}{dx} \int_{500}^{500} \sin(t^2) dt + \frac{d}{dx} \int_{500}^{2x^3} \sin(t^2) dt$$

$$= -\frac{d}{dx} \int_{500}^{1-2x} \sin(t^2) dt + 6x^2 \sin(4x^6)$$

$$= -\sin((1-2x)^2) \cdot (-2) + 6x^2 \sin(4x^6) = 2\sin((1-2x)^2) + 6x^2 \sin(4x^6)$$

The value 500 could be ANY number.

4. Find a formula for $f(x)$: $2x^3 - 3x^2 + x - 1 = \int_{-1}^x f(t) dt$

Differentiate.

$$6x^2 - 6x + 1 = f(x)$$

$$\frac{d}{dx} \int_a^{u(x)} f(t) dt = f(u(x)) u'(x)$$

Note:

$$\int_a^b f(t) dt + \int_b^c f(t) dt = \int_a^c f(t) dt$$

$$5. \int_2^7 x\sqrt{x^2+2} dx = \frac{1}{2} \int_2^7 \underbrace{(x^2+2)}_u \underbrace{2x dx}_{du}$$

$$u = x^2 + 2$$

$$du = 2x dx$$

$$x=7 \Rightarrow u=51$$

$$x=2 \Rightarrow u=6$$

$$= \frac{1}{2} \int_6^{51} u^{1/2} du = \frac{1}{2} \cdot \frac{2}{3} u^{3/2} \Big|_6^{51} \\ = \frac{1}{3} (51^{3/2} - 6^{3/2})$$

$$6. \int (3\sec^2(2x) - 2\sqrt{x-1}) dx = \int 3\sec^2(2x) dx - \int 2\sqrt{x-1} dx$$

$$= \frac{3}{2} \tan(2x) - 2 \cdot \frac{2}{3} (x-1)^{3/2} + C$$

$$= \frac{3}{2} \tan(2x) - \frac{4}{3} (x-1)^{3/2} + C$$

you can use u-sub if you aren't comfortable.

7. Give the average value of $f(x) = x^2 - 2x + 4$ on the interval $[-1, 2]$, and verify the conclusion of the mean value theorem for integrals for this function on this interval.

$$\rightarrow = \frac{1}{2-(-1)} \int_{-1}^2 (x^2 - 2x + 4) dx = \frac{1}{3} \left(\frac{1}{3} x^3 - x^2 + 4x \right) \Big|_{-1}^2$$

$$= \frac{1}{3} \left[\left(\frac{8}{3} - 4 + 8 \right) - \left(-\frac{1}{3} - 1 - 4 \right) \right]$$

$$= \frac{1}{3} [3 + 9] = 4 \leftarrow \text{Average value of } f(x) \text{ on } [-1, 2].$$

Now, find a value c so that

$-1 < c < 2$ and

$$f(c) = 4$$

$$c^2 - 2c + 4 = 4$$


$$\Rightarrow c^2 - 2c = 0$$

$$c(c-2) = 0$$

$$c=0 \text{ or } c=2$$

Note: $-1 < 0 < 2$.

8. The graph of $f(x)$ is shown below. The area of region A is $\frac{7}{3}$, the area of region B is $\frac{34}{3}$, and the area of region C is $\frac{7}{3}$.




- a. Give the area of the region bounded between the graph of $f(x)$ and the x -axis on the interval $[-2, 4]$.

$$\begin{aligned} & \text{Area}(A) + \text{Area}(B) + \text{Area}(C) \\ &= \frac{7}{3} + \frac{34}{3} + \frac{7}{3} = \frac{48}{3} \end{aligned}$$

b. $\int_{-1}^4 f(x) dx = -\text{Area}(B) + \text{Area}(C)$

$$= -\frac{34}{3} + \frac{7}{3} = -9.$$

9. Find the **area** bounded between the graphs of $f(x) = x^3 - x^2$ and $g(x) = 2x$ on the interval $[0,2]$. The graph is shown below.



$$\begin{aligned}x^3 - x^2 &= 2x \\x^3 - x^2 - 2x &= 0 \\x(x^2 - x - 2) &= 0 \\x(x-2)(x+1) &= 0\end{aligned}$$

$$\text{Area} = \int_0^2 (\text{Top} - \text{Bottom}) dx$$

$$= \int_0^2 (2x - (x^3 - x^2)) dx$$

$$= \int_0^2 (2x - x^3 + x^2) dx = \left(x^2 - \frac{1}{4}x^4 + \frac{1}{3}x^3 \right) \Big|_0^2$$


$$= 4 - 4 + \frac{8}{3} - 0$$

$$= \frac{8}{3}$$

$$y = x^3 - x^2 \quad y = 2x$$

10. The region bounded between the graphs of $f(x) = x^3 - x^2$ and $g(x) = 2x$ on the interval $[0, 2]$

(see the previous problem) is rotated around the y-axis to generate a solid. Find the volume.



Shell
 Thickness = dx
 Surface Area = $2\pi r h$
 $= 2\pi x (2x - (x^3 - x^2))$

Inf. volume = $2\pi x (2x - x^3 + x^2) dx$
 (shell)

Full volume = $\int_0^2 2\pi x (2x - x^3 + x^2) dx$

$= 2\pi \int_0^2 (2x^2 - x^4 + x^3) dx$

$= 2\pi \left(\frac{2}{3}x^3 - \frac{1}{5}x^5 + \frac{1}{4}x^4 \right) \Big|_0^2$


$= 2\pi \left[\left(\frac{16}{3} - \frac{32}{5} + 4 \right) - 0 \right] = \underline{\underline{\text{you do it.}}}$

11. Sketch the region in the first quadrant bounded between the graphs of $f(x) = x + 3$ and $g(x) = (x + 1)^2$. Then rotate this region around the y -axis to generate a solid.

$x = y - 3$

a. Give a formula involving integral(s) in y for the volume generated. Do not compute the integral(s)!

$\rightarrow dy$ integration (Need horiz. line segs)



$1 \leq y \leq 3$

Disk, Thickness = dy



Area = $\pi r^2 = \pi (-1 + \sqrt{y})^2$

Inf. volume = $\int \pi (-1 + \sqrt{y})^2 dy$

$3 \leq y \leq 4$ washer

See next page.


b. Give a formula involving integral(s) in x for the volume generated. Do not compute the integral(s)!

$\rightarrow dx$ integration use vertical line segs.

$x + 3 = (x + 1)^2$
 $x + 3 = x^2 + 2x + 1$
 $x^2 + x - 2 = 0$
 $(x + 2)(x - 1) = 0$
 $x = -2, x = 1$

Shells: Thickness = dx


Surface Area = $2\pi r h = 2\pi x (x + 3 - (x + 1)^2)$
 $= 2\pi x (x + 3 - (x + 1)^2)$



\Rightarrow Inf. volume = $\int 2\pi x (x + 3 - (x + 1)^2) dx$

Full volume = $\int_0^1 2\pi x (x + 3 - (x + 1)^2) dx$

Washer ($3 \leq y \leq 4$)



$$r = y - 3$$

$$R = -1 + \sqrt{y}$$


Thickness = dy

$$\begin{aligned} \text{Area} &= \pi (R)^2 - \pi (r)^2 \\ &= \pi (-1 + \sqrt{y})^2 - \pi (y - 3)^2 \end{aligned}$$

$$\text{Inf. Volume} = \int (\pi (-1 + \sqrt{y})^2 - \pi (y - 3)^2) dy$$

$$\text{Full Volume} = \underbrace{\int_1^3 \pi (-1 + \sqrt{y})^2 dy}_{1 \leq y \leq 3} + \underbrace{\int_3^4 (\pi (-1 + \sqrt{y})^2 - \pi (y - 3)^2) dy}_{3 \leq y \leq 4}$$

12. Repeat the previous problem, assuming that the region is rotated around the x-axis to generate a volume.



a) dy integral(s)
 \Rightarrow horiz line segs.

r $1 \leq y \leq 3$
 shell.

Thickness = dy

$$\text{Surface area} = 2\pi r h$$

$$= 2\pi y (-1 + \sqrt{y})$$

$$\text{Inf. Volume} = \int 2\pi y (-1 + \sqrt{y}) dy$$

$3 \leq y \leq 4$ shell

Thickness = dy


$$\text{Surface Area} = 2\pi r h$$

$$= 2\pi y ((-1 + \sqrt{y}) - (y - 3))$$

$$\text{Inf. Volume} = \int 2\pi y (2 + \sqrt{y} - y) dy$$


$$\text{Full Volume} = \underbrace{\int_1^3 2\pi y (-1 + \sqrt{y}) dy}_{1 \leq y \leq 3} + \underbrace{\int_3^4 2\pi y (2 + \sqrt{y} - y) dy}_{3 \leq y \leq 4}$$

b) dx integration. \Rightarrow vertical line segments



washers:

Thickness = dx



$$\text{Area} = \pi R^2 - \pi r^2$$


$$= \pi (x+3)^2 - \pi ((x+1)^2)^2$$

$$\text{Int. Volume (washer)} = \left(\pi (x+3)^2 - \pi (x+1)^4 \right) dx$$

$$\text{Full Volume} = \int_0^1 \left(\pi (x+3)^2 - \pi (x+1)^4 \right) dx .$$

13. a. Give the Upper Riemann sum for the function $f(x) = \begin{cases} 1-x, & -1 \leq x < 0 \\ x+1, & 0 \leq x < 1 \\ 3-x, & 1 \leq x \leq 2 \end{cases}$ on the interval

$[-1, 2]$ with respect to the partition $P = \{-1, -1/4, 1/4, 1, 3/2, 2\}$.




$$f(-1) \cdot \frac{3}{4} + f(-\frac{1}{4}) \cdot \frac{1}{2} + f(\frac{1}{4}) \cdot \frac{3}{4} + f(1) \cdot \frac{1}{2} + f(\frac{3}{2}) \cdot \frac{1}{2}$$

put the values in.
...

- b. Give the Lower Riemann sum for the function $f(x) = \begin{cases} 1-x, & -1 \leq x < 0 \\ x+1, & 0 \leq x < 1 \\ 3-x, & 1 \leq x \leq 2 \end{cases}$ on the interval


$[-1, 2]$ with respect to the partition $P = \{-1, -1/4, 1/4, 1, 3/2, 2\}$.



$$f(-\frac{1}{4}) \cdot \frac{3}{4} + f(0) \cdot \frac{1}{2} + f(\frac{1}{4}) \cdot \frac{3}{4} + f(\frac{3}{2}) \cdot \frac{1}{2} + f(2) \cdot \frac{1}{2}$$

Get the values
...

14. Sketch the region bounded between the graphs of $x + y = 2$ and $x = y^2$. Then give a formula for the area of the region involving integral(s) in x . Repeat the process with integral(s) in y .



$$2 - y = y^2$$

$$0 = y^2 + y - 2$$

$$0 = (y + 2)(y - 1)$$

$$y = -2, y = 1$$

$$y = 1 \Rightarrow x = 1$$

$$y = -2 \Rightarrow x = 4$$

$$\text{Area} = \int_0^1 (\overset{\text{Top}}{\sqrt{x}} - \overset{\text{bottom}}{-\sqrt{x}}) dx + \int_1^4 (\overset{\text{Top}}{2-x} - \overset{\text{bottom}}{-\sqrt{x}}) dx$$

$$= \int_0^1 2\sqrt{x} dx + \int_1^4 (2-x+\sqrt{x}) dx.$$

$$\text{Area} = \int_{-2}^1 ((2-y) - y^2) dy$$

\uparrow right \uparrow left