Today: *More* Section 6.2/6.3

- **Test 4 , Dec. 6 - 8**
- **Final Exam, Dec. 17 - 19**
- Dates are subject to change...
- **Practice Test 4** (counts as a quiz) and **Practice Final Exam** (counts as 2 quizzes) will be posted soon.
- **EMCFs** are posted for the rest of the week, and **Homework** is posted for Monday.
- **Practice Problems** will be posted soon for Test 4.
- **Friday's Quiz** will cover volumes of revolution.

Example: Create a right circular cone with height h and base radius r as a volume of revolution. Then use your creation to derive the formula for the volume of a cone.



Thickness $= dx$

Surface Area $= 2\pi x\left(-\frac{h}{r}x + h\right)$

Inf. Volume (shell) $= 2\pi x\left(-\frac{h}{r}x + h\right)dx$

Full Volume $= \int_0^r 2\pi x\left(-\frac{h}{r}x + h\right)dx$

$$= \int_0^r \left(-2\pi\frac{h}{r}x^2 + 2\pi h x\right)dx$$

$$= \left(-2\pi\frac{h}{r}\cdot\frac{x^3}{3} + \pi h x^2\right)\Big|_0^r$$

$$= -2\pi\frac{h}{r}\cdot\frac{r^3}{3} + \pi h r^2$$

$$= -\frac{2}{3}\pi h r^2 + \pi h r^2$$

$$= \frac{1}{3}\pi r^2 h$$

Volumes by Slicing: Main Setting

A solid S intersects the xy-plane in the region R shown below. Every cross section of the solid S taken perpendicular to a given line can be described. **Find the volume of S.**



Typically, the solid object is aligned so that the cross sections perpendicular to one of the axes is known.



Example: A solid object is aligned so that its base is the region in the xy-plane bounded between the x-axis and the curve $y = 1 - x^2$. Cross sections taken perpendicular to the x-axis are squares. Find the volume of the region.





Thickness = dx

Area = $(1-x^2)^2$

Inf. Volume = $(1-x^2)^2\,dx$

Full Volume = $\int_{-1}^{1}(1-x^2)^2\,dx = \cdots$

Example: A solid object is aligned so that its base is the region in the *xy*-plane bounded between the *x*-axis and the curve $y = 1 - x^2$. Cross sections taken perpendicular to the *y*-axis are semi-circles with their diagonals lying in the *xy*-plane. Find the volume of the region.



$y = 1 - x^2$

$x^2 = 1 - y$

$x = \pm\sqrt{1-y}$

Thickness = dy

$\text{Area} = \frac{1}{2}\pi r^2 = \frac{1}{2}\pi\left(\sqrt{1-y}\right)^2$

$\text{Inf Volume} = \frac{1}{2}\pi(1-y)\,dy$

$$\text{Full Volume} = \int_0^1 \frac{1}{2}\pi(1-y)\,dy$$