

MATH 4331
Introduction to Real Analysis
Fall 2015

First name: _____ Last name: _____

Points:

Assignment 6, due Thursday, October 22, 2:30pm

Please staple this cover page to your homework. When asked to prove something, make a careful step-by-step argument. You can quote anything we covered in class in support of your reasoning.

Problem 1

If a function $f : [a, b] \rightarrow \mathbb{R}$ is Lipschitz-continuous with Lipschitz constant C , then prove that for any partition P of $[a, b]$, we have $U(f, P) - L(f, P) \leq C(b - a)\text{mesh}(P)$.

Problem 2

Show that if a real-valued function f is bounded and Riemann integrable on $[a, b]$, so is $|f|$, $|f|(x) = |f(x)|$.

Problem 3

Show that if f and g are real-valued, bounded and Riemann integrable on $[a, b]$ and $f(x) \leq g(x)$ for each $x \in [a, b]$, then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

Problem 4

Prove the mean value theorem for integrals: If a real-valued function f is continuous on $[a, b]$, then there is $c \in (a, b)$ such that

$$f(c) = \frac{1}{b - a} \int_a^b f(x) dx.$$

Problem 5

Show that the function defined by $f(x) = \sin(1/x)$ for $x > 0$ and $f(0) = 0$ is Riemann integrable on $[0, 1]$.