

## 2017 Houston Summer School on Dynamical Systems

### *Problem set: statistical properties*

1. Let  $\mu$  be a probability measure on a measurable space  $X$ , and let  $f: X \rightarrow X$  be a measurable map. Prove that the following definitions of “ $f$ -invariant” are equivalent.
  - (a) For every measurable  $E \subset X$  we have  $\mu(f^{-1}E) = \mu(E)$ .
  - (b) For every measurable  $\varphi: X \rightarrow \mathbb{R}$  we have  $\int \varphi d\mu = \int \varphi \circ f d\mu$ .
2. Prove that Lebesgue measure is invariant for each of the following.
  - (a) the doubling map  $E_2: S^1 \rightarrow S^1$ ;
  - (b) a circle rotation  $R_\alpha: S^1 \rightarrow S^1$ ;
  - (c) the toral automorphism given by  $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ ;
  - (d) the twist  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  on the torus;
  - (e) the map  $(x, y) \mapsto (x + \alpha, y + x)$  on the torus.
3. Let  $\mu$  be an invariant probability measure for  $f$ , and prove that the following definitions of “ergodic” are equivalent.
  - (a) If  $E$  is a measurable set such that  $f^{-1}(E) = E$ , then  $\mu(E) = 0$  or  $\mu(E) = 1$ .
  - (b) If  $\varphi$  is a measurable function  $\varphi$  such that  $\varphi(x) = \varphi(f(x))$  for  $\mu$ -a.e.  $x$ , then  $\varphi$  is constant  $\mu$ -a.e.
  - (c) If  $\nu_1, \nu_2$  are invariant measures such that  $\mu = a_1\nu_1 + a_2\nu_2$  for some  $a_1, a_2 \geq 0$  with  $a_1 + a_2 = 1$ , then  $\nu_1 = \nu_2 = \mu$ .
4. Recall that given a measure  $\mu$  on the circle  $S^1 = \mathbb{R}/\mathbb{Z}$ , the Fourier transform of  $\mu$  is the function  $\hat{\mu}: \mathbb{Z} \rightarrow \mathbb{C}$  given by  $\hat{\mu}(\kappa) = \int e^{-2\pi i \kappa x} d\mu(x)$ , and that the map  $\mu \mapsto \hat{\mu}$  is 1-1. Use this fact to prove that an irrational rotation is uniquely ergodic.
5. Prove that  $x \in \mathbb{R}/\mathbb{Z}$  is periodic for the doubling map if and only if it is rational.
6. Recall that the Fourier transform of  $\varphi \in L^2(S^1)$  is  $\hat{\varphi}(\kappa) = \int e^{-2\pi i \kappa x} \varphi(x) dx$ , where the integral is taken with respect to Lebesgue measure, and that the map  $\varphi \mapsto \hat{\varphi}$  is a bijection (in fact a unitary isomorphism) between  $L^2(S^1)$  and  $\ell^2(\mathbb{Z})$ . Use this to prove that Lebesgue measure is ergodic for the doubling map.
7. Let  $R_\theta(x) = x + \theta \pmod{1}$  where  $\theta$  is irrational. Let  $dx$  be Lebesgue measure on  $[0, 1]$  and suppose that  $\phi$  is a continuous function on the circle with  $\int \phi dx = 0$ . Show that  $\frac{1}{n} S_n \phi \rightarrow 0$  uniformly; that is, for every  $\epsilon > 0$  there is  $N \in \mathbb{N}$  such that for all  $n \geq N$ , we have  $|\frac{1}{n} S_n \phi(x)| < \epsilon$  for every  $x$ .
8. Use Fourier analysis to prove that Lebesgue measure is ergodic for the toral automorphism  $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ .

9. Consider the map  $f: \mathbb{T}^2 \rightarrow \mathbb{T}^2$  given by  $f(x, y) = (x + \alpha, y + x)$ , where  $\alpha$  is irrational. Prove that  $f$  is uniquely ergodic by following the steps below.
- (a) Prove that Lebesgue measure  $m$  is ergodic using Fourier analysis: use the fact that the characteristic function of any subset of  $\mathbb{T}^2$  is in  $L^2$ , and hence has a Fourier transform in  $\ell^2$ , so that in particular the Fourier coefficients decay at infinity.
  - (b) Let  $\nu$  be any invariant measure for  $f$  and let  $\nu_t$  be the image of  $\nu$  under a rotation by  $t$  in the second coordinate. Show that each  $\nu_t$  is also  $f$ -invariant, and that the average over all values of  $t$  is Lebesgue measure  $m$  on  $\mathbb{T}^2$ . Use ergodicity to conclude that  $\nu_t = m$  for a.e.  $t$ , and hence  $\nu = m$ .
10. Let  $E_2$  be the doubling map. Prove that there are infinitely many ergodic measures  $\mu$  such that (1)  $\mu$  is not periodic, and (2) the support of  $\mu$  is not the whole circle.
- ◊ Recall that the support of  $\mu$  is  $\text{supp}(\mu) = \{x \mid \mu(B(x, \epsilon)) > 0 \text{ for every } \epsilon > 0\}$ .
  - ◊ Hint: code the system by the full 2-shift and consider Markov measures.
- Prove that in fact, there are infinitely many closed subsets  $A \subset S^1$  such that (1)  $A$  is infinite, and (2) there is an ergodic measure  $\mu$  such that  $\text{supp}(\mu) = A$ .
11. Prove that every orbit for  $R_\theta$  is dense if  $\theta$  is irrational. Prove that not every orbit for  $E_2$  is dense. Construct a dense orbit for  $E_2$  using the symbolic representation on the full 2-shift.
12. Let  $X$  be a separable metric space and  $T: X \rightarrow X$  be continuous. Given  $x \in X$  and  $n \in \mathbb{N}$ , let  $\mathcal{E}_{x,n} = \frac{1}{n} \sum_{k=0}^{n-1} \delta_{T^k x}$  be the  $n$ th *empirical measure* for  $x$ . That is, the measure  $\mathcal{E}_{x,n}$  is defined by

$$\int \phi(y) d\mathcal{E}_{x,n}(y) = \frac{1}{n} \sum_{k=0}^{n-1} \phi(T^k x)$$

for every continuous  $\phi: X \rightarrow \mathbb{R}$ . Suppose  $n_j \rightarrow \infty$  and  $\mu \in \mathcal{M}(X)$  are such that  $\mathcal{E}_{x,n_j} \rightarrow \mu$  in the weak\* topology. (Note that if  $X$  is compact then existence of a convergent subsequence follows from compactness of  $\mathcal{M}(X)$ .)

- (a) Show that  $\mu$  is  $T$ -invariant.
- (b) Give an example to show that this may fail if  $T$  is not continuous.
- (c) Say that  $x$  is *generic* for  $\mu$  if  $\mathcal{E}_{x,n} \rightarrow \mu$  (without passing to a subsequence). Birkhoff's ergodic theorem says that if  $\mu$  is ergodic and  $G_\mu$  is the set of generic points for  $\mu$ , then  $\mu(G_\mu) = 1$ . Give an example showing that  $G_\mu$  may be empty if  $\mu$  is invariant but not ergodic.
- (d) Let  $\Sigma$  be the full shift on two symbols and let  $\mu$  be any  $\sigma$ -invariant measure (not necessarily ergodic). Show that  $\mu$  has a generic point.

13. Let  $A$  be a finite alphabet and  $\Sigma \subset A^{\mathbb{Z}}$  (or  $A^{\mathbb{N}}$ ) be a closed  $\sigma$ -invariant subset. Given  $n \in \mathbb{N}$ , let

$$L_n = \{w = w_1 \cdots w_n \mid w_i \in A \text{ for each } 1 \leq i \leq n, \text{ and } w \text{ appears as a subword of some } x \in \Sigma\}.$$

- (a) Let  $a_n = \log \#L_n$  and show that

$$a_{n+m} \leq a_n + a_m \text{ for every } n, m \in \mathbb{N}. \quad (\star)$$

- (b) A sequence satisfying  $(\star)$  is called *subadditive*. Prove *Fekete's lemma*: if  $a_n$  is a subadditive sequence, then  $\lim_{n \rightarrow \infty} \frac{1}{n} a_n$  exists and is equal to  $\inf_{n \in \mathbb{N}} \frac{1}{n} a_n$  (which may be  $-\infty$ ).
- (c) Deduce that  $h(\Sigma) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \#L_n(\Sigma)$  exists for every shift  $\Sigma$ . This limit is called the *topological entropy* of the shift  $\Sigma$ .
- (d) Let  $\Sigma$  be the SFT on two symbols  $\{0, 1\}$  where the only forbidden word is 11. Show that  $\#L_n(\Sigma)$  is the Fibonacci sequence.
- (e) Let  $\Sigma$  be a topological Markov chain with transition matrix  $M$  – that is, a sequence  $x$  is in  $\Sigma$  if and only if  $M_{x_n, x_{n+1}} = 1$  for every  $n$ . Show that  $\#L_n$  is the sum of all the entries of  $M^{n-1}$ .
- (f) Let  $\lambda$  be a positive real eigenvalue of  $M$  with the property that  $|\chi| \leq \lambda$  for all eigenvalues  $\chi$ . (Existence of such an eigenvalue is part of the Perron–Frobenius theorem.) Prove that  $h(\Sigma) = \log \lambda$ .

14. Let  $(X, \mathcal{B}, \mu)$  be a probability space and  $T: X \rightarrow X$  a measure-preserving map. Given measurable sets  $A, B \subset X$ , let

$$C_n(A, B) := |\mu(A \cap T^{-n}B) - \mu(A)\mu(B)|$$

be the  $n$ th correlation function of  $A, B$ . Similarly, given  $L^2$  test functions  $\phi, \psi$ , let

$$C_n(\phi, \psi) := \left| \int \phi \cdot (\psi \circ T^n) d\mu - \int \phi d\mu \int \psi d\mu \right|.$$

- (a) Show that  $C_n(A, B) \rightarrow 0$  for every  $A, B$  if and only if  $C_n(\phi, \psi) \rightarrow 0$  for every  $\phi, \psi \in L^2$ . In this case the measure is called *mixing*.
- (b) Our results on decay of correlations all involve  $C_n(\phi, \psi)$  for sufficiently regular test functions, instead of  $C_n(A, B)$ , or  $C_n(\phi, \psi)$  for arbitrary  $L^2$  functions. This is because even when  $C_n(\phi, \psi)$  decays exponentially for Hölder continuous functions (or some other nice class), we may have very slow decay for  $C_n(A, B)$ , or for arbitrary measurable functions.

Demonstrate this phenomenon as follows: let  $T: [0, 1] \rightarrow [0, 1]$  be the doubling map  $T(x) = 2x \pmod{1}$ , and let  $\mu$  be Lebesgue measure on  $[0, 1]$ . Find measurable sets  $A, B \subset [0, 1]$  such that  $C_n(A, B)$  only decays polynomially – that is, there are  $\gamma, c > 0$  such that  $C_n(A, B) \geq cn^\gamma$  for all  $n$ .

Note that this is equivalent to answering the same question where  $X$  is the full shift on two symbols and  $\mu$  is  $(\frac{1}{2}, \frac{1}{2})$ -Bernoulli.

15. The *Dyck shift*  $\Sigma$  is the two-sided shift on the four-symbol alphabet  $A$  whose letters are the brackets  $(, [, ],$  and  $)$ , and whose allowable sequences are precisely those in which the brackets are opened and closed in the right order. That is, the word  $( [ ] )$  is allowable, but the word  $( [ ) ]$  is not. Similarly,  $(( ([$  is allowable, but  $( ) ([$  is not.

Let  $B \subset A^* := \bigcup_n A^n$  be the set of *balanced* words; that is, the set defined by the following recursive procedure:

- ◇ the empty word is in  $B$ ;
- ◇ if  $w \in B$  then  $(w) \in B$ ;
- ◇ if  $w \in B$  then  $[w] \in B$ .

Consider the subsets

$$\Sigma^R = \{x \in \Sigma \mid \text{for every } n \in \mathbb{N} \text{ there is } m < n \text{ with } x_m x_{m+1} \cdots x_n \in B\},$$

$$\Sigma^L = \{x \in \Sigma \mid \text{for every } m \in \mathbb{N} \text{ there is } n > m \text{ with } x_m x_{m+1} \cdots x_n \in B\}.$$

That is,  $\Sigma^R$  is the set of sequences where every right bracket has a matching left bracket, and  $\Sigma^L$  is the set of sequences where every left bracket has a matching right bracket.

- (a) Let  $X = \{0, 1, 2\}^{\mathbb{Z}}$  be the full shift on three symbols and define  $h: \Sigma \rightarrow X$  by  $h(x)_n = H(x_n)$ , where  $H: A \rightarrow \{0, 1, 2\}$  maps the symbol  $($  to 1, the symbol  $[$  to 2, and both symbols  $)$ ,  $]$  to 0. Show that  $h$  is 1-1 on  $\Sigma^R$ .
- (b) Let  $\mu$  be an  $\sigma$ -invariant probability measure on  $\Sigma$  and show that  $\mu(\Sigma^R \cup \Sigma^L) = 1$ .
- (c) Consider the directed graph  $G$  whose vertices are non-negative integers and which has the following edges:
  - ◇ 2 edges from 0 to 0;
  - ◇ 2 edges from  $n$  to  $n + 1$  for every  $n \geq 0$ ;
  - ◇ 1 edge from  $n$  to  $n - 1$  for every  $n \geq 1$ .
 Let  $a_n$  be the number of paths of length  $n$  on this graph that start at 0. Show that  $\#L_n(\Sigma) = a_n$ .  
*Hint:* it may help to label the two edges from 0 to 0 with the right brackets  $)$  and  $]$ , the two edges from  $n$  to  $n + 1$  with  $($  and  $[$ , and the edge from  $n$  to  $n - 1$  with “ $)$  or  $]$ ”.

- (d) Show that  $h(\Sigma) = \log 3$ .
- (e) If you know about measure-theoretic entropy and the variational principle, show that  $\Sigma$  has two ergodic measures of maximal entropy, and that both are fully supported on  $\Sigma$ .