#### Math 2413 - Review for Test 2

Schedule your exam; students are responsible for scheduling their tests BEFORE the testing window begins. Do not be late for your test. If you're more than 10 minutes late, CASA will not let you in. Your instructor does not control the scheduler; don't miss your scheduled time. If you miss it, try rescheduling online; you can do this only if there are available seats.

Material: Chapter 3

When: By reservation; check your confirmation email.

Where: CASA Testing center (check your confirmation email for exact location)

What to bring: Picture ID, writing utensils.

Time: 75 minutes

Number of questions: Approximately 18.

MC: TBA

FR: 3-4 questions

The grade you'll see right away will be for MC part only (out of ?? points).

Take practice test 2! 5% of your best score will be added to your test grade.

## How to study?

Go over class notes, rework past HW assignments and labquizzes. Work on the main review sheet posted on CASA and take the practice test. Solve all problems on this review sheet. Check the keys for WHW (posted on CASA) to see how to show work properly.

#### What to know:

- Related Rates
- How to determine the intervals of increase/decrease, concave up/down.
- Critical points, local min/max, absolute min/max, points of inflection.
- First derivative test, Second derivative test.
- Given the graph of f', what can you conclude about f?
- Given the graph of f', what can you conclude about f?
- Finding vertical tangents, cusps.
- Graphing a function (with all details).

#### **Related Rates**

- Work all the problems in your homework and quizzes (solve: shadow length, ladder, area of a rectangle, area of a triangle, area of a circle, volume of a sphere, volume of a cone, etc....)
- Must know basic area and volume formulas from geometry.

Area: Circle, Square, Triangle, Rectangle

Surface area: Cube, Rectangular Prism

Volume: Cube, Cone, Cylinder, Sphere, Rectangular Prism

1. Given  $K = 2hx^3$ . Find the derivative of K with respect to time t; assume that h stays constant with time, and x varies with time.

2. The radius of a circle is increasing at a rate of 2inches/sec. How fast is the perimeter changing when the radius is 10 inches?

# Solve all related rates questions on the main review sheet.

**3.** Water is poured into a conical tank at the rate of 64 cubic feet per minute. If the tank is 10 feet tall and the top of the tank has a radius of 6 feet, how fast does the radius of the water surface change when the water is 2 feet deep?

4. Sand is falling onto a conical pile at a rate of 40 cubic feet per minute. If the diameter of the base of the cone is 4 times its altitude. At what rate is the height of the pile changing when it is 20 feet high?

5. Homer is 4 ft tall and he is walking towards a lamp post that is 20 ft high; if he is approaching the lamp at a rate of ½ feet/sec, how fast is his shadow changing when he is 5 feet away from the lamp?

Exercise: A 5 ft ladder is leaning against a wall. If it is pulled away from the wall
from the bottom tip at a rate of 0.5 ft/sec, how fast is the height of the ladder on the
wall changing when the bottom tip is 4 feet away from the wall?

Exercise: The altitude of a triangle is increasing at a rate of 5 in/min while the base is decreasing at a rate of 1 in/min. At what rate is the area of the triangle changing when the altitude is 10 in and the base is 6 in?

6. A particle is moving on the x-axis with position function  $x(t) = -t^3 + t^2 + 1$ .

What is the **velocity** of this particle when t=5?

What is the **speed** of the particle when t=5?

What is the **acceleration** of the particle when t=5?

7. Given:  $f(x) = 2x^3 + 3x^2 - 4$ . Find the intervals over which this function is increasing.

8. Given: f(x) is a polynomial with the **first derivative**:  $f'(x) = -2(x+1)(x-2)(x-3)^2$ . Find the intervals over which this function is decreasing.

9. Given:  $f(x) = 5x^4 + 5x^3 - 4x + 1$ . Find the intervals over which this function is concave up.

10. Given f(x) is a polynomial with **second derivative function**: f''(x) = 5(x+2)(x+1)(x-4). Find the intervals over which this function is concave down.

11. Given:  $f(x) = x^3 + 6x^2 - 9$ .

Find the critical numbers and classify the them.

Find the critical numbers; Classify the critical points.  $f(x) = (x^2 - 6x)^{1/3}$ 12.

$$f(x) = \left(x^2 - 6x\right)^{1/3}$$

13. Given: 
$$f(x) = (x-12)^{2/3}$$
; classify the point (12,  $f(12)$ ).

- a) Local max
- b) Point of inflection
- c) Vertical tangent
- d) Vertical cusp
- e) None of the above.

14. Given 
$$f(x) = \frac{1}{20}x^5 - \frac{1}{4}x^4 + 2$$
, State the x-coordinates of any points of inflection.

15. Find absolute minimum and maximum values (if any) of f on the given interval and state where those values occur.

$$f(x) = 2x^3 - 3x^2 + 2$$
 [-1,2]

16. Given f(x) is a polynomial with:

$$f(2) = 4, f'(2) = 0, f''(2) = -5$$
. Classify the point (2,4).

- a) Intercept
- b) Local min
- c) Local max
- d) Point of inflection
- e) Vertical cusp
- f) None of the above.

17. If the function  $f(x) = Bx^3 + 4x + 1$  has a critical point at x = 2, find the value of B.

18. Verify that Mean Value Theorem applies. Find the c satisfying the Mean Value theorem on this interval.

$$f(x) = x^3 - x + 1$$
 over [1,3].

Exercise: Does MVT apply over the given interval?

$$f(x) = x^3 + x$$
 over [1,3].

$$f(x) = \sqrt{x-2}$$
 over [1,3].


$$f(x) = \frac{x}{x-2}$$
 over [1,3].

$$f(x) = 2|x-1|$$
 over [0,3].

19. The graph of f'(x) is given.

When is f(x) increasing?

When is f(x) decreasing?




Critical numbers of f(x):

Local min/max points for f(x):

20. The graph of f''(x) is given.

When is f(x) concave up?



When is f(x) concave down?

Points of inflection for f(x):

21	.State the domain, any asymptotes, critical numbers, the intervals where the
	function is increasing/decreasing, concave up/down, and any Points of
	inflection.

$$f(x) = \frac{-10x}{\left(x-4\right)^2}$$

Derivatives:

$$f'(x) = \frac{10x + 40}{(x - 4)^3}$$
$$f''(x) = \frac{-20x - 160}{(x - 4)^4}$$

Domain:	
Asymptotes:	
Intercepts:	

Critical Points:

The function is Increasing on:

The function is Decreasing on:

Local extreme points:

The function is Concave up on:				
The function is Concave down on:				
Points of inflection:				
Exercise: Graph the function.				

EXERCISE: State the domain, any asymptotes, critical numbers, the intervals where the function is increasing/decreasing, concave up/down, and GRAPH this function.

$$f(x) = \frac{x^2}{(x+2)^2}$$

$$f'(x) = \frac{4x}{(x+2)^3}$$

$$f''(x) = \frac{-8x+8}{(x+2)^4}$$

### **EXERCISE**:

What can you say about a function with these properties:

- 1. The domain is all real numbers except 3 and -3.
- 2. The function has vertical asymptotes at x = 3 and x = -3
- 3. The function is symmetric about the y-axis

$$4. \lim_{x\to\infty} f(x) = -1$$

$$5. f(0) = 0, f(2) = 0, f(4) = 0$$

6. 
$$f'(x) < 0$$
 for  $0 < x < 1$  and  $x > 3$ 

7. 
$$f'(x) > 0$$
 for  $1 < x < 3$ 

8. 
$$f''(x) \le 0$$
 for  $0 \le x \le 1/2$ 

## **POPPER#**

**Question#** Is there a vertical tangent or a vertical cusp at the critical number for

$$f(\mathbf{x}) = (\mathbf{x} + \mathbf{3})^{\frac{2}{3}}?$$

- a. Vertical tangent
- b. cusp


**Question#** Is there a vertical tangent or a vertical cusp at the critical number for

$$f(x) = (x-5)^{\frac{1}{5}}$$
?

- a. Vertical tangent
- b. cusp

**Question#** Classify x = -1 if the graph is the derivative of f.

- a. relative maximum
- b. relative minimum
- c. POI
- d. who cares?



Question# The graph of **the first derivative of** f(x) is given below (note the derivative is given!). Which point is a local minimum for f(x)?

- a) P
- b) M
- c) N
- d) Q
- e) None of the above

