

**MATH 6361**  
**Applied Analysis**  
**Spring 2019**

First name: \_\_\_\_\_ Last name: \_\_\_\_\_

Points:

## Assignment 1, due Friday, January 25, 10am

Please staple this problem sheet to your homework. When asked to prove something, make a careful step-by-step argument. You can quote anything we covered in class in support of your reasoning.

### Problem 1

Establish the identity

$$\|z - x\|^2 + \|z - y\|^2 = \frac{1}{2}\|x - y\|^2 + 2\|z - \frac{1}{2}(x + y)\|^2$$

for  $x, y, z$  in a real or complex inner product space.

### Problem 2

Let  $P$  be a bounded linear map on a real or complex Hilbert space  $H$ . Show that if  $P^2 = P$  and for each  $x, y \in H$ ,  $\langle Px, y \rangle = \langle x, Py \rangle$ , then the range of  $P$  is closed and  $P$  is the orthogonal projection onto its range. Hint: If  $Px = 0$  then  $x \perp \text{ran}(P)$ .

### Problem 3

Let  $K$  be a closed, convex set in a real Hilbert space  $H$  and  $x_0 \in H \setminus K$ . Recall that there is a unique  $y_0 \in K$  such that  $\|y_0 - x_0\| = \inf_{y \in K} \|y - x_0\|$ . Show that  $z \in K$  satisfies  $z = y_0$  if and only if for each  $y \in K$ ,  $\langle x_0 - z, y \rangle \leq \langle x_0 - z, z \rangle$ . Hint: Consider for  $t \in [0, 1]$  the squared distance between  $x_0$  and  $ty_0 + (1 - t)z$ .

### Problem 4

The complex exponentials  $u_n(x) = \frac{1}{\sqrt{2\pi}}e^{inx}$ , define an orthonormal basis for  $L^2([-\pi, \pi])$ , so for  $f \in L^2([-\pi, \pi])$ ,  $\|f\|_2^2 = \sum_{n=-\infty}^{\infty} |\langle f, u_n \rangle|^2$ . Evaluate both sides of this identity for (a)  $f(x) = 1$ , if  $x \geq 0$  and  $f(x) = 0$ , if  $x < 0$  and (b)  $f(x) = \pi - |x|$ . Conclude a series summation formula for series of real numbers in both cases.