Let 
$$f(x) = x^{2} + 10 \times +24$$
 be a quadratic function. Find:  
 $f(x) = x^{2} + 10 \times +24 = 0$   
 $f(x) = x^{2} + 10 \times +24 = 0$   
 $(x+6)(x+4) = 0 = x = -6$   
 $x = -4$   
A.  $\{-6, -4\}$  B.  $75. \ 26, 4\}$  C. 24 D. none

(2) y-intercept = 
$$f(0) = 0^2 + 10 \cdot 0 + 24 = 24 \Longrightarrow$$
 (0,24)  
A. (0,0) B. (24,0) C. (0,24) D. none  
 $X = -\frac{b}{10} = -\frac{10}{2 \cdot 1} = -5$ 

(3) Vertex coordinates  

$$y = f(-5) = (-5)^{2} + 10(-5) + 24 = -1$$
  
A.  $(5, -1)$  B.  $(-5, 1)$  C.  $(-5, -1)$  D. none

## Math 1330 - Section 2.2 Polynomial Functions

Our objectives in working with polynomial functions will be, first, to gather information about the graph of the function and, second, to use that information to generate a reasonably good graph without plotting a lot of points. In later examples, we'll use information given to us about the graph of a function to write its equation.

 $\int f(x) = \frac{2}{2}x^{3} + 3x^{2} - 5$ 

 $\sqrt{f(x)} = \frac{2}{2} \times \sqrt{4} \sqrt{3} \times \sqrt{4} - \times \sqrt{4} \sqrt{2}$ 

 $f(x) = x^4 + \sqrt{x}$ 

A polynomial function is a function of the form

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_2 x^2 + a_1 x + a_0$$

where  $a_n \neq 0$ ,  $a_0, a_1, ..., a_n$  are real numbers and *n* is a whole number.

The number  $a_n$  is called the **leading coefficient**. The degree of the polynomial function  $\int f(x) = 5 x^{3} - 3 x^{\dagger} + 2 x - 5 = -5$ <u>is n</u>. -> if even, then "even-degree" polynomial  $P(0) = a_0$  and this number is called the **constant coefficient**. >Highest exponent gives the degree of polynomial -> if odd, they **Example:** The graph of  $f(x) = x(x-2)^3(x+1)^2$  is given: "odd-degree" polynomial -> find degree : Need "leading term  $f(x) = x(x-2)^3(x+1)^2$  $= x \left( \begin{array}{c} x^{3} + \dots \end{array} \right) \left( \begin{array}{c} x^{2} + \dots \end{array} \right)$   $\implies \text{leading term: } x \cdot x \cdot x = x \qquad = x^{6}$ 0 -2 Ś 1 Ź It is a 6<sup>th</sup> degree polynomial i.e. <u>even</u> degree polynomial -2-





If the degree of the function is even and  $a_n > 0$ , then the end behavior of the function is **Degree: Even, Coefficient:** +

*n* even and  $a_n > 0$  (even degree and leading coefficient positive)



If the degree of the function is even and  $a_n < 0$ , then the end behavior of the function is **Degree: Even, Coefficient:** -



If the degree of the function is odd and  $a_n > 0$ , then the end behavior of the function is **Degree: Odd, Coefficient:** +





If the degree of the function is odd and  $a_n < 0$ , then the end behavior of the function is **Degree: Odd, Coefficient:** -



3



**Example 1:** Determine the end behavior of the function:

a) 
$$f(x) = x^4 - 5x^2 + 4$$
.  
• degree = 4 even  
positive coefficient  
 $(x^2 \text{ like})$   
b)  $f(x) = -4x - x^2 + 2x^3 - x^5$ .  
• degree = 5 odd  
negative coefficient =)  
 $(-x^2 \text{ like})$   
c)  $f(x) = 2x(x-1)^2(2x-1)^3(4-x)^2$ .  
degree =  $1+2 + 3 + 1 = 7$  odd  
leading coefficient = $2 \cdot 1^2 \cdot 2^3 \cdot (-1)$   
 $= -16$  negative  $(-x^3 \text{ like})$ 

Next, you should be able to find the *x* intercept(s) and the *y* intercept of a polynomial function.

You will need to set the function equal to zero and then use **the Zero Product Property** to find the *x* intercept(s). That means if ab = 0, then either a = 0 or b = 0. To find the *y* intercept of a function, you will find f(0).

To be continued  
on Friday, 22405 (look for graph on next elide)  
Example 2: Find the x and y intercept(s) of 
$$f(s) = -2(s-3)(s+4)(2-s)$$
.  
• x-intercepts :  $f(x)=0 \Rightarrow x-3=0$ ,  $x+4=0$ ,  $2-x=0$  (respected  
 $|x=3, -4|, 2| \iff$  tike a line on (a)  
• y-intercept =  $f(o) = -2(o-3)(o+4)(2-0) = 43 \Rightarrow [(o,43)]$   
degree =  $1+1+1=3$ , leading coefficient =  $-2\cdot1\cdot1\cdot(-1)=2$  positive.  
In some problems, one or more of the factors will appear more than once when the  
function is factored. The power of a factor is called its multiplicity. So  
given  $F(s) = x^2(x-3)(x+1)$ , then the multiplicity of the third factor is 1.  
If the multiplicity of a factor is 1: the graph trouches the x-axis (looks like a line there),  
the factor have degree :  
 $(x-c)^{-1} \to (x-c)$   
If the multiplicity of a factor is codd and greater than 1: the graph crosses the x-axis  
and it looks like a cubic there.  
He factor have odd degree:  
 $(x-c)^{-1} \to (x-c)^{-1} \to - \cdots$   
Example 3: Find the x intercept(s) of the function and state the multiplicity of each.  
Indicate the possible bettere:  
 $(x-c)^{-1} \to (x-c)^{-1} \to - \cdots$   
Example 3: Find the x intercept(s) of the function and state the multiplicity of each.  
Indicate the possible bettere:  
 $(x-c)^{-1} \to (x-c)^{-1} \to - \cdots$   
Example 3: Find the x intercept(s) of the function and state the multiplicity of each.  
Indicate the possible bettere:  
 $(x-c)^{-1} \to (x-c)^{-1} \to - \cdots$   
Example 3: Find the x intercept(s) of the function and state the multiplicity of each.  
Indicate the possible bettere:  
 $(x-c)^{-1} \to (x-c)^{-1} \to - \cdots$   
 $x-idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x+5=0$   
 $x=idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x+5=0$   
 $x=idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x+5=0$   
 $x=idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x+5=0$   
 $x=idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x+5=0$   
 $x=idercept : f(x)=0 \to x^{-1}=0$ ,  $x+4=0$ ,  $x=5$  (calde)  
• y-istercept :  $f(x)=0 \to x^{-1}=0$ ,  $x=1$ ,  $(e_1f)^{-1}=-2$  regetive  $f(x)$ .





Now we'll put all of this information together to generate the graph of a polynomial function. For each problem, you'll need to state

- Follow these Steps beforc graph
- the degree of the function
- the leading coefficient of the function
- the end behavior of the function
- the *x* and *y* intercepts (and multiplicities)
- behavior of the function through each of the *x* intercepts (zeros) of the function

## Your graph should be smooth, with no sharp corners. Note that graphs of polynomial functions may have peaks or valleys, but without additional information, you will not be able to determine how high or low these points are.

**Example 4:** Find the *x* and *y* intercepts of the graph of the function. State the multiplicities of the zeros of the function. State the degree of the function and find the leading coefficient. Indicate the end behavior of the function and the behavior of the function through each zero. Use all of this information to graph the function.



**Example 5:** Write the equation of the cubic polynomial P(x) with leading coefficient -2 whose graph passes through (2, 16) and is tangent to the x-axis at the origin.

-> cubic function 
$$\rightarrow$$
 should have 3 factors  
 $\rightarrow a = -21$   
 $\Rightarrow (2,16) - given point$   
 $\Rightarrow tangent to x-axis at origin$   
 $x=0, mult. 2 - parabola shapes$   
 $f(x) = -2x^2(x-c)$   
 $f(x) = -2x^2(x-d)$ 

**Example 6:** Write the equation of a 5<sup>th</sup> degree polynomial with leading coefficient -1 given that the graph of the polynomial is tangent to the x-axis at the points 2 and 4 and the graph passes through the origin.



$$= f(x) = -x(x-2)^{2}(x-4)^{2}$$

7

**Example 7:** Given the graph of a polynomial, try to determine the equation of the polynomial.



We may not have enough time to solve all the examples here. Since graphing polynomials is a subject covered in College Algebra, we assume that you are already familiar with this subject. If you are not comfortable with graphing polynomials, please study! You can check Chapter 4 of the online textbook for College Algebra (the link is on your CASA account!). Please solve these extra problems to practice.

(Extra) Example: Find the *x* and *y* intercepts of the graph of the function. State the multiplicities of the zeros of the function. State the degree of the function and find the leading coefficient. Indicate the end behavior of the function and the behavior of the function through each zero. Use all of this information to graph the function.



(Extra) Example: Find the *x* and *y* intercepts of the graph of the function. State the multiplicities of the zeros of the function. State the degree of the function and find the leading coefficient. Indicate the end behavior of the function and the behavior of the function through each zero. Use all of this information to graph the function.



(Extra) Example: Find the *x* and *y* intercepts of the graph of the function. State the multiplicities of the zeros of the function. State the degree of the function and find the leading coefficient. Indicate the end behavior of the function and the behavior of the function through each zero. Use all of this information to graph the function.



NOTE: With some problems, you can use transformations to graph polynomial functions.

