

Popper # 19:

① A

② A


③ A

### Section 5.3a - Graphs of Secant and Cosecant Functions

Using the identity  $\csc(x) = \frac{1}{\sin(x)}$ , you can conclude that the graph of  $g$  will have a vertical asymptote whenever  $\sin(x) = 0$ . This means that the graph of  $g$  will have vertical asymptotes at  $x = 0, \pm\pi, \pm 2\pi, \dots$ . The easiest way to draw a graph of  $g(x) = \csc(x)$  is to draw the graph of  $f(x) = \sin(x)$ , sketch asymptotes at each of the zeros of  $f(x) = \sin(x)$ , then sketch in the cosecant graph.

$g(x) = \csc(x) = \frac{1}{\sin(x)}$ ; if  $\sin(x) = 0$ , then  $g(x)$  has a vertical asymptote.

Here's the graph of  $f(x) = \sin(x)$  on the interval  $\left(-\frac{5\pi}{2}, \frac{5\pi}{2}\right)$ .




$f(x) = \csc x$   
 $= \frac{1}{\sin x}$   
 can be zero

$\sin x = 0$   
 $x = 0, \pi, 2\pi, 3\pi, \dots$   
 $-\pi, -2\pi, -3\pi$

$x = k\pi$   
 Vertical Asymptotes.

Next, we'll include the asymptotes for the cosecant graph at each point where  $\sin(x) = 0$ .




$f(x) = \csc x$   
 period =  $2\pi$   
 V.A.  $x = k\pi$ .

Recall:

$\sin\left(\frac{\pi}{6}\right) = \sin\left(\frac{6\pi}{6}\right) = \frac{1}{2}$

$\sin\left(\frac{7\pi}{6}\right) = \sin\left(\frac{11\pi}{6}\right) = -\frac{1}{2}$

Now we'll include the graph of the cosecant function.



Period:  $2\pi$

Vertical Asymptote:  $x = k\pi$ ,  $k$  is an integer

$x = 0, \pi, 2\pi, 3\pi, \dots$   
 $-\pi, -2\pi, -3\pi, \dots$

$x = k\pi$  V.A.

Graph

x-intercepts: None

y-intercept: None

Domain:  $x \neq k\pi$ ,  $k$  is an integer

Range:  $(-\infty, -1] \cup [1, \infty)$

Typically, you'll just graph over one period  $(0, 2\pi)$ .

one period is always enough.

To graph  $y = A \csc(Bx - C) + D$ , first graph, **THE HELPER GRAPH**:  $y = A \sin(Bx - C) + D$ .

ex  $y = 2 \csc(2x - \pi) + 1$

Always  $\rightarrow$


Graph  $y = 2 \sin(2x - \pi) + 1$   
 then put "the parabola" shapes on top of sine function.  
 Do not forget V.A.

You'll also be able to take advantage of what you know about the graph of  $f(x) = \cos(x)$  to help you graph  $g(x) = \sec(x)$ . Using the identity  $\sec(x) = \frac{1}{\cos(x)}$ , you can conclude that the graph of  $g$  will have a vertical asymptote whenever  $\cos(x) = 0$ .

This means that the graph of  $g$  will have vertical asymptotes at  $x = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$ . The easiest way to draw a graph of  $g(x) = \sec(x)$  is to draw the graph of  $f(x) = \cos(x)$ , sketch asymptotes at each of the zeros of  $f(x) = \cos(x)$ , then sketch in the secant graph.

$$g(x) = \sec(x) = \frac{1}{\cos(x)}; \quad \text{if } \cos(x) = 0, \text{ then } g(x) \text{ has a vertical asymptote.}$$

Here's the graph of  $f(x) = \cos(x)$  on the interval  $\left(-\frac{5\pi}{2}, \frac{5\pi}{2}\right)$ .



$$f(x) = \sec x = \frac{1}{\cos x}$$

can be zero


$$\cos x = 0$$

$$x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$$

$$-\frac{\pi}{2}, -\frac{3\pi}{2}, -\frac{5\pi}{2}, \dots$$

These are V.A.  
for  $f(x) = \sec x$

Next, we'll include the asymptotes for the secant graph.



$$x = \frac{\pi}{2}, \frac{3\pi}{2} \quad \underline{\underline{\text{V.A.}}}$$

$$-\frac{\pi}{2}, -\frac{3\pi}{2}$$

$$\cos 0 = 1$$

$$\cos \frac{\pi}{3} = \frac{1}{2}$$

$$\cos \frac{2\pi}{3} = -\frac{1}{2}$$


$$\cos \frac{4\pi}{3} = -\frac{1}{2}$$

$$\cos \frac{5\pi}{3} = \frac{1}{2}$$

$$\cos \pi = -1$$

$$\cos 2\pi = 1$$

Now we'll include the graph of the secant function.



Period:  $2\pi$

Vertical Asymptote:

$x = k\pi/2$  is an odd integer,  $k$  is odd

x-intercepts: None

y-intercept:  $(0, 1)$

Domain:  $x \neq k\pi/2$ ,  $k$  is an odd integer

Range:  $(-\infty, -1] \cup [1, \infty)$

$$x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$$

$$x = \frac{-\pi}{2}, \frac{-3\pi}{2}, \frac{-5\pi}{2}, \dots$$

$$x = \frac{k\pi}{2}, k \text{ odd}$$

graph

Typically, you'll just graph over one period  $(0, 2\pi)$ .

To graph  $y = A \sec(Bx - C) + D$ , first graph, **THE HELPER GRAPH**:  $y = A \cos(Bx - C) + D$ .

$y = A \sec(Bx - C) + D$  Always  $\rightarrow$

graph  $y = A \cos(Bx - C) + D$   
and then put "parabola"  
shapes on top of cosine!  
Do not forget V.A.

Example 1: Sketch  $f(x) = 4 \sec\left(\frac{x}{2}\right)$

Helper graph:


$$y = 4 \cos\left(\frac{x}{2}\right)$$

Vertical Stretch

$$B = \frac{1}{2}$$

Horizontal Stretch

$$\text{period} = \frac{2\pi}{\frac{1}{2}} = 4\pi$$



Note  $f(x) = 4 \sec\left(\frac{x}{2}\right)$  has V.A. at

$$x = \pi, 3\pi, 5\pi, \dots$$

$$- \pi, -3\pi, -5\pi$$

because  $\cos\left(\frac{x}{2}\right) = 0$  at those points !!!

Example 2: Sketch  $f(x) = -2 \csc\left(\frac{\pi x}{2} - \frac{\pi}{2}\right)$

Helper graph:

$$y = -2 \sin\left(\frac{\pi}{2}x - \frac{\pi}{2}\right)$$


•  $A = -2 \rightarrow$  vertical stretching and reflection

•  $B = \frac{\pi}{2} \rightarrow$  horizontal shrinking  
 period =  $\frac{2\pi}{\pi/2} = 4$

•  $C = \frac{\pi}{2} \rightarrow$  horizontal shift

by  $\frac{C}{B} = \frac{\pi/2}{\pi/2} = 1$  unit right.

$\Rightarrow$  Then put  $y = -2 \csc\left(\frac{\pi}{2}x - \frac{\pi}{2}\right)$ .



Note  $f(x) = -2 \csc\left(\frac{\pi}{2}x - \frac{\pi}{2}\right)$  has V.A.


at  $x = 1, 3, 5, 7, \dots$   
 $-1, -3, -5, -7, \dots$

because  $\sin\left(\frac{\pi}{2}x - \frac{\pi}{2}\right) = 0$  at these values.

To be continued on Thursday, 03/24

**Example 3:** Give an equation of the form  $y = A \csc(Bx - C) + D$  and  $y = A \sec(Bx - C) + D$  that could describe the following graph.

Complete the helper graph!



$$f(x) = 3 \csc(x) - 1$$


Let's find helper:

$$y = A \sin(Bx - C) + D$$

- $A = \frac{6}{2} = 3 \Rightarrow \boxed{A=3}$
- $\boxed{B=1}$  since period =  $2\pi$ .
- $\boxed{C=0}$  since no shift
- $\boxed{D=-1}$ , 1 unit down  
 $D = \frac{-4+2}{2} = -1$

**Exercise:** Give an equation of the form  $y = A \csc(Bx - C) + D$  and  $y = A \sec(Bx - C) + D$  that could describe the following graph.

Viewing as a **secant** function:



• amplitude =  $\frac{7 - (-1)}{2} = \frac{8}{2} = 4 \Rightarrow A = 4$

• period =  $8 = \frac{2\pi}{B} \Rightarrow B = \frac{\pi}{4}$

• shift 1 to right,  $\frac{C}{B} = 1 \Rightarrow C = B = \frac{\pi}{4}$


• Vertical shift =  $\frac{(-1) + 7}{2} = 3 \rightarrow D = 3$

$$y = 4 \cos\left(\frac{\pi}{4}(x-1)\right) + 3$$

$$= 4 \cos\left(\frac{\pi x}{4} - \frac{\pi}{4}\right) + 3$$


$$\Rightarrow y = 4 \sec\left(\frac{\pi x}{4} - \frac{\pi}{4}\right) + 3$$

**Example 3:** Give an equation of the form  $y = A\csc(Bx - C) + D$  and  $y = A\sec(Bx - C) + D$  that could describe the following graph.



**Exercise:** Give an equation of the form  $y = A\csc(Bx - C) + D$  and  $y = A\sec(Bx - C) + D$  that could describe the following graph

Viewed as cosecant Function



$$f(x) = 4 \sin\left(\frac{\pi}{4}x + \frac{\pi}{4}\right) + 3$$

$$\Rightarrow f(x) = 4 \csc\left(\frac{\pi}{4}x + \frac{\pi}{4}\right) + 3$$

Let's find helper:

$$y = A \sin(Bx - C) + D$$

- amplitude =  $\frac{8}{2} = 4$

$$A = 4$$

- period =  $3 - (-5) = 8$

$$\frac{2\pi}{B} = 8 \Rightarrow B = \frac{\pi}{4}$$

- shifted 1 unit left

$$\frac{C}{B} = -1 \Rightarrow C = -\frac{\pi}{4}$$

- $D = 3$  units up!



**Exercise:** Find the vertical asymptotes of:

$$a) f(x) = 2\sec\left(\frac{x}{2} - \pi\right) = 2 \cdot \frac{1}{\cos\left(\frac{x}{2} - \pi\right)}$$

Vertical Asymptotes:

$$\cos\left(\frac{x}{2} - \pi\right) = 0 \Rightarrow$$

$$\Rightarrow x = k\pi, \quad k \text{ odd}$$

$$\frac{x}{2} - \pi = \frac{\pi}{2} \Rightarrow \frac{x}{2} = \frac{3\pi}{2} \Rightarrow x = 3\pi$$

$$\frac{x}{2} - \pi = \frac{3\pi}{2} \Rightarrow \frac{x}{2} = \frac{5\pi}{2} \Rightarrow x = 5\pi$$

$$\frac{x}{2} - \pi = -\frac{\pi}{2} \Rightarrow \frac{x}{2} = \frac{\pi}{2} \Rightarrow x = \pi$$

$$\frac{x}{2} - \pi = -\frac{3\pi}{2} \Rightarrow \frac{x}{2} = -\frac{\pi}{2} \Rightarrow x = -\pi$$

$$b) f(x) = 2\csc\left(x - \frac{\pi}{4}\right)$$

$$= 2 \cdot \frac{1}{\sin\left(x - \frac{\pi}{4}\right)}$$

Vertical asymptotes:  $\sin\left(x - \frac{\pi}{4}\right) = 0$

$$x - \frac{\pi}{4} = 0 \Rightarrow x = \frac{\pi}{4}$$

$$x - \frac{\pi}{4} = -\pi \Rightarrow x = -\frac{3\pi}{4}$$

$$x - \frac{\pi}{4} = \pi \Rightarrow x = \frac{5\pi}{4}$$

$$x - \frac{\pi}{4} = -2\pi \Rightarrow x = -\frac{7\pi}{4}$$

$$x - \frac{\pi}{4} = 2\pi \Rightarrow x = \frac{9\pi}{4}$$

$$x = \frac{k\pi}{4}, \quad k = \dots, -7, -3, 1, 5, 9, \dots$$

$$\text{or } x = \frac{\pi}{4} + k\pi, \quad k \text{ integer.}$$