

Math 3331 Differential Equations

2.7 Existence and Uniqueness of Solutions

Blerina Xhabli

Department of Mathematics, University of Houston

`blerina@math.uh.edu`
`math.uh.edu/~blerina/teaching.html`



2.7 Existence and Uniqueness of Solutions

- Existence of Solution
 - Existence for Linear Equation
 - Existence when the Right-hand Side is Discontinuous
- Interval of Existence of a Solution
- Uniqueness of Solution
- Worked out Examples from Exercises:
 - 1, 3, 5, 7



Existence and Uniqueness Theorem

Basic Existence and Uniqueness Theorem (EUT):

Suppose $f(t, x)$ is defined and continuous, and has a continuous partial derivative $\partial f(t, x)/\partial x$ on a rectangle R in the tx -plane. Then, given any initial point (t_0, x_0) in R , the initial value problem

$$x' = f(t, x), \quad x(t_0) = x_0$$

has a unique solution $x(t)$ defined in an interval containing t_0 . Furthermore, the solution will be defined at least until the solution leaves R .



Example

Ex.: $tx' = x + 3t^2 \Rightarrow x' = x/t + 3t$

- f and $\partial f/\partial x$ are defined and continuous for any (t, x) if $t \neq 0$
- General solution (use Sec. 2.6):

$$x(t) = 3t^2 + Ct$$

- For any C : $x(0) = 0$, hence
 - no solution for $x(0) = x_0 \neq 0$
 - ∞ solutions for $x(0) = 0$

- Solution for $x(t_0) = x_0$, $t_0 > 0$:

$$3t_0^2 + Ct_0 = x_0 \Rightarrow C = x_0/t_0 - 3t_0$$

$$\Rightarrow x(t) = 3t^2 + (x_0/t_0 - 3t_0)t$$

unique solution with IoE $(0, \infty)$

- EUT applies to any rectangle that is not intersected by the vertical line $t = 0$.




Figure 1 All solutions of (7.2) pass through $(0,0)$.



Example: Non-Uniqueness of Solution

Ex.: $x' = x^{1/3}$

S.o.V.: $\int x^{-1/3} dx = (3/2)x^{2/3} = t + D$

$\Rightarrow x_{\pm}(t) = \pm[(2/3)t + C]^{3/2} \quad (C = 2D/3)$



- Let $C = 0 \Rightarrow x_{\pm}(0) = 0$
- Other solution with $x(0) = 0$:

$$x(t) = 0$$

\Rightarrow At least 3 solutions for IC

$$x(0) = 0$$

- EUT doesn't apply to any rectangle that is intersected by the horizontal line $x = 0$



Interval of Existence

Interval of Existence:

Largest interval in which a solution of a first order ODE can be defined.



Example

$$\text{Ex.: } x' = -x^2, \quad x(0) = x_0$$

$$\text{S.o.V.: } -\int (1/x^2)dx = 1/x = t + C$$

$$\Rightarrow x = 1/(t + C)$$

$$x(0) = 1/C = x_0 \Rightarrow C = 1/x_0$$

$$\Rightarrow x(t) = x_0/(1 + x_0 t)$$

$$\text{If } \left. \begin{array}{l} x_0 > 0 \\ x_0 < 0 \end{array} \right\} \Rightarrow \text{IoE: } \left\{ \begin{array}{l} (-1/x_0, \infty) \\ (-\infty, -1/x_0) \end{array} \right.$$

$$\text{If } x_0 = 0 \Rightarrow x(t) = 0, \text{ IoE: } (-\infty, \infty)$$

- $f(t, x) = -x^2$ satisfies hypotheses of EUT in any rectangle

\Rightarrow Unique solution for any x_0

- $x(t)$ leaves any rectangle in finite time

\Rightarrow Solution is not defined for *all* reals if $x_0 \neq 0$



Existence When the RHS is Discontinuous

Ex.: IVP $y' = -2y + f(t)$, $y(0) = 3$

$$f(t) = \begin{cases} 0 & \text{if } t < 1 \\ 5 & \text{if } t \geq 1 \end{cases}$$

$$t < 1: y' = -2y \Rightarrow y(t) = 3e^{-2t}$$

$$\text{For } t \rightarrow 1: y(1) = 3e^{-2}$$

Continue solution beyond $t = 1$:


$$t \geq 1: y' = -2y + 5, y(1) = 3e^{-2}$$

$$\begin{aligned} \Rightarrow y(t) &= 3e^{-2t} + e^{-2t} \int_1^t e^{2t'} 5 dt' \\ &= 5/2 + (3 - 5e^2/2)e^{-2t} \end{aligned}$$

Combine:

$$y(t) = \begin{cases} 3e^{-2t} & \text{if } t \leq 1 \\ 5/2 + (3 - 5e^2/2)e^{-2t} & \text{if } t \geq 1 \end{cases}$$

- f is discontinuous at $t = 1$, but unique solution exists for all t
- $y'(t)$ is discontinuous at $t = 1$



Exercise 2.7.1

Ex. 1: $y' = 4 + y^2$, $y(0) = 1$. Does IVP have a unique solution?

Yes, because $f = 4 + y^2$ and $\partial f/\partial y = 2y$ are continuous everywhere.



Exercise 2.7.3

Ex. 3: $y' = t \tan^{-1}(y)$, $y(0) = 2$. Does IVP have a unique solution?

Yes (as Ex. 1).



Exercise 2.7.5

Ex. 5: $x' = t/(x + 1)$, $x(0) = 0$. Does IVP have a unique solution?

Yes, because f and $\partial f/\partial x = -t/(x + 1)^2$ are continuous in any rectangle away from the horizontal line $x = -1$, and $x(0) \neq -1$.



Exercise 2.7.7

Ex. 7: $ty' - y = t^2 \cos t$, $y(0) = -3$.

- (i) Find general solution and sketch several solutions.
 (ii) Show IVP has no solution and explain why this doesn't contradict EUT.

Answer (i): $y' - y/t = t \cos t$, use integrating factor:

$$u(t) = \exp\left(-\int (1/t)dt\right) = \exp(-\ln t) = 1/t$$

$$\Rightarrow (y/t)' = \cos t \Rightarrow y/t = \sin t + C \Rightarrow y(t) = t \sin t + Ct$$

Answer (ii): Since $y(0) = 0$ for any C , there is no solution that satisfies $y(0) = -3$. This doesn't contradict EUT because f is not continuous at $t = 0$.

