

# Math 3331 Differential Equations

## 6.1 Euler's Method

**Blerina Xhabli**

Department of Mathematics, University of Houston

`blerina@math.uh.edu`  
`math.uh.edu/~blerina/teaching.html`



## 6.1 Euler's Method

- Basic Idea
  - Euler's Method: Basic Idea
  - Euler's Method: Iteration Scheme
- Examples
- Errors
  - Euler's Method: Errors
  - Errors in Euler's Method: First Order
- Worked out Examples from Exercises:
  - 1, 7



# Euler's Method: Basic Idea

## Basic Idea

- ODE:  $y' = f(t, y)$
- Assume  $y(t)$  is known
- For small  $h$  approximate

$$\frac{y(t+h) - y(t)}{h} \approx y'(t) \\ = f(t, y(t))$$


$$\Rightarrow y(t+h) \approx y_{ap}(t+h)$$

where

$$y_{ap}(t+h) = y(t) + h f(t, y(t))$$

- **Truncation Error:**

$$|y(t+h) - y_{ap}(t+h)|$$



# Euler's Method: Iteration Scheme

## Iteration Scheme

$$\text{IVP: } y' = f(t, y), \quad y(t_0) = y_0$$

Approximate  $y(t_k) \approx y_k$  at  $t_k$ :

$$y_1 = y_0 + h f(t_0, y_0), \quad t_1 = t_0 + h$$

$$y_2 = y_1 + h f(t_1, y_1), \quad t_2 = t_1 + h$$

$$\vdots$$

$$y_{k+1} = y_k + h f(t_k, y_k)$$

$$t_{k+1} = t_k + h$$



# Example 1

**Ex:** Approximate the solution to

$$y' = y, \quad y(0) = 1$$

in  $0 \leq t \leq 1$ . Start:  $t_0 = 0, y_0 = 1$

**h = 1**

$$y_1 = y_0 + h f(0, 1) = 1 + 1 \cdot 1 = 2$$

$$t_1 = t_0 + h = 0 + 1 = 1$$

**h = 0.5**

$$y_1 = 1 + 0.5 \cdot 1 = 1.5$$

$$t_1 = 0 + 0.5 = 0.5$$

$$y_2 = 1.5 + 0.5 \cdot 1.5 = 2.25$$

$$t_2 = 0.5 + 0.5 = 1$$

**h = 0.25**

$$y_1 = 1 + 0.25 \cdot 1 = 1.25$$

$$t_1 = 0 + 0.25 = 0.25$$

$$y_2 = 1.25 + 0.25 \cdot 1.25 = 1.5625$$

$$t_2 = 0.25 + 0.25 = 0.5$$

$$y_3 = 1.5625 + 0.25 \cdot 1.5625$$


$$= 1.953125$$

$$t_3 = 0.5 + 0.25 = 0.75$$

$$y_4 = 1.953125 + 0.25 \cdot 1.953125$$

$$= 2.44140625$$

$$t_4 = 0.75 + 0.25 = 1$$



# Example 2

**Ex:** Approximate the solution to

$$y' = t - y, \quad y(0) = 0.5$$

in  $0 \leq t \leq 1$  using  $h = 0.25$

**Start:**  $y_0 = 0.5, t_0 = 0$

$$y_1 = 0.5 + 0.25 \cdot (0 - 0.5) = 0.375$$

$$t_1 = 0 + 0.25 = 0.25$$

$$\begin{aligned} y_2 &= 0.375 + 0.25 \cdot (0.25 - 0.375) \\ &= 0.3438 \end{aligned}$$

$$t_2 = 0.25 + 0.25 = 0.5$$

$$\begin{aligned} y_3 &= 0.3438 + 0.25 \cdot (0.5 - 0.3438) \\ &= 0.3828 \end{aligned}$$

$$t_3 = 0.5 + 0.25 = 0.75$$

$$\begin{aligned} y_4 &= 0.3828 + 0.25 \cdot (0.75 - 0.3828) \\ &= 0.4746 \end{aligned}$$

$$t_4 = 0.75 + 0.25 = 1$$




# Euler's Method: Errors

## Errors

Three error sources:

- Truncation error at each Euler step
- Propagated (accumulated) truncation error
- Roundoff error (not controllable)



# Errors in Euler's Method: First Order

**Ex.:**  $y' = t - y$ ,  $y(0) = .5$

Approximate  $y(1)$  for stepsizes

$$h = 1/m, \quad m = 1, 2, 4, 8, 16, 32$$

**Exact Value:**  $y(1) = 0.5518$

**Error:**  $E(h) = |y(1) - y_m|$

$h$	$y_m$	$E(h)$
1	0	0.5518
1/2	0.375	0.1768
1/4	0.4746	0.0772
1/8	0.5154	0.0364
1/16	0.5341	0.0177
1/32	0.5431	0.0087

$$E(h/2) \approx E(h)/2 \Rightarrow E(h) \approx Ch$$

**Theorem:** There  $\exists C > 0$  s.t.

$$E(h) \leq Ch$$

(Euler method is first order method)





# Exercise 6.1.1

**Ex. 1:**  $y' = ty$ ,  $y(0) = 1$ .

Compute five Euler-iterates for  $h = 0.1$ .  
Arrange computation and results in a table.

$k$	$t_k$	$y_k$	$f(t_k, y_k) = t_k y_k$	$h$	$f(t_k, z_k)h$
0	0	1	0	0.1	0
1	0.1	1	0.1000	0.1	0.0100
2	0.2	1.0100	0.2020	0.1	0.0202
3	0.3	1.0302	0.3091	0.1	0.0309
4	0.4	1.0611	0.4244	0.1	0.0424
5	0.5	1.1036	0.5518	0.1	0.0552



# Exercise 6.1.7 (i)

**Ex. 7:**  $y' + 2xy = x$ ,  $y(0) = 8$

- (i) Compute Euler-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and Euler approximations as points.

```

h=0.2;
m=1/h;x=0;y=8;
xv=x;yv=y;
for k=1:m
    f=-2*x*y+x;
    y=y+h*f;yv=[yv y];
    x=x+h;xv=[xv x];
end
x0_2=xv;y0_2=yv;
  
```

(i) In Matlab, Euler approximation for  $h = 0.2$  is computed and stored in arrays  $x0\_2$ ,  $y0\_2$  via

Analogously for  $h = 0.1$  and  $h = 0.05$  (arrays  $x0\_1$ ,  $y0\_1$  and  $x0\_05$ ,  $y0\_05$ ).



# Exercise 6.1.7 (ii)

**Ex. 7:**  $y' + 2xy = x$ ,  $y(0) = 8$

- (i) Compute Euler-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and Euler approximations as points.

(ii) Variation of Parameter:

$$y'_h = -2xy \Rightarrow$$

$$y_h(x) = \exp\left(\int_0^x (-2x)dx\right) = e^{-x^2}$$

$$\begin{aligned} y(x) &= y_h(x) \left( 8 + \int_0^x [f(\xi)/y_h(\xi)]d\xi \right) \\ &= 8e^{-x^2} + e^{-x^2} \int_0^x \xi e^{\xi^2} d\xi \\ &= 8e^{-x^2} + e^{-x^2} (e^{x^2} - 1)/2 \\ &= (15/2)e^{-x^2} + 1/2 \end{aligned}$$



# Exercise 6.1.7 (iii)

**Ex. 7:**  $y' + 2xy = x$ ,  $y(0) = 8$

- (i) Compute Euler-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and Euler approximations as points.

(iii) Matlab plot commands:

```
x=linspace(0,1,100);
y=1/2+15/2*exp(-x.^2);
plot(x0_2,y0_2,'ko',x0_1,y0_1,'k*',...
      x0_05,y0_05,'k+',x,y,'k'),
xlabel('x'),ylabel('y')
axis([0 1 3.5 8])
```

