

# Math 3331 Differential Equations

## 6.3 Numerical Error Comparisons

**Blerina Xhabli**

Department of Mathematics, University of Houston

`blerina@math.uh.edu`  
`math.uh.edu/~blerina/teaching.html`



## 6.3 Numerical Error Comparisons

- Numerical Error Comparisons
  - Examples
  - Least Square Fit
- Worked out Examples from Exercises:
  - 3, 7,



# Example

**Ex.**  $y' = t - y, y(0) = 0.5$

$$y(1) \approx y_m \rightarrow E(h) = |y(1) - y_m|$$

$$h = 1/m, \quad m = 1, 2, 4, 8, 16, 32$$

$h$	EM	RKM2	RKM4
1	0.5518	0.198181	0.010680838
1/2	0.1768	0.034118	0.000437105
1/4	0.0772	0.006974	0.000022137
1/8	0.0364	0.001581	0.000001246
1/16	0.0177	0.000377	0.000000074
1/32	0.0087	0.000092	0.000000005

$E(h)$  for



EM: Euler Method

RKM2: 2nd order RKM

RKM4: 4th order RKM



# Example (cont.): Least Square Fit of $E(h)$



# Exercise 6.2.3

**Ex. 3:**  $y' = ty$ ,  $y(0) = 1$ .

Compute five RK2-iterates for  $h = 0.1$ .  
Arrange computation and results in a table.

$k$	$t_k$	$y_k$	$s_l$	$s_r$	$h$	$h(s_l + s_r)/2$
0	0	1	0	0.1	0.1	0.005
1	0.1	1.0050	0.1005	0.2030	0.1	0.0152
2	0.2	1.0202	0.2040	0.3122	0.1	0.0258
3	0.3	1.0460	0.3138	0.4309	0.1	0.0372
4	0.4	1.0832	0.4333	0.5633	0.1	0.0498
5	0.5	1.1331	0.5665	0.7138	0.1	0.0640



# Exercise 6.2.7 (i)

**Ex. 7:**  $z' + z = \cos x$ ,  $z(0) = 1$

- (i) Compute RK2-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and RK2 approximations as points.

(i) In Matlab, the RK2 approximation for  $h = 0.2$  is computed and stored in arrays  $x0\_2$ ,  $z0\_2$  via

Analogously for  $h = 0.1$  and  $h = 0.05$  (arrays  $x0\_1$ ,  $z0\_1$  and  $x0\_05$ ,  $z0\_05$ ).

```

h=0.2;
m=1/h;x=0;z=1;
xv=x;zv=z;
for k=1:m
    sl=cos(x)-z;
    sr=cos(x+h)-(z+sl*h);
    z=z+h*(sl+sr)/2;zv=[zv z];
    x=x+h;xv=[xv x];
end
x0_2=xv;z0_2=zv;

```



## Exercise 6.2.7 (ii)

**Ex. 7:**  $z' + z = \cos x$ ,  $z(0) = 1$

- (i) Compute RK2-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and RK2 approximations as points.

(ii) Variation of Parameter:

$$z'_h = -z \Rightarrow z_h(x) = e^{-x}$$

$$\begin{aligned} z(x) &= e^{-x} + \int_0^x e^{\xi} \cos(\xi) d\xi \\ &= (\cos x + \sin x + e^{-x})/2 \end{aligned}$$




# Exercise 6.2.7 (iii)

**Ex. 7:**  $z' + z = \cos x$ ,  $z(0) = 1$

- (i) Compute RK2-approximations in  $0 \leq x \leq 1$  for  $h = 0.2$ ,  $h = 0.1$ ,  $h = 0.05$ .
- (ii) Find exact solution
- (iii) Plot exact solution as curve and RK2 approximations as points.

(iii) Plot:

(see CN Sec. 6.1 for commands)





# Exercise 6.2.7a (i)

**Ex. 7a:**  $z' + z = \cos x$ ,  $z(0) = 1$

- (i) Compute RK4-approximation in  $0 \leq x \leq 1$  for  $h = 0.2$ .
- (iii) Plot exact solution as curve and RK4 approximation as points.

(i) RK4 approximation for  $h = 0.2$  is computed and stored in arrays  $xv$ ,  $zv$ :

```

h=0.2;
m=1/h;x=0;z=1;
xv=x;zv=z;
for k=1:m
    s1=cos(x)-z;
    s2=cos(x+h/2)-(z+s1*h/2);
    s3=cos(x+h/2)-(z+s2*h/2);
    s4=cos(x+h)-(z+s3*h);
    z=z+h*(s1+2*s2+2*s3+s4)/6;
    zv=[zv z];
    x=x+h;xv=[xv x];
end
  
```



# Exercise 6.2.7a (ii)

**Ex. 7a:**  $z' + z = \cos x$ ,  $z(0) = 1$

- (i) Compute RK4-approximation in  $0 \leq x \leq 1$  for  $h = 0.2$ .  
 (iii) Plot exact solution as curve and RK4 approximation as points.

(iii) Matlab plot commands:

```
x=linspace(0,1,100);
z=1/2*(cos(x)+sin(x)+exp(-x));
plot(xv,zv,'ko',x,z,'k'),
xlabel('x'),ylabel('z'),
axis([0 1 0.86 1])
```

