

# MATH 1314

Section 3.1

# Functions: Basic Ideas

Basic Definition: (Non-formal):

Plug in one  $x$ -value, you get one answer.

The rest of this course deals with **functions**.

**Definition:** A function,  $f$ , is a rule that assigns to each element  $x$  in a set  $A$  exactly one element, called  $f(x)$ , in a set  $B$ .

Functions are so important that we use a special notation when working with them. We'll write  $f(x)$  to denote the value of function  $f$  at  $x$ . We read this as "f of x." We can use letters other than  $f$  to denote a function, so you may see a function such as  $g(x)$ ,  $h(x)$  or  $P(x)$ .

- You cannot have one  $x$  with 2 answers

• You can have two  $x$  with the same answer

$$y = 2x + 5$$

$$f(x) = 2x + 5$$

$f(x)$ : "f of x"

- This is a function
- Ind. variable is  $x$  Named as "f"

**Definition:** The set A is called the **domain** and is the set of all valid inputs for the function.

**Definition:** The set B is called the **range** and is the set of all possible values of  $f(x)$  as  $x$  varies throughout the domain.

Sets A and B will consist of real numbers.

Domain: All inputs (that evaluate to answers). Typically,  $x$ -values.


Range: All possible output of a function. Typically,  $y$ -values.

Sometimes, Domain and Range will be All Real Numbers.

Sometimes, they will be restrictions based on the nature of the function.

**Example 1:**

a. Given:



Is  $f$  a function?

$$f(A) = 2$$


$$f(B) = 1$$

$$f(C) = 2$$

This is a function.

Every element of the domain has exactly 1 answer.

b. Given:



Is  $g$  a function?

$$\begin{aligned}g(A) &= 1 \\g(B) &= 1 \\g(B) &= 3 \\g(C) &= 2\end{aligned}$$

B has 2 different answers.

$g$  is Not a function.

(It only takes part of the domain to make this not a function)

Next we'll consider some things you'll need to be able to do when working with functions. First, you'll need to be able to evaluate all types of functions when given a specific value for the variable.

**Example 2:** Let  $f(x) = x^2 - 4x$  Calculate

a.  $f(-3) \rightarrow$  Replace  $x$  with  $-3$  in  $f(x)$

$$f(-3) = (-3)^2 - 4(-3)$$

$$\begin{array}{c} \uparrow \\ x \end{array} = 9 + 12 = 21 \rightarrow (-3, 21)$$

b.  $-2f(x)$  (Notice:  $-2$  is outside the function)

$$-2f(x) = -2(x^2 - 4x)$$

$$= -2x^2 + 8x$$

$\rightarrow$  Replacing  $x$  with  $3x$

$$\begin{aligned} \text{c. } f(3x) &= (3x)^2 - 4(3x) \\ &= \boxed{9x^2 - 12x} \end{aligned}$$

$\rightarrow$  Replace  $x$  with  $x+2$

$$\begin{aligned} \text{d. } f(x+2) &= (x+2)^2 - 4(x+2) \\ &= (x+2)(x+2) - 4(x+2) \\ &= x^2 + 2x + 2x + 4 - 4x - 8 \\ &= \boxed{x^2 - 4} \end{aligned}$$

Example 3: Suppose  $g(x) = \begin{cases} 2x - 6, & x < -2 \\ x^2 + 2x + 3, & x \geq -2 \end{cases}$  Calculate the following

piecewise function

a.  $g(-5)$

Is  $-5 < -2$ ?  $\rightarrow 2x - 6$   
 $-5 \geq -2$

$$g(-5) = 2(-5) - 6 = -10 - 6 = \boxed{-16}$$

$(-5, -16)$

Popper #9:

1.  $g(-2) = (-2)^2 + 2(-2) + 3 =$   
 $-2 \geq -2$   $4 - 4 + 3 = \boxed{3}$

2.  $g(3) = (3)^2 + 2(3) + 3$   
 $3 \geq -2$   $= 9 + 6 + 3 = \boxed{18}$

a. 18

b. 3

c. -10

d. 0

B

ZA

## Finding the Domain of a Function

Recall: The domain is the set of all real numbers for which the expression is defined as a real number. Exclude from a function's domain real numbers that cause division by zero or real numbers that result in an even root of a negative number.

We express the set of real numbers as  $(-\infty, \infty)$ .

The domain of any polynomial function is  $(-\infty, \infty)$ .

2 Things we look for to eliminate:

- Zeros in the denominator
- Negative inside  $\sqrt[n]{\quad}$  powered roots.

(If neither of these two occur, then the domain is  $(-\infty, \infty)$ .)



Example 4: Find the domain of each function below and express your answer in interval notation.

a.  $f(x) = -17$  Domain:  $(-\infty, \infty)$

- No fractions
- No roots

c.  $h(x) = \frac{5x}{2x-8}$

Fraction

Domain:  
 $(-\infty, 4) \cup (4, \infty)$

Denominator  $\neq 0$

$$2x + 8 \neq 0$$

$$\frac{2x + 8}{+8 \quad +8}$$


---


$$\frac{\cancel{2}x + \cancel{8}}{\cancel{2}} \neq \frac{\cancel{8}}{\cancel{2}}$$


---


$$x \neq 4$$

b.  $f(x) = 3x - 4$  Domain:  $(-\infty, \infty)$

- No fractions
- No roots

d.  $f(x) = \frac{x-1}{2x-6}$

Domain:  
 $(-\infty, 3) \cup (3, \infty)$

Denominator  $\neq 0$

$$2x + 6 \neq 0$$

$$\frac{2x + 6}{+6 \quad +6}$$


---


$$\frac{\cancel{2}x + \cancel{6}}{\cancel{2}} \neq \frac{\cancel{6}}{\cancel{2}}$$


---


$$x \neq 3$$

$$e. p(x) = \frac{x^2 - 16}{x^2 - 4x - 12}$$

Domain:

$$(-\infty, -2) \cup (-2, 6) \cup (6, \infty)$$

Denominator  $\neq 0$

$$x^2 - 4x - 12 \neq 0$$

$$(x - 6)(x + 2) \neq 0$$

$$\begin{array}{r} x + 6 \neq 0 \\ +6 \quad +6 \\ \hline x \neq -6 \end{array}$$

$$\begin{array}{r} x + 2 \neq 0 \\ -2 \quad -2 \\ \hline x \neq -2 \end{array}$$

$$f. h(x) = \sqrt{x}$$

Root inside  $\geq 0$   
 $x \geq 0$   
 $[0, \infty)$

$$g. j(x) = \frac{\sqrt{x - 5}}{x + 2}$$

• Root: inside  $\geq 0$

$$\begin{array}{r} x - 5 \geq 0 \\ +5 \quad +5 \\ \hline x \geq 5 \end{array}$$

• Denominator  $\neq 0$

$$\begin{array}{r} x + 2 \neq 0 \\ -2 \quad -2 \\ \hline x \neq -2 \end{array}$$

$$\text{Domain: } [5, \infty)$$

$$h. k(x) = \frac{\sqrt{x - 5}}{x - 8}$$

• Root:  $x - 5 \geq 0$   
 $x \geq 5$

• Den:  $x - 8 \neq 0$   
 $x \neq 8$

$$[5, 8) \cup (8, \infty)$$

Note: odd roots have domain:  $(-\infty, \infty)$

# Popper #9...continued

3.  $q(x) = \sqrt{x-4}$  D

$$\begin{array}{r} x-4 \geq 0 \\ +4 \quad +4 \\ \hline x \geq 4 \end{array} \quad [4, \infty)$$

4.  $f(x) = \sqrt[3]{2x+4}$  C

↳ Like Root

odd roots have domain of  $(-\infty, \infty)$

→ Even Root

5.  $g(x) = \sqrt[10]{42-2x}$  A

$$\begin{array}{r} 42-2x \geq 0 \\ -42 \quad -42 \\ \hline -2x \geq -42 \\ \hline -2 \quad -2 \\ \hline x \leq 21 \end{array}$$

a.  $(-\infty, 21]$

b.  $[21, \infty)$

c.  $(-\infty, \infty)$

d.  $[4, \infty)$

$(-\infty, 21]$