# MATH 1314

Section 4.1

string of torms that are added on Polynomial Functions:

Subtracted

Positive Mole Munchers

A polynomial function is a function of the form

Example

Geograph

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^n + \dots + a_1 x^1 + a_n$$

where  $a_n \neq 0$ ,  $a_0, a_1, ..., a_n$  are real numbers and n is a whole number.

The degree of the polynomial function is n. We call the term  $a_n x^n$  the leading term, and  $a_n$  is called the leading coefficient. Degree Largest Exponent

[P(0) = a0] Leading Torm Term with largest constant term term without the x

Our objectives in working with polynomial functions will be, first, to gather information about the graph of the function and, second, to use that information to generate a reasonably good graph without plotting a lot of points. In later examples, we'll use information given to us about the graph of a function to write its equation.

P(x)=  $-3x^4 + 2x^3 - 5x + 2$ (Pegree: 4)

Constant Term: 2

### Factored Form:

$$p(x) = 2(x - 4)^{2}(x^{2} + 1)^{3}$$

Leading Term: "First" Step of Folk

2 (x) (x2) [Each Langert Pours x and who outside experient]

 $2\left(x^{2}\chi\chi^{6}\right)$ 

2x8 Degree 8

Constant Term >> >-intercept

$$P(0) = 2(0-4)^{2}(0^{2}+1)^{3}$$

$$= 2(-4)^{2}(1)^{3}$$


$$= 2(16)(1) = 32, \quad (0,32)$$

#### **Graph Properties of Polynomial Functions**

Let *P* be any *n*th degree polynomial function with real coefficients. The graph of *P* has the following properties.

- 1. *P* is continuous for all real numbers, so there are no breaks, holes, jumps in the graph.  $\mathcal{D}: (-\infty, \infty)$
- 2. The graph of *P* is a smooth curve with rounded corners and no sharp corners.
- 3. The graph of P has at most n x-intercepts.
- 4. The graph of P has at most n-1 turning points.





Example 1: Given the following polynomial functions, state the leading term, the degree of the polynomial and the leading coefficient, constant term

a.  $P(x) = 6x^4 - 4x^3 + 7x - 2$ Leading Torm 6x4 Degae: 4 Lending Coefficient 6

constant term - 7

b.  $P(x) = (3x + 4)(x + 1)^2(x - 5)^3$ 

Leading Term:  $(3\times)(x)^2(x)^3 = 3\times -x^2 \cdot x^3 = 3\times 6$ 

Degree 6

Lealing Coefficient: 3


( onstant Term: P(a) - (3.014) (0+1) (0-5) = 4(1) 2(-5) = 4(1)(-125)

-500

We'll start with the shapes of the graphs of functions of the form  $f(x) = x^n$ , n > 0.

You should be familiar with the graphs of  $f(x) = x^2$  and  $g(x) = x^3$ .

The graph of  $f(x) = x^n$ , n > 0., n is even, will resemble the graph of  $f(x) = x^2$ , and the graph of  $f(x) = x^n$ , n > 0, n is odd, will resemble the graph of  $f(x) = x^3$ .



Next, you will need to be able to describe the end behavior of a function.


### End Behavior of Polynomial Functions


The behavior of a graph of a function to the far left or far right is called its end behavior.

The end behavior of a polynomial function is revealed by the leading term of the polynomial End Behavior: what is


function.

1. Even-degree polynomials look like  $y = \pm x^2$ 





### End behavior of Polynomials:



### Popper 20

For the following, refer to  $f(x) = -8x_3 - 2x_2 + 9x - 3$ 

- What is the degree of the polynomial?
- First a.
- b. Sixth
- c. Third
- d. Eighth

- 2. What is the leading coefficient?
- a.

b. -2

c. 9

d. -3

- 3. What is the end behavior on the left?
- Rising a.

- b. Falling
- 4. What is the end behavior on the right?
- Rising a.

b. Falling

Leading Coefficient: -8 Degree: 3

Left: Rising Right: Falling

Describe the End Behavior of the following:

$$g(x) = (x^{2} + 3)(x - 2)^{4}$$
Leading Term:  $(\chi^{2})(\chi)^{4} = \chi^{2} \cdot \chi^{4} = \chi^{6}$ 
Leading Coefficient: 1

Pagne: 6

Lett: Rising

Right! Rising

Next, you should be able to find the x intercept(s) and the y intercept of a polynomial function.

## Zeros of Polynomial Functions (x-intercepts or Roots)

You will need to set the function equal to zero and then use the Zero Product Property to find the x-intercept(s). That means if ab = 0, then either a = 0 or b = 0. To find the y intercept of a function, you will find f(0).

a. 
$$f(x) = x^4 - x^2$$

$$\times^2 \left( \times^2 - 1 \right)$$

b. 
$$f(x) = -3x \left(x + \frac{1}{2}\right) (x - 4)^3$$
 [Factored Form]  
 $-3x = 0$   $\times +\frac{1}{2} = 0$   $\times -4 = 0$   
 $\times -6$   $\times -\frac{1}{2}$   $\times = 4$   
 $3 \times -\frac{1}{2} + \frac{1}{2} = 0$ 

$$\gamma - 1$$
 tercept:  
 $f(0) = -3(0)(0+\frac{1}{2})(0-4)^{3}$   
 $= -3(0)(1/2)(-4)^{3} = 0$ 

In some problems, one or more of the factors will appear more than once when the function is factored. The power of a factor is called its multiplicity. So given  $P(x) = x^3(x-3)^2(x+2)^1$ , then the multiplicity of the first factor is 3, the multiplicity of the second factor is and the multiplicity of  $P(0) = 0^{3}(0-3)^{2}(0+2)=0 \qquad X=0 \qquad X-3=0 \qquad X+2=0$   $M: 3 \qquad X=3 \qquad X=-2$ havior at Each x-intercept  $M: 1 \qquad V=0$ the third factor is 1.

Description of the Behavior at Each x-intercept

1. Even Multiplicity: The graph touches the x-axis, but does not cross it. It looks like a parabola Tangent to the x-axis there.

2. Multiplicity of 1: The graph crosses the *x*-axis. It looks like a line there.

3. Odd Multiplicity greater than or equal 3: The graph crosses the x-axis. It looks like a cubic there.

You can use all of this information to generate the graph of a polynomial function.

- degree of the function
- · end behavior of the function
- $\cdot x$  and y intercepts (and multiplicities)
- behavior of the function through each of the x intercepts (zeros) of the function

(the best we can draw is a rough sketch of a polynomial)