

# Answers to Odd-Numbered Problems

## CHAPTER 1

### Exercises 1.2

1. ordinary, first order
3. partial, second order
5. ordinary, third order
7. ordinary, second order
9. Both  $y$  and  $z$  are solutions.
11. Both  $y$  and  $z$  are solutions.
13. Both  $u_1$  and  $u_2$  are solutions.
15.  $u_1$  is a solution;  $u_2$  is not a solution.
17.  $r = -4, 2$ ;  $y_1(x) = e^{-4x}$  and  $y_2(x) = e^{2x}$  are solutions.
19.  $r = -3, 5$ ;  $y_1(x) = e^{-3x}$  and  $y_2(x) = e^{5x}$  are solutions.
21. No real values of  $r$ ;  $r = 1 \pm 2i$  are complex values.
23.  $r = -3, -2, 2, 3$ ;  $y_1(x) = e^{-3x}$ ,  $y_2(x) = e^{-2x}$ ,  $y_3 = e^{2x}$ ,  $y_4 = e^{3x}$  are solutions.
25.  $r = 1, 1, 2$ ;  $y_1(x) = e^x$ ,  $y_2 = xe^x$ ,  $y_3 = e^{2x}$  are solutions.
27.  $r = 2, 3$ ;  $y_1 = x^2$ ,  $y_2 = x^3$  are solutions.
29.  $r = 3, 3$ ;  $y_1 = x^3$ ,  $y_2 = x^3 \ln x$  are solutions.
31.  $r = -2, 0, 4$ ;  $y_1 = x^{-2}$ ,  $y_2 = x^0 = 1$ ,  $y_3 = x^4$  are solutions.
33.  $A = 3/2$ ;  $z = \frac{3}{2}e^{2x}$  is a solution.
35.  $A = 1/5$ ;  $z = \frac{1}{5}e^{-x}$  is a solution.

### Exercises 1.3

1.  $y = x^2 + x \ln x - x + C$ .
3.  $y = x^3 - \frac{1}{4} \cos 2x + C_1 x + C_2$ .
5.  $y = \frac{C}{x}$ .
9.  $y' = \frac{3y - 3}{x}$ .
11.  $y' = \frac{2y^3 - 6}{3xy^2}$ .

13.  $y' = \frac{3y^4 + 4x}{4xy^3}$ .
15.  $y' = y + \cos x - \sin x$ .
17.  $y'' + y' - 2y = 0$ .
19.  $y'' - 4y' + 4y = 0$ .
21.  $y'' + \frac{1}{x}y' - \frac{1}{x^2}y = 0$ .
23.  $y'' + 9y = 0$ .
25.  $x^3y''' - 3x^2y'' + 6xy' - 6y = 0$ .

#### Exercises 1.4

1. (b)  $y = 2e^{5x}$
3. (b)  $y = \frac{e}{e - 2e^x}$
5. (b)  $y = -\cos 2x - \frac{1}{2}\sin 2x$ .
7. (b)  $y = -\frac{17}{4} + 9x^{1/2}$   
 (c)  $y'$  is not defined at  $x = 0$ ; the initial-value problem does not have a solution.

### CHAPTER 2

#### Exercises 2.2

1.  $y = -\frac{1}{2} + Ce^{2x}$ .
3.  $y = 1 + Ce^{-x^2}$ .
5.  $y = e^{-x} + Ce^x$ .
7.  $y = x^{-2}\sin x + Cx^{-2}$ .
9.  $y = \frac{2}{9}(x+1)^{5/2} + C(x+1)^{-2}$ .
11.  $y = \sin x \cos x + C \cos x = \frac{1}{2}\sin 2x + C \cos x$ .
13.  $y = e^x + \frac{C}{x}$ .
15.  $y = x(\ln x)^2 + Cx$ .
17.  $y = 1 + Ce^{-e^x}$ .
19.  $y = x - 1 + 2e^{-x}$ .
21.  $y = \frac{\ln(1+e^x)}{e^x} + (e - \ln 2)e^{-x}$ .
23.  $y = \frac{5 - \cos 2x}{2 \sin x}$ .
25. (a)  $y = \frac{c}{b} + Ce^{-bx/a}$       (c)  $y = \frac{c}{b} + \left(\alpha - \frac{c}{b}e^{-bx/a}\right)$

### Exercises 2.3

1.  $y = \left(\frac{x^2}{4} + C\right)^2$ .
3.  $\tan^{-1} y = x^3 + C$  or  $y = \tan(x^3 + C)$ .
5.  $\cot y = \ln \sqrt{\frac{1-x}{1+x}} + C$ .
7.  $e^{-y} = e^x - xe^x + C$ .
9.  $\sqrt{y^2 - 1} = \frac{x}{1 + Cx}$ .
11.  $y^2 = C(\ln x)^2 - 1$ .
13.  $\ln |y| = -\ln |x| - \frac{1}{x} - 1$ .
15.  $y = xe^{x^2-1}$ .
17.  $y + \ln |y| = \frac{1}{3}x^3 - x - 5$ .
19.  $y = \frac{x + C}{1 - Cx}$

### Exercises 2.4

1.  $y = \frac{2}{Cx - 3x^3}$ .
3.  $y = (Ce^{2x} - e^x)^2$ .
5.  $y = \frac{1}{\sqrt[3]{Cx^3 - 2x^3 \ln x}}$ .
7.  $y^2 = Cx + x^2$ .
9.  $x \ln x + \frac{x+y}{e^{y/x}} = Cx$ .
11.  $\csc(y/x) - \cot(y/x) = Cx$ .
13.  $y^2 = \frac{C}{1+x^2} - 1$ .
15.  $y = \frac{\ln |\sec x + \tan x| + C}{x}$ .
17.  $y + \ln |1-y| = C - x - \ln |1+x|$ .
19.  $y = -x \ln(C - \ln x)$ .
21.  $y = C(3x^2 + 1)^{1/3} - 3$ .
23.  $y = \frac{1}{Cx + \ln x + 1}$ .
25.  $2y^3 = x^3 - Cx$ .
27. (a)  $u = \sin y$       (b)  $\sin y = e^{-x^2}(4x + C)$ .

### EXERCISES 2.5

#### Exercises 2.5.1

1.  $x^2 + 3y^2 = C$ .
3.  $\frac{x^2}{2} + y^2 - 4y = C$ .

5.  $y^2 = \ln(\sin^2 x) + C$ .
7.  $y = -\frac{1}{2}x^2 + C$ .
9.  $x^2 + \frac{1}{2}y^2 = C$ ; ellipses, center at the origin, major axis horizontal.
11.  $x^2 + y^2 - Cy = 0$ .

### Exercises 2.5.2

1. (a)  $A(t) = 50 \left(\frac{9}{10}\right)^{t/2} \approx 50e^{-0.05268t}$ . (b)  $A(4) = 50 \left(\frac{9}{10}\right)^2 = 40.5$  grams.  
 (c)  $T \approx 13.16$  hours.
3.  $t = \frac{2 \ln 10}{\ln 2} \approx 6.64$  hours.
5. (a)  $P(t) \approx 0.25e^{0.0421t}$  (b)  $\approx 1.6573$  square centimeters (c)  $\approx 16.464$  hours
7. (a)  $P(t) \approx 4.5e^{0.01438t}$ . (b) 48.19 years (c)  $\approx 6.93$  billion.

### Exercises 2.5.3

1. (a)  $40.1^\circ$ . (b) 1.62 minutes.
3. (a)  $u(t) = 150 - 100e^{\frac{t}{10} \ln(3/4)} = 150 - 100 \left(\frac{3}{4}\right)^{t/10}$   
 (b)  $t = \frac{10 \ln(1/2)}{\ln(3/4)} \approx 24.09$  minutes  
 (c) The temperature will never reach  $200^\circ$ ;  $\lim_{t \rightarrow \infty} u(t) = 150$
5. (a) Approximately 12.12 (b) Approximately 12:48

### Exercises 2.5.4

1. (a)  $v = \left(v_0 + \frac{g}{r}\right) e^{-rt} - \frac{g}{r}$  (b)  $\lim_{t \rightarrow \infty} v = -\frac{g}{r}$ .  
 (c)  $y = y_0 + \frac{1}{r} \left(v_0 + \frac{g}{r}\right) (1 - e^{-rt}) - \frac{g}{r}t$
3.  $k \approx 17.8$

### Exercises 2.5.5

1. (a)  $A(t) = 10,000 (1 - e^{-t/200})$  (b)  $t = 200 \ln 5 \approx 322$  minutes
3. (a)  $A(t) = \frac{9}{2} (1 - e^{-t/150})$  (b)  $t = 150 \ln 3 \approx 165$  minutes
5. (a)  $A(t) = \frac{3}{20} t(100 - t)$  (b)  $\max = A(50) = 375$

### Exercises 2.5.6

1. (a) 3259 people. (b)  $\approx 6.89$  days.
3. (a)  $\frac{d^2y}{dt^2} = k \frac{dy}{dt} (M - 2y)$ ;  $\frac{d^2y}{dt^2} > 0$  for  $0 < y < M/2$ ,  $\frac{d^2y}{dt^2} < 0$  for  $y > M/2$ .  
 $dy/dt$  has a maximum when  $y = M/2$
5.  $k \approx 0.0006$

**Exercises 2.5.7**

1. (a)  $\frac{dP}{dt} = kt(1000 - R) - 950e^{-kt^2/2}$       (b)  $P(t) = 1000 - Ce^{-kt^2/2}$       (c)  $P(t) = 1000 -$


3.  $A(t) = 1000 \left(\frac{4}{5}\right)^{t^2/100}$

5.  $A(t) = \frac{140,000}{140 + 3t}$

7.  $a = \frac{\ln 2}{24}$ ,  $b = \frac{\ln 2}{6}$


**Exercises 2.6**

1. (a) and (b)




(c)  $y = e^x$

3. (a) and (b)



(c)  $y = e^{x^2}$

5. Initial conditions:  $y(0) = 0$ ,  $u(0) = 1$ ,  $v(0) = 2$ .



## CHAPTER 3

### Exercises 3.2

1. Yes
3. Yes
5. Yes
7. (a)  $r = -1, r = 4$ .  
(b) Fundamental set:  $y_1(x) = x^{-1}, y_2(x) = x^4$ ; general solution:  $y = C_1x^{-1} + C_2x^4$ .  
(c)  $y = \frac{9}{5}x^{-1} + \frac{1}{5}x^4$ .  
(d) The trivial solution:  $y \equiv 0$ .
9.  $y'' - 2y' - 3y = 0$ .
11.  $y'' = 0$ .
13.  $x^2y'' - 2xy' + 2y = 0$ .
15.  $W[y_1, y_2](x) = e^{-\int_a^x p(t) dt} \neq 0$  for all  $x$ .
17.  $\{y_1(x) = x, y_2(x) = x^2\}$ .
19.  $\{y_1(x) = e^{x^2}, y_2(x) = e^{-x^2}\}$ .
21.  $\alpha\delta - \beta\gamma \neq 0$ .
23.  $W[y_1 + y_2, y_1 - y_2] = -2W[y_1, y_2]$ .
25. Set  $u(x) = \frac{y_2(x)}{y_1(x)}$ . Then

$$u'(x) = \frac{y_1y_2' - y_2y_1'}{y_1^2} = \frac{W[y_1, y_2]}{y_1^2} \equiv 0.$$

Therefore,  $u \equiv \lambda$  constant, which implies that  $y_2 = \lambda y_1$ .

### Exercises 3.3

1.  $y = C_1e^{2x} + C_2e^{-4x}$ .
3.  $y = C_1e^{5x} + C_2xe^{5x}$ .
5.  $y = e^{-2x}[C_1 \cos 3x + C_2 \sin 3x]$ .
7.  $y = C_1 + C_2e^{-2x}$ .
9.  $y = C_1e^{2\sqrt{3}x} + C_2e^{-2\sqrt{3}x}$ .
11.  $y = e^x[C_1 \cos x + C_2 \sin x]$ .
13.  $y = C_1e^{6x} + C_2e^{-5x}$ .
15.  $y = e^{-x/2}[C_1 \cos x/2 + C_2 \sin x/2]$ .
17.  $y = C_1e^{4x} + C_2xe^{4x}$ .
19.  $y = 2e^{2x} - e^{3x}$ .
21.  $y = -3e^{-x} - 2xe^{-x}$ .
23.  $y = -e^x \cos x$ .
25.  $y'' + 3y' - 10y = 0$ .

27.  $y'' + 4y = 0$ .
29.  $y'' - \frac{5}{2}y' + y = 0$ .
31.  $y'' + 2y' + 10y = 0$ .
33.  $y'' + 16y = 0$ .
35.  $y = (1 + \beta)e^{x/2} + (1 - \beta)e^{-x/2}$ ;  $\beta = -1$ .
37. (a)  $a^2 - 4b > 0$  (b)  $a^2 - 4b = 0$  (c)  $a^2 - 4b < 0$
39. (a)  $y'' + by = 0$ ,  $b > 0$ ; general solution:  $y = C_1 \cos x\sqrt{b} + C_2 \sin x\sqrt{b}$ , all solutions are bounded.

(b)  $y'' + ay' = 0$ ; general solution:  $y = C_1 + C_2e^{-ax}$  and  $\lim_{x \rightarrow \infty} y = C_1$ .

The solution that satisfies the initial conditions is:  $y = \left(\alpha + \frac{\beta}{a}\right) - \frac{\beta}{a}e^{-ax}$ ;  $k = \alpha + \frac{\beta}{a}$ .

41.  $r_1, r_2 = \frac{-a \pm \sqrt{a^2 - 4b}}{2} = \frac{-a}{2} \pm \frac{\sqrt{a^2 - 4b}}{2} = \alpha \pm \beta$ .

General solution:

$$\begin{aligned} y &= C_1 e^{(\alpha+\beta)x} + C_2 e^{(\alpha-\beta)x} = C_1 e^{\alpha x} e^{\beta x} + C_2 e^{\alpha x} e^{-\beta x} \\ &= e^{\alpha x} \left[ (C_1 + C_2) \frac{e^{\beta x} + e^{-\beta x}}{2} + (C_1 - C_2) \frac{e^{\beta x} - e^{-\beta x}}{2} \right] \\ &= e^{\alpha x} (K_1 \cosh \beta x + K_2 \sinh \beta x). \end{aligned}$$

43.  $y = C_1 x^{-2} + C_2 x^4$ .
45.  $y = C_1 x^2 + C_2 x^2 \ln x$ .

### Exercises 3.4

1.  $z(x) = x^2 \ln x + \frac{1}{2}$ ;  $y = C_1 x^2 + C_2 x^{-1} + x^2 \ln x + \frac{1}{2}$ .
3.  $z(x) = -x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2$ ;  $y = C_1 x + C_2 x^2 - x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2$ .
5.  $z(x) = -(1 + x^2)$ ;  $y = C_1 x + C_2 e^x - (1 + x^2)$ .
7.  $y = C_1 e^{-x} + C_2 e^{2x} - \frac{2}{3} x e^{-x}$ .
9.  $y = C_1 \cos 2x + C_2 \sin 2x + \frac{1}{4} \cos 2x \ln(\cos 2x) + \frac{1}{2} x \sin 2x$ .
11.  $y = C_1 e^x + C_2 x e^x - e^x \cos x$ .
13.  $y = C_1 e^{-2x} + C_2 x e^{-2x} - e^{-2x} \ln x$ .
15.  $y = C_1 \cos 3x + C_2 \sin 3x + \sin 3x \ln(\sec 3x + \tan 3x) - 1$ .
17.  $y = C_1 x + C_2 x^{-1} + x \ln x$ .
19.  $y = C_1 x + C_2 x \ln x + x^2$ .

### Exercises 3.5

1.  $y = C_1 e^{-x} + C_2 e^{3x} - e^{2x}$ .
3.  $y = C_1 e^{-3x} + C_2 x e^{-3x} + \frac{1}{4} e^{3x}$ .
5.  $y = C_1 e^{-2x} + C_2 - \frac{1}{2} \cos 2x - \frac{1}{2} \sin 2x$ .
7.  $y = C_1 e^{-x/2} + C_2 e^{-x} + x^2 - 6x + 14 - \frac{9}{10} \cos x - \frac{3}{10} \sin x$ .

9.  $y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{1}{2}x + \frac{1}{4}$ .
11.  $y = C_1 e^{-2x} + C_2 e^{-4x} + \frac{3}{2}x e^{-2x}$ .
13.  $y = C_1 \cos 3x + C_2 \sin 3x + \frac{2}{3} + \frac{1}{162}(9x^2 - 6x + 1)e^{3x}$ .
15.  $y = e^x(C_1 \cos 2x + C_2 \sin 2x) + \frac{1}{10}e^{-x} \cos 2x + \frac{1}{20}e^{-x} \sin 2x$ .
17.  $y = e^x - \frac{1}{2}e^{-2x} - x - \frac{1}{2}$ .
19.  $y = \frac{13}{15}e^{-x} + \frac{1}{12}e^{2x} + \frac{1}{20} \cos 2x - \frac{3}{20} \sin 2x$ .
21.  $z = A + (Bx^2 + Cx)e^{-x} + D \cos 3x + E \sin 3x$ .
23.  $z = Ax^2 + Bx + C + Dx \cos x + Ex \sin x$ .
25.  $z = (Ax^3 + Bx^2)e^{2x} + Cx^2 + Dx + E + (Fx + G) \cos 2x + (Hx + I) \sin 2x$ .
27.  $z = Ae^{-x} + Bxe^{-x} \cos x + Cxe^{-x} \sin x + D$ .
29.  $y = C_1 e^{2x} + C_2 x e^{2x} + \frac{8}{25} \cos x + \frac{6}{25} \sin x + 3x e^{2x} \ln x$ .
31.  $y = C_1 \cos 3x + C_2 \sin 3x + \frac{3}{8} \cos x - \sin 3x \ln(\sec 3x + \tan 3x) + 1$ .
33.  $y_1 - y_2$  is a solution of the reduced equation  $y'' + ay' + by = 0$  with  $a, b > 0$ . As shown in Exercises 3.3, Problem 38,  $y_1 - y_2 \rightarrow 0$  as  $x \rightarrow \infty$ . If  $a = 0, b > 0$ , then all solutions of the reduced equation are bounded (Problem 39 (a), Exercises 3.3).

### Exercises 3.6

1. The equation of motion is  $y(t) = \sin(8t + \frac{1}{2}\pi)$ . The amplitude is 1 and the frequency is  $8/2\pi = 4/\pi$ .
3.  $\pm 2\pi A/T$ .
5.  $y = C_1 \cos \omega t + C_2 \sin \omega t =$

$$\sqrt{C_1^2 + C_2^2} \left( \frac{C_1}{\sqrt{C_1^2 + C_2^2}} \cos \omega t + \frac{C_2}{\sqrt{C_1^2 + C_2^2}} \sin \omega t \right) \quad (*)$$

Let  $A = \sqrt{C_1^2 + C_2^2}$ . Since  $\left(\frac{C_1}{\sqrt{C_1^2 + C_2^2}}\right)^2 + \left(\frac{C_2}{\sqrt{C_1^2 + C_2^2}}\right)^2 = 1$  there is an angle  $\phi_0$  such that

$$\sin \phi_0 = \frac{C_1}{\sqrt{C_1^2 + C_2^2}} \quad \text{and} \quad \cos \phi_0 = \frac{C_2}{\sqrt{C_1^2 + C_2^2}}$$

(a) Substituting into (\*), we get

$$y = C_1 \cos \omega t + C_2 \sin \omega t = A(\cos \omega t \sin \phi_0 + \sin \omega t \cos \phi_0) = A \sin(\omega t + \phi_0)$$

(b)  $A \sin(\omega t + \phi_0) = A \cos(\omega t + \phi_0 - \frac{\pi}{2}) = A \cos(\omega t + \psi_0)$  where  $\psi_0 = \phi_0 - \frac{1}{2}\pi$ .

7. Assume that  $r_1 > r_2$ . If  $C_1 = 0$  or  $C_2 = 0$ , then  $y = C_1 e^{r_1 t} + C_2 e^{r_2 t}$  can never be zero. If both  $C_1$  and  $C_2$  are nonzero, then  $C_1 e^{r_1 t} + C_2 e^{r_2 t} = 0$  implies  $e^{(r_1 - r_2)t} = -\frac{C_2}{C_1}$ . Since  $e^{(r_1 - r_2)t}$  is an increasing function ( $r_1 > r_2$ ), it can take the value  $\frac{C_2}{C_1}$  at most once. By the same reasoning,  $y'(t) = C_1 r_1 e^{r_1 t} + C_2 r_2 e^{r_2 t}$  can be zero at most once. Therefore the motion can change direction at most once.



9. Suppose  $\gamma \neq \omega$ . Set  $z = A \cos \gamma t + B \sin \gamma t$  and use undetermined coefficients. The result is

$$z = \frac{F_0/m}{\omega^2 - \gamma^2} \cos \gamma t.$$

11. Suppose  $\gamma = \omega$ . Then a particular solution  $z$  has the form  $z = At \cos \omega t + Bt \sin \omega t$ . Substituting  $z$  into the equation, we get

$$A = 0, B = \frac{F_0}{2\omega m}, \quad \text{and so} \quad z = \frac{F_0}{2\omega m} t \sin \omega t.$$

### Exercises 3.7

1.  $y = C_1 e^x + C_2 e^{2x} + C_3 e^{3x}$ .
3.  $y = C_1 e^{2x} + C_2 e^{-2x} + e^x [C_3 \cos 2x + C_4 \sin 2x]$ .
5.  $y = C_1 \cos x + C_2 \sin x + e^{2x} [C_3 \cos 3x + C_4 \sin 3x]$ .
7.  $y = C_1 + C_2 x + C_3 e^x + C_4 e^{-x} + C_5 \cos x + C_6 \sin x$ .
9.  $y = 2x - 1$ .
11.  $y = \frac{1}{5} e^x - \frac{1}{5} \cos 3x - \frac{1}{15} \sin x$
13.  $y^{(4)} + 4y''' + 7y'' + 42y' + 90y = 0$ .
15.  $y^{(5)} - 2y^{(4)} - 2y''' - 2y'' - 3y' = 0$ .
17.  $y^{(5)} - 2y^{(4)} + y''' - 2y'' = 0$ .
19.  $y^{(4)} - y'' = 0$ .
21.  $y = C_1 e^{-x} + C_2 \cos x + C_3 \sin x + \frac{1}{4} e^x + 4$ .
23.  $y = C_1 \cos x + C_2 \sin x + C_3 x \cos x + C_4 x \sin x + 6 + \frac{1}{9} \cos 2x$ .
25.  $y = -\frac{1}{24} e^{2x} + \frac{1}{12} x e^{2x} + \frac{\sqrt{3}}{72} e^{-x} \sin \sqrt{3}x + \frac{1}{24} e^{-x} \cos \sqrt{3}x$

## CHAPTER 4

### Exercises 4.2

1.  $\frac{1}{s^2}$ .
3.  $\frac{1}{s^2 + 1}$ .
5.  $\frac{1}{2(s-1)} - \frac{1}{2(s+1)}$ .
7.  $\frac{s-a}{(s-a)^2 + b^2}$ .

### Exercises 4.3

1.  $\frac{3}{s} - \frac{2}{s^2} + \frac{2}{s^3}$ .
3.  $\frac{3}{s} + \frac{4}{s-3} - \frac{2s}{s^2+4}$ .
5.  $\frac{10}{s^3} - \frac{4}{(s+3)^2+4}$ .

7.  $\frac{2s}{(s^2+1)^2} + \frac{2(s^2-4)}{(s^2+4)^2}$ .
9.  $\sinh \beta x = \frac{e^{\beta x} - e^{-\beta x}}{2}$ ;  $\frac{1}{2} \left[ \frac{1}{s-\beta} - \frac{1}{s+\beta} \right] = \frac{\beta}{s^2 - \beta^2}$ .
11.  $\frac{1}{2} \left[ \frac{1}{s-3} + \frac{1}{s-2} - \frac{1}{s-1} + \frac{1}{s-4} \right]$ .
15.  $Y(s) = \frac{1}{s-2}$ .
17.  $Y(s) = \frac{2}{(s-2)(s+4)} - \frac{9}{(s^2+9)(s+4)} - \frac{3}{s+4}$ .
19.  $Y(s) = \frac{2}{(s+3)^2}$ .
21.  $Y(s) = \frac{3}{s(s-5)(s+3)} + \frac{4}{(s-5)(s+3)^2} + \frac{s-5}{(s-5)(s+3)}$ .
23. Set  $g(x) = \int_0^x f(t) dt$ . Then  $g'(x) = f(x)$  and  $g(0) = 0$ .

$$F(s) = \mathcal{L}[f(x)] = \mathcal{L}[g'(x)] = s\mathcal{L}[g(x)] - g(0) = s\mathcal{L}[g(x)].$$

$$\text{Therefore, } \mathcal{L}[g(x)] = \frac{1}{s}F(s).$$

#### Exercises 4.4

1.  $f(x) = 6e^{-7x}$ .
3.  $f(x) = 2 \cos 5x + \frac{1}{5} \sin 5x$ .
5.  $f(x) = e^{-4x} \cos x$ .
7.  $f(x) = e^{-2x} \cos 2x + e^{-2x} \sin 2x$ .
9.  $f(x) = 2xe^{-2x} - e^x \cos x - e^x \sin x$ .
11.  $f(x) = \frac{1}{2} e^{-x} - \frac{1}{2} \cos x + \frac{1}{2} \sin x$ .
13.  $f(x) = \frac{1}{4} - \frac{1}{4} \cos 2x$ .
15.  $f(x) = \frac{1}{2} - e^x + \frac{3}{2} e^{-2x}$ .
17.  $f(x) = e^{2x} - 4e^x + 2x + 3$ .
19.  $y = \frac{2}{3} e^{-2x} + \frac{1}{3} e^x$ .
21.  $y = \frac{3}{2} e^{-x} - \frac{1}{2} \cos x + \frac{1}{2} \sin x$ .
23.  $y = e^x \sin x$ .
25.  $y = \frac{3}{4} e^{-x} + \frac{1}{4} e^x + \frac{1}{2} x e^x$ .
27.  $y = \frac{1}{4} e^x + x e^{-x} + x - 2$ .
29.  $y = e^{-2x} + e^x$ .
31.  $y = -\frac{1}{5} e^{-2x} \cos 2x - \frac{1}{10} e^{-2x} \sin 2x + \frac{1}{5} e^{-x}$ .
33.  $\alpha = \frac{1}{4}$ .
35.  $\beta = -\frac{26}{5}$ .
37.  $y = \frac{7}{4} e^{2(x-1)} - 3e^{x-1} + \frac{1}{2} x + \frac{3}{4}$ .

**Exercises 4.5**

1.  $F(s) = 2e^{-5s} \frac{1}{s}$ .
3.  $F(s) = \frac{2}{s^2} - 2e^{-3s} \frac{1}{s^2} - 5e^{-3s} \frac{1}{s}$ .
5.  $f(x) = 0 + 5u(x-4)$ ;  $F(s) = 5e^{-4s} \frac{1}{s}$ .
7.  $f(x) = 0 + (x-2)u(x-2) + 2u(x-2)$ ;  $F(s) = e^{-2s} \frac{1}{s^2} + 2e^{-2s} \frac{1}{s}$ .
9.  $f(x) = 8 + 2(x-5)u(x-5) + 2u(x-5)$ ;  $F(s) = \frac{8}{s} + 2e^{-5s} \frac{1}{s^2} + 2e^{-5s} \frac{1}{s}$ .
11.  $f(x) = x^2 - (x-3)^2 u(x-3) - 3(x-3)u(x-3)$ ;  $F(s) = \frac{2}{s^3} - e^{-3s} \frac{2}{s^3} - 3e^{-3s} \frac{1}{s^2}$ .
13.  $f(x) = x - 1 - (x-2)u(x-2) - u(x-2) + e^{-2} e^{-(x-2)} u(x-2)$ ;  
 $F(s) = \frac{1}{s^2} - \frac{1}{s} - e^{-2s} \frac{1}{s^2} - e^{-2s} \frac{1}{s} + e^{-2} e^{-2s} \frac{1}{s+1}$ .
15.  $f(x) = \sin 2x - \sin 2(x-\pi)u(x-\pi) + (x-\pi)u(x-\pi) + \pi u(x-\pi)$ ;  
 $F(s) = \frac{2}{s^2+4} - e^{-\pi s} \frac{2}{s^2+4} + e^{-\pi s} \frac{1}{s^2} + \pi e^{-\pi s} \frac{1}{s}$ .
17.  $f(x) = x - (x-2)u(x-2) - 2u(x-2) + (x-4)^2 u(x-4)$ ;  
 $F(s) = \frac{1}{s^2} - e^{-2s} \frac{1}{s^2} - 2e^{-2s} \frac{1}{s} + e^{-4s} \frac{2}{s^3}$ .
19.  $f(x) = 1 - u(x-2) + (x-2)u(x-2) - (x-4)u(x-4) - 2u(x-4) + e^{-(x-4)} u(x-4)$ ;  
 $F(s) = \frac{1}{s} - e^{-2s} \frac{1}{s} + e^{-2s} \frac{1}{s^2} - e^{-4s} \frac{1}{s^2} - 2e^{-4s} \frac{1}{s} + e^{-4s} \frac{1}{s+1}$ .

**Exercises 4.6**

1.  $f(x) = 2 + 2u(x-3) = \begin{cases} 2, & 0 \leq x < 3 \\ 4, & x \geq 3 \end{cases}$ .
3.  $f(x) = \sin x - \sin x u(x-\pi) = \begin{cases} \sin x, & 0 \leq x < \pi \\ 0, & x \geq \pi. \end{cases}$
5.  $f(x) = \cos x - \cos x u(x-\pi) + \sin x u(x-\pi) = \begin{cases} \cos x, & 0 \leq x < \pi \\ \sin x, & x \geq \pi. \end{cases}$
7.  $f(x) = \cos \pi x - \sin \pi x u(x-2) = \begin{cases} \cos \pi x, & 0 \leq x < 2 \\ \cos \pi x - \sin \pi x, & x \geq 2. \end{cases}$
9.  $f(x) = 3e^{3(x-3)} u(x-3) - 2e^{2(x-3)} u(x-3) = \begin{cases} 0, & 0 \leq x < 3 \\ 3e^{3(x-3)} - 2e^{2(x-3)}, & x \geq 3. \end{cases}$
11.  $f(x) = 2 + e^{(x-1)} u(x-1) - e^2 e^{(x-2)} u(x-2) = \begin{cases} 2, & 0 \leq x < 1 \\ 2 + e^{(x-1)}, & 1 \leq x < 2 \\ 2 + e^{(x-1)} - e^x, & x \geq 2. \end{cases}$
13.  $f(x) = \cos 2x - 1 + u(x-2) - \cos 2(x-2) u(x-2) = \begin{cases} \cos 2x - 1, & 0 \leq x < 2 \\ \cos 2x - \cos 2(x-2), & x \geq 2. \end{cases}$
15.  $f(x) = 2e^\pi e^{-2x} \cos 3x u(x-\pi/2) - e^\pi e^{-2x} \sin 3x u(x-\pi/2)$

$$= \begin{cases} 0, & 0 \leq x < \pi/2 \\ 2e^\pi e^{-2x} \cos 3x - e^\pi e^{-2x} \sin 3x & x \geq \pi/2. \end{cases}$$

### Exercises 4.7

1.  $y = -\frac{1}{2} + \frac{5}{2}e^{2x} + u(x-1) \left[-\frac{1}{2} + \frac{1}{2}e^{2(x-1)}\right].$   

$$= \begin{cases} -\frac{1}{2} + \frac{5}{2}e^{2x}, & 0 \leq x < 1 \\ -1 + \frac{5}{2}e^{2x} + \frac{1}{2}e^{2(x-1)}, & x \geq 1 \end{cases}$$
3.  $y = 1 - \cos x + \sin x - u(x-1)[\cos(x-1) - 1].$   

$$= \begin{cases} 1 - \cos x + \sin x, & 0 \leq x < \pi \\ 2 \cos x, & x \geq \pi \end{cases}$$
5.  $y = 1 - e^{-x} - xe^{-x} + u(x-2) \left[x - 4 + xe^{-(x-2)}\right].$   

$$= \begin{cases} 1 - e^{-x} - xe^{-x}, & 0 \leq x < 2 \\ -3 - e^{-x} - xe^{-x} + x + xe^{-(x-2)}, & x \geq 2 \end{cases}$$
7.  $y = -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x + u(x-1) \left[\frac{1}{3} + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{x-1}\right].$   

$$= \begin{cases} -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x, & 0 \leq x < 1 \\ -\frac{1}{6}e^{3x} + \frac{1}{2}e^x + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{(x-1)}, & x \geq 1 \end{cases}$$
9.  $y = \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + u(x-1) \left[xe^{-(x-1)} - 1\right].$   

$$= \begin{cases} \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x}, & 0 \leq x < 1 \\ \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + xe^{-(x-1)} - 1, & x \geq 1 \end{cases}$$

## CHAPTER 5

### Exercises 5.2

1.  $x = 4, y = 1.$
3.  $x = 4 - 2a, y = a, a$  any real number.
5.  $x = -3, y = 1.$
7. No solution.
9.  $x = 2a - 3, y = a, a$  any real number.
11.  $x = \frac{3}{7}a + 1, y = \frac{5}{7}a - 1, z = a, a$  any real number; a line in 3-space.
13. No solution.
15.  $x = 2, y = 1, z = 1;$  a point in 3-space.

### Exercises 5.3

1. matrix of coefficients: 3; augmented matrix: 3;  $x = 5, y = 3, z = -1.$
3. matrix of coefficients: 2; augmented matrix: 2;  $x = 4 - 2a, y = a, z = -2, a$  an real number.

5. matrix of coefficients: 3; augmented matrix: 3;  $x_1 = -1$ ,  $x_2 = -1 - 2a$ ,  $x_3 = 3 + a$ ,  $x_4 = a$ ,  $a$  any real number.
7. matrix of coefficients: 3; augmented matrix: 3;  $x_1 = 8 + 2a - 3b$ ,  $x_2 = a$ ,  $x_3 = 3 - 1 - 2b$ ,  $x_4 = b$ ,  $x_5 = -3$ ,  $a$ ,  $b$  any real numbers.
9.  $x = 2$ ,  $y = 5$ .
11.  $x = -3 - a$ ,  $y = 2 + 2a$ ,  $z = a$ ,  $a$  any real number.
13.  $x = \frac{10}{7}$ ,  $y = \frac{2}{7}$ ,  $z = \frac{3}{2}$ .
15.  $x_1 = 11 - 2a + b$ ,  $x_2 = a$ ,  $x_3 = 3 - b$ ,  $x_4 = b$ ,  $a$ ,  $b$  any real numbers.
17.  $x_1 = -2$ ,  $x_2 = -5$ ,  $x_3 = -1$ ,  $x_4 = 5$ .
19.  $x_1 = 3 - 2a$ ,  $x_2 = a$ ,  $x_3 = 2$ ,  $x_4 = 1$ .
21. No solution.
23. (i)  $k \neq -3$ , 2 (ii)  $k = -3$  (iii)  $k = 2$ .
25. (i)  $a \neq -3$ , 3 (ii)  $a = -3$  (iii)  $a = 3$ .
27.  $a$  any real number;  $b \neq a$ .
29. (a) No (b) No (c) Yes

#### Exercises 5.4

1. Yes.
3. No. The leading 1 in the third row is not to the right of the leading 1 in the second row.
5. No. The leading 1 in the last column is not the only nonzero in its column.
7. Yes
9.  $x = 10$ ,  $y = -9$ ,  $z = -7$
11.  $x = -3 - a$ ,  $y = 2 + 2a$ ,  $z = a$ ,  $a$  any real number.
13.  $x_1 = 11 - 2a + b$ ,  $x_2 = a$ ,  $x_3 = 3 - b$ ,  $x_4 = b$ ,  $a$ ,  $b$  any real numbers.
15.  $x_1 = 7 - 2a - b$ ,  $x_2 = 1 + 3a - 4b$ ,  $x_3 = a$ ,  $x_4 = b$ ,  $a$ ,  $b$  any real numbers.
17.  $x = y = 0$ .
19.  $x = y = z = 0$ .
21.  $x_1 = 2a - b$ ,  $x_2 = -a + 4b$ ,  $x_3 = a$ ,  $x_4 = b$ ,  $a$ ,  $b$  any real numbers.
23.  $x_1 = x_2 = x_3 = x_4 = 0$ .
25. Consider the system

$$\begin{aligned}x + y &= 0 \\2x + 2y &= 0 \\3x + 3y &= 0\end{aligned}$$

This system has the solutions  $x = -a$ ,  $y = a$ ,  $a$  any real number.

27.  $a = 4$ .
29. (a) Solution set  $\mathcal{S}$ :  $x = 1 + a$ ,  $y = -1 - a$ ,  $z = a$ ,  $a$  any real number.

$$(b) \mathcal{S} : \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} + a \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}.$$

### Exercises 5.5

1. (a)  $\begin{pmatrix} 0 & 4 \\ 3 & 5 \\ 1 & -1 \end{pmatrix}$ . (c)  $\begin{pmatrix} 8 & -4 \\ 1 & 4 \end{pmatrix}$ . (e)  $\begin{pmatrix} 2 & 8 \\ 2 & 8 \\ 5 & -3 \end{pmatrix}$ .
3. (a)  $\begin{pmatrix} -4 & -3 \\ 28 & -6 \\ -20 & 24 \end{pmatrix}$ . (c) Not defined. (e)  $\begin{pmatrix} 1 & 3 \\ -3 & -12 \\ -41 & 21 \end{pmatrix}$ .
5. (a)  $c_{32} = 2$  (b)  $c_{13} = 34$  (c)  $d_{21} = 5$  (d)  $d_{22} = 1$ .
7. (a)  $d_{22} = 6$  (b)  $d_{12} = -4$  (c)  $d_{23} = -18$ .
11. (a)  $AB = \begin{pmatrix} 4 & 7 & 10 \\ 0 & -5 & -14 \end{pmatrix}$ ,  $BA$  not defined.
- (b)  $AC = \begin{pmatrix} 14 & 5 \\ -2 & -3 \end{pmatrix}$ ,  $CA = \begin{pmatrix} -1 & 14 \\ 5 & 12 \end{pmatrix}$ .
- (c)  $AD = DA = \begin{pmatrix} 4 & 4 \\ -2 & 2 \end{pmatrix}$ .
13.  $A(BD) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}$
- $(AB)D = \begin{pmatrix} 0 & -2 \\ 3 & 5 \end{pmatrix} \begin{pmatrix} -2 & 3 & -2 \\ 3 & 0 & -4 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}$ .
15. (a)  $3 \times 3$  (c) Does not exist (e)  $2 \times 3$ .

### Exercises 5.6

1.  $A^{-1} = \begin{pmatrix} 1/2 & 0 \\ -3/2 & 1 \end{pmatrix}$ .
3.  $A^{-1} = \begin{pmatrix} -2 & 1 \\ 3/2 & -1/2 \end{pmatrix}$ .
5.  $A^{-1} = \begin{pmatrix} -11 & 2 & 2 \\ -4 & 0 & 1 \\ 6 & -1 & -1 \end{pmatrix}$ .
7. No inverse.
9. No inverse.
11.  $\det A = \pm 1$ .
13.  $x = 5, y = 0$ .
15.  $x = \frac{9}{2}, y = -5$ .
17.  $x = \frac{7}{9}, y = \frac{1}{3}, z = -\frac{5}{9}$ .
19.  $-31$ .

21.  $-45$ .  
 23.  $30$ .  
 25.  $-21$ .  
 27.  $-18$ .  
 29.  $26$ .  
 31.  $x = 0, 1, -3$ .  
 33.  $y = -\frac{25}{37}$ .  
 35. Cramer's rule does not apply.  
 37.  $x = 0$ .  
 39.  $\lambda = -4, 7$ .

### Exercises 5.7

3. Dependent;  $(-4, 8, 9) = 2(1, -2, 3) + 3(-2, 4, 1)$ .  
 5. Dependent;  $(-2, 6, 3) = (1, -1, 3) + 2(0, 2, 3) - 3(1, -1, 2)$ .  
 7. Dependent;  $(7, -4, 1) = 3(1, -2, 1) + 2(2, 1, -1)$ .  
 9. Dependent;  $(4, -2, 0, 2) = 2(2, -1, 0, 1)$ .  
 11.  $b \neq -\frac{1}{3}$ .  
 13.  $b = 0, -7$ .  
 17. No; a linearly dependent set can have linearly independent subsets. For example,  $\{(1, -2, 3), (-2, 4, 1)\}$  is a linearly independent subset of  $\{(1, -2, 3), (-2, 4, 1)\}, (-4, 8, 9)$ .  
 19.  $W(x) = -a$ ; linearly independent.  
 21.  $W(x) = -2x^{-6}$ ; linearly independent.  
 23.  $W(x) = e^{2x}(x - 2)$ ; linearly independent.  
 25. (a) False (b) True (c) True.

### Exercises 5.8

1.  $2, \begin{pmatrix} 1 \\ 0 \end{pmatrix}; 3, \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ .  
 3.  $-1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}; 4, \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ .  
 5.  $1, 1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ .  
 7.  $2, 2, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .  
 9.  $2 + i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} + i \begin{pmatrix} 0 \\ -1 \end{pmatrix}; 2 - i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} - i \begin{pmatrix} 0 \\ -1 \end{pmatrix}$ .  
 11.  $8, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}; 1, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}; 2, \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ .

13.  $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}.$
15.  $1+i, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + i \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}; 1-i, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - i \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}; 0, \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}.$
17.  $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$
19.  $7, \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}; -2, \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix}; -2, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$
21.  $2, \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}; 2, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}; 6, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}; 4, \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}.$

## CHAPTER 6

### Exercises 6.1

1.  $\begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -3 & t \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 0 \\ \sin 2t \end{pmatrix}.$
3.  $\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ e^t \end{pmatrix}.$
5. 
$$\begin{aligned} x'_1 &= 2x_1 - x_2 + e^{2t} \\ x'_2 &= 3x_1 + 2e^{-t} \end{aligned}.$$
7. 
$$\begin{aligned} x'_1 &= 2x_1 + 3x_2 - x_3 + e^t \\ x'_2 &= -2x_1 + x_3 + 2e^{-t} \\ x'_3 &= 2x_1 + 3x_2 + e^{2t} \end{aligned}.$$
9.  $\begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} \sin t \\ -2 \cos t \end{pmatrix}.$
11.  $\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 2 & 1 & 3 \\ 1 & -3 & 0 \\ 2 & -1 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 3e^{2t} \\ -2 \cos t \\ t \end{pmatrix}.$

### Exercises 6.2

1. Independent
3. Independent
5. Dependent
7. Dependent
9. Dependent



11. (c)  $\mathbf{x}(t) = c_1\mathbf{u} + c_2\mathbf{v}$ , where  $c_1, c_2$  are arbitrary constants.

$$(d) \mathbf{x}(t) = -2 \begin{pmatrix} e^{2t} \\ e^{2t} \end{pmatrix} + \begin{pmatrix} 3e^{3t} \\ 2e^{3t} \end{pmatrix}.$$

$$13. (b) \mathbf{x}(t) = \begin{pmatrix} -4te^{-t} \\ 2 - e^{-t} \\ 2 \end{pmatrix}.$$

15. The system is equivalent to  $y''' + y'' - 4y' - 4y = 0$ . A fundamental set of solutions of this equation is  $\{y_1 = e^{-2t}, y_2 = e^{-t}, y_3 = e^{2t}\}$ . Therefore, linearly independent solutions of the given system are:

$$\mathbf{x}_1 = \begin{pmatrix} e^{-2t} \\ -2e^{-2t} \\ 4e^{-2t} \end{pmatrix} = e^{-2t} \begin{pmatrix} 1 \\ -2 \\ 4 \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} e^{-t} \\ -e^{-t} \\ e^{-t} \end{pmatrix} = e^{-t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} e^{2t} \\ 2e^{2t} \\ 4e^{2t} \end{pmatrix} = e^{2t} \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix}.$$

17. The system is equivalent to  $y'' + \frac{6}{t}y' + \frac{6}{t^2}y = 0$ . A fundamental set of solutions of this equation is  $\{y_1 = t^{-3}, y_2 = t^{-2}\}$ . Therefore, linearly independent solutions of the given system are:

$$\mathbf{x}_1 = \begin{pmatrix} t^{-3} \\ -3t^{-4} \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} t^{-2} \\ -2t^{-3} \end{pmatrix}.$$

### Exercises 6.3

$$1. \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-3t} \begin{pmatrix} 4 \\ -1 \end{pmatrix}.$$

$$3. \mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + C_2 e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}; \quad \mathbf{x}(t) = -3e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + 4e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

$$5. \mathbf{x}(t) = C_1 e^{5t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 2 \\ -1 \end{pmatrix}.$$

$$7. \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$$

$$9. \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}; \quad \mathbf{x}(t) = -e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + 2e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}.$$

$$11. \mathbf{x}(t) = C_1 e^{10t} \begin{pmatrix} 9 \\ 7 \\ 4 \end{pmatrix} + C_2 e^{6t} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + C_3 e^{5t} \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}.$$

$$13. \quad (a) \mathbf{x}' = \begin{pmatrix} 0 & 1 \\ -b & -a \end{pmatrix} \mathbf{x}; \quad (b) \lambda^2 + a\lambda + b = 0; \quad (c) \text{ They are the same.}$$

#### Exercises 6.4

$$1. \mathbf{x}(t) = C_1 \left[ \cos 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right] + C_2 \left[ \cos 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} \right].$$

$$3. \mathbf{x}(t) = C_1 e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + C_2 \left[ e^t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right].$$

$$5. \mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} + C_2 \left[ e^{-t} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} \right].$$

$$7. \mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + C_3 e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix};$$

$$\mathbf{x}(t) = e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$$

$$9. \mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} + C_2 \left[ \cos 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] + C_3 \left[ \cos 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} \right].$$

$$11. \mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + C_3 e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix};$$

$$\mathbf{x}(t) = \frac{7}{2} e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \frac{9}{2} e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \frac{3}{2} e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}$$

$$13. \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_3 \left[ e^t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$$

$$15. \mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \left[ \cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + C_3 e^t \left[ \cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right];$$

$$\mathbf{x}(t) = e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + e^t \left[ \cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + 3e^t \left[ \cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$$

17.  $\mathbf{x}(t) = C_1 e^{6t} \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} + C_2 e^{-2t} \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} + C_3 e^{-2t} \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix}.$