

# Math 2331 – Linear Algebra

## 3.1 Introduction to Determinants

**Jiwen He**

Department of Mathematics, University of Houston

`jiwenhe@math.uh.edu`  
`math.uh.edu/~jiwenhe/math2331`



## 3.1 Introduction to Determinants

- Determinant via Cofactor Expansion
  - Examples
  - Theorem
- Triangular Matrices
  - Examples
  - Theorem



# Introduction to Determinants

Notation:  $A_{ij}$  is the matrix obtained from matrix  $A$  by deleting the  $i$ th row and  $j$ th column of  $A$ .

## Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix} \quad A_{23} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$$

Recall that  $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$  and we let  $\det [a] = a$ .

For  $n \geq 2$ , the **determinant** of an  $n \times n$  matrix  $A = [a_{ij}]$  is

$$\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{1+n} a_{1n} \det A_{1n}$$

$$= \sum_{j=1}^n (-1)^{1+j} a_{1j} \det A_{1j}$$


# Determinants: Example

## Example

Compute the determinant of  $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$

## Solution

$$\det A = 1 \det \begin{bmatrix} -1 & 2 \\ 0 & 1 \end{bmatrix} - 2 \det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} + 0 \det \begin{bmatrix} 3 & -1 \\ 2 & 0 \end{bmatrix}$$

$$= \text{-----} = \text{-----}$$

Common notation:  $\det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} = \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix}$ .

So

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1 \begin{vmatrix} -1 & 2 \\ 0 & 1 \end{vmatrix} - 2 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix}$$



# Determinants: Cofactor Expansion

## Cofactor

The **(i, j)-cofactor** of  $A$  is the number  $C_{ij}$  where

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

## Example (Cofactor Expansion)

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1C_{11} + 2C_{12} + 0C_{13}$$

(cofactor expansion across row 1)



# Cofactor Expansion: Theorem

## Theorem (Cofactor Expansion)

The determinant of an  $n \times n$  matrix  $A$  can be computed by a cofactor expansion across any row or down any column:

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}$$

(expansion across row  $i$ )

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}$$

(expansion down column  $j$ )

Use a matrix of signs to determine  $(-1)^{i+j}$

$$\begin{bmatrix} + & - & + & \cdots \\ - & + & - & \cdots \\ + & - & + & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$



# Cofactor Expansion: Example

## Example

Compute the determinant of  $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$  using cofactor expansion down column 3.

## Solution

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 3 & -1 \end{vmatrix} = 1.$$



# Cofactor Expansion: Example

## Example

Compute the determinant of  $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{bmatrix}$

## Solution:

$$\begin{aligned} & \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \\ &= 1 \begin{vmatrix} 2 & 1 & 5 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} + 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 2 & 1 \end{vmatrix} \\ &= 1 \cdot 2 \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} = 14 \end{aligned}$$



# Triangular Matrices

*Method of cofactor expansion is not practical for large matrices - see Numerical Note on page 167.*

## Triangular Matrices

$$\begin{bmatrix} * & * & \cdots & * & * \\ 0 & * & \cdots & * & * \\ 0 & 0 & \ddots & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & * \end{bmatrix}$$

(upper triangular)

$$\begin{bmatrix} * & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 \\ * & * & \ddots & 0 & 0 \\ * & * & \cdots & * & 0 \\ * & * & \cdots & * & * \end{bmatrix}$$

(lower triangular)

## Theorem

*If  $A$  is a triangular matrix, then  $\det A$  is the product of the main diagonal entries of  $A$ .*



# Triangular Matrices: Example

## Example

$$\begin{vmatrix} 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & -3 & 5 \\ 0 & 0 & 0 & 4 \end{vmatrix} = \text{-----} = -24$$

