

Math 2331 – Linear Algebra

4.2 Null Spaces, Column Spaces, & Linear Transformations

Jiwen He

Department of Mathematics, University of Houston

`jiwenhe@math.uh.edu`
`math.uh.edu/~jiwenhe/math2331`



4.2 Null Spaces, Column Spaces, & Linear Transformations

- Null Spaces
 - Definition
 - Theorem
 - Examples
- Column Spaces
 - Definition
 - Theorem
 - Examples
- The Contrast Between Nul A and Col A
- Null Spaces & Column Spaces: Review
- Null Spaces & Column Spaces: Examples
- Kernel and Range of a Linear Transformation



Null Space

Null Space

The **null space** of an $m \times n$ matrix A , written as $\text{Nul } A$, is the set of all solutions to the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

$$\text{Nul } A = \{\mathbf{x} : \mathbf{x} \text{ is in } \mathbf{R}^n \text{ and } A\mathbf{x} = \mathbf{0}\} \quad (\text{set notation})$$

Theorem (2)

The null space of an $m \times n$ matrix A is a subspace of \mathbf{R}^n . Equivalently, the set of all solutions to a system $A\mathbf{x} = \mathbf{0}$ of m homogeneous linear equations in n unknowns is a subspace of \mathbf{R}^n .

Proof: $\text{Nul } A$ is a subset of \mathbf{R}^n since A has n columns. Must verify properties a, b and c of the definition of a subspace.

Property (a) Show that $\mathbf{0}$ is in $\text{Nul } A$. Since _____, $\mathbf{0}$ is in

Null Space (cont.)

Property (b) If \mathbf{u} and \mathbf{v} are in $\text{Nul } A$, show that $\mathbf{u} + \mathbf{v}$ is in $\text{Nul } A$.
 Since \mathbf{u} and \mathbf{v} are in $\text{Nul } A$,

_____ and _____.

Therefore

$$A(\mathbf{u} + \mathbf{v}) = \text{_____} + \text{_____} = \text{_____} + \text{_____} = \text{_____}.$$

Property (c) If \mathbf{u} is in $\text{Nul } A$ and c is a scalar, show that $c\mathbf{u}$ is in $\text{Nul } A$:

$$A(c\mathbf{u}) = \text{---}A(\mathbf{u}) = c\mathbf{0} = \mathbf{0}.$$

Since properties a, b and c hold, A is a subspace of \mathbf{R}^n .

Solving $A\mathbf{x} = \mathbf{0}$ yields an *explicit description* of $\text{Nul } A$.



Null Space: Example

Example

Find an explicit description of $\text{Nul } A$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}$$

Solution: Row reduce augmented matrix corresponding to $A\mathbf{x} = \mathbf{0}$:

$$\left[\begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \sim \dots \sim \left[\begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right]$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 13x_4 - 33x_5 \\ x_2 \\ 6x_4 + 15x_5 \\ x_4 \\ x_5 \end{bmatrix}$$



Null Space: Example (cont.)

$$= x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}$$

Then

$$\text{Nul } A = \text{span}\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$$



Null Space: Observations

Observations:

- Spanning set of $\text{Nul } A$, found using the method in the last example, is automatically linearly independent:

$$c_1 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

\implies

$$c_1 = \text{-----} \quad c_2 = \text{-----} \quad c_3 = \text{-----}$$

- If $\text{Nul } A \neq \{\mathbf{0}\}$, the the number of vectors in the spanning set for $\text{Nul } A$ equals the number of free variables in $A\mathbf{x} = \mathbf{0}$.



Column Space

Column Space

The **column space** of an $m \times n$ matrix A ($\text{Col } A$) is the set of all linear combinations of the columns of A .

If $A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_n]$, then

$$\text{Col } A = \text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$$

Theorem (3)

The column space of an $m \times n$ matrix A is a subspace of \mathbf{R}^m .

Why? (Theorem 1, page 194)

Recall that if $A\mathbf{x} = \mathbf{b}$, then \mathbf{b} is a linear combination of the columns of A . Therefore

$$\text{Col } A = \{\mathbf{b} : \mathbf{b} = A\mathbf{x} \text{ for some } \mathbf{x} \text{ in } \mathbf{R}^n\}$$



Column Space: Example

Example

Find a matrix A such that $W = \text{Col } A$ where

$$W = \left\{ \begin{bmatrix} x - 2y \\ 3y \\ x + y \end{bmatrix} : x, y \text{ in } \mathbf{R} \right\}.$$

Solution:

$$\begin{aligned} \begin{bmatrix} x - 2y \\ 3y \\ x + y \end{bmatrix} &= x \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + y \begin{bmatrix} -2 \\ 3 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -2 \\ 0 & 3 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \end{aligned}$$



The Contrast Between Nul A and Col A

Example

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 7 \\ 3 & 6 & 10 \\ 0 & 0 & 1 \end{bmatrix}.$$

- (a) The column space of A is a subspace of \mathbf{R}^k where $k = \dots$.
- (b) The null space of A is a subspace of \mathbf{R}^k where $k = \dots$.
- (c) Find a nonzero vector in Col A . (There are infinitely many possibilities.)

$$\text{---} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix} + \text{---} \begin{bmatrix} 2 \\ 4 \\ 6 \\ 0 \end{bmatrix} + \text{---} \begin{bmatrix} 3 \\ 7 \\ 10 \\ 1 \end{bmatrix} = \begin{bmatrix} \\ \\ \\ \end{bmatrix}$$



The Contrast Between Nul A and Col A (cont.)

Example (cont.)

(d) Find a nonzero vector in Nul A. Solve $A\mathbf{x} = \mathbf{0}$ and pick one solution.

$$\begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 7 & 0 \\ 3 & 6 & 10 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \text{ row reduces to } \begin{bmatrix} 1 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$x_1 = -2x_2$$

x_2 is free

\implies let $x_2 = \dots \implies$

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}$$

$$x_3 = 0$$

*Contrast Between Nul A and Col A where A is $m \times n$
(see page 204)*



Null Spaces & Column Spaces: Review

Review

A **subspace** of a vector space V is a subset H of V that has three properties:

- The zero vector of V is in H .
- For each \mathbf{u} and \mathbf{v} in H , $\mathbf{u} + \mathbf{v}$ is in H .
(In this case we say H is closed under vector addition.)
- For each \mathbf{u} in H and each scalar c , $c\mathbf{u}$ is in H .
(In this case we say H is closed under scalar multiplication.)

If the subset H satisfies these three properties, then H itself is a vector space.



Null Spaces & Column Spaces: Review (cont.)

Theorem (1, 2 and 3 in Sections 4.1 & 4.2)

- If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .
- The null space of an $m \times n$ matrix A is a subspace of \mathbf{R}^n .
- The column space of an $m \times n$ matrix A is a subspace of \mathbf{R}^m .



Null Spaces & Column Spaces: Examples

Example

Determine whether each of the following sets is a vector space or provide a counterexample.

$$(a) H = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : x - y = 4 \right\}$$

Solution: Since

$$\begin{bmatrix} 4 \\ 0 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \end{bmatrix}$$

is not in H , H is not a vector space.



Null Spaces & Column Spaces: Examples (cont.)

Example

$$(b) \quad V = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : \begin{array}{l} x - y = 0 \\ y + z = 0 \end{array} \right\}$$

Solution: Rewrite

$$x - y = 0$$

$$y + z = 0$$

as

$$\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

So $V = \text{Nul } A$ where $A = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$. Since $\text{Nul } A$ is a subspace of \mathbf{R}^3 , V is a vector space.



Null Spaces & Column Spaces: Examples (cont.)

Example

$$(c) \ S = \left\{ \begin{bmatrix} x + y \\ 2x - 3y \\ 3y \end{bmatrix} : x, y, z \text{ are real} \right\}$$

One *Solution*: Since

$$\begin{bmatrix} x + y \\ 2x - 3y \\ 3y \end{bmatrix} = x \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + y \begin{bmatrix} 1 \\ -3 \\ 3 \end{bmatrix},$$

$$S = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 3 \end{bmatrix} \right\};$$

therefore S is a vector space by Theorem 1.



Null Spaces & Column Spaces: Examples (cont.)

Another Solution: Since

$$\begin{bmatrix} x + y \\ 2x - 3y \\ 3y \end{bmatrix} = x \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + y \begin{bmatrix} 1 \\ -3 \\ 3 \end{bmatrix},$$

$$S = \text{Col } A \quad \text{where } A = \begin{bmatrix} 1 & 1 \\ 2 & -3 \\ 0 & 3 \end{bmatrix};$$

therefore S is a vector space, since a column space is a vector space.



Kernal and Range of a Linear Transformation

Linear Transformation

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V ;
2. $T(c\mathbf{u}) = cT(\mathbf{u})$ for all \mathbf{u} in V and all scalars c .

Kernal and Range

The *kernel* (or **null space**) of T is the set of all vectors \mathbf{u} in V such that $T(\mathbf{u}) = \mathbf{0}$. The *range* of T is the set of all vectors in W of the form $T(\mathbf{u})$ where \mathbf{u} is in V .

So if $T(\mathbf{x}) = A\mathbf{x}$, $\text{col } A = \text{range of } T$.

