

Math 2331 – Linear Algebra

4.3 Linearly Independent Sets; Bases

Jiwen He

Department of Mathematics, University of Houston

`jiwenhe@math.uh.edu`
`math.uh.edu/~jiwenhe/math2331`



4.3 Linearly Independent Sets; Bases

- Linearly Independent Sets
 - Definition
 - Facts
 - Examples
- A Basis Set: Definition
- A Basis Set: Examples
 - Nul A : Examples and Theorem
 - Col A : Examples and Theorem
- The Spanning Set Theorem



Linearly Independent Sets

Linearly Independent Sets

- A set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ in a vector space V is said to be **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}$$

has only the trivial solution $c_1 = 0, \dots, c_p = 0$.

- The set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ is said to be **linearly dependent** if there exists weights c_1, \dots, c_p , not all 0, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}.$$



Linearly Independent Sets: Facts

The following results from Section 1.7 are still true for more general vectors spaces.

Fact 1

A set containing the zero vector is linearly dependent.

Fact 2

A set of two vectors is linearly dependent if and only if one is a multiple of the other.

Fact 3

A set containing the zero vector is linearly independent.



Linearly Independent Sets: Examples

Example

$\left\{ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 3 & 0 \end{bmatrix} \right\}$ is a linearly _____ set.

Example

$\left\{ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 6 \\ 9 & 11 \end{bmatrix} \right\}$ is a linearly _____ set

since $\begin{bmatrix} 3 & 6 \\ 9 & 11 \end{bmatrix}$ is not a multiple of $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$.



Linearly Independent Sets: Examples

Theorem (4)

An indexed set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ of two or more vectors, with $\mathbf{v}_1 \neq \mathbf{0}$, is linearly dependent if and only if some vector \mathbf{v}_j ($j > 1$) is a linear combination of the preceding vectors $\mathbf{v}_1, \dots, \mathbf{v}_{j-1}$.

Example

Let $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ be a set of vectors in \mathbf{P}_2 where $\mathbf{p}_1(t) = t$, $\mathbf{p}_2(t) = t^2$, and $\mathbf{p}_3(t) = 4t + 2t^2$. Is this a linearly dependent set?


Solution: Since $\mathbf{p}_3 = \text{---}\mathbf{p}_1 + \text{---}\mathbf{p}_2$, $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ is a linearly _____ set.



A Basis Set

Let H be the plane illustrated below. Which of the following are valid descriptions of H ?

- (a) $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ (b) $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_3\}$
(c) $H = \text{Span}\{\mathbf{v}_2, \mathbf{v}_3\}$ (d) $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$



A *basis set* is an “efficient” spanning set containing no unnecessary vectors. In this case, we would consider the linearly independent sets $\{\mathbf{v}_1, \mathbf{v}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_3\}$ to both be examples of basis sets or bases (plural for basis) for H .



A Basis Set: Definition and Examples

A Basis Set

Let H be a subspace of a vector space V . An indexed set of vectors $\beta = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a basis for H if

- i. β is a linearly independent set, and
- ii. $H = \text{Span}\{\mathbf{b}_1, \dots, \mathbf{b}_p\}$.

Example

Let $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$. Show that

$\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is a basis for \mathbf{R}^3 . The set $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is called a **standard basis** for \mathbf{R}^3 .

Solutions: (Review the IMT, page 112)

$$\text{Let } A = [\mathbf{e}_1 \quad \mathbf{e}_2 \quad \mathbf{e}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$



A Basis Set: Definition and Examples

Since A has 3 pivots,

- the columns of A are linearly _____ by the IMT,
- and the columns of A _____ by IMT;
- therefore, $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is a basis for \mathbf{R}^3 .

Example

Let $S = \{1, t, t^2, \dots, t^n\}$. Show that S is a basis for \mathbf{P}_n .

Solution: Any polynomial in \mathbf{P}_n is in span of S . To show that S is linearly independent, assume

$$c_0 \cdot 1 + c_1 \cdot t + \cdots + c_n \cdot t^n = \mathbf{0}.$$

Then $c_0 = c_1 = \cdots = c_n = 0$. Hence S is a basis for \mathbf{P}_n .



A Basis Set: Example

Example

Let $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$, $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}$.

Is $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ a **basis** for \mathbf{R}^3 ?

Solution: Let $A = [\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3] = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$. By row reduction,

$$\begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 1 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 5 \end{bmatrix}$$

and since there are 3 pivots, the columns of A are linearly independent and they span \mathbf{R}^3 by the IMT. Therefore $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is a **basis** for \mathbf{R}^3 .



A Basis Set: Example

Example

Explain why each of the following sets is **not** a basis for \mathbf{R}^3 .

$$(a) \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 7 \end{bmatrix} \right\}$$

Example

$$(b) \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\}$$



Bases for Nul A : Example

Example

Find a basis for Nul A where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

Solution: Row reduce $[A \ \mathbf{0}]$:

$$\begin{bmatrix} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{bmatrix} \implies$$

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5$$

x_2 , x_4 and x_5 are free



Bases for Nul A : Example (cont.)

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 13x_4 - 33x_5 \\ x_2 \\ 6x_4 + 15x_5 \\ x_4 \\ x_5 \end{bmatrix} = \\
 x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \\
 \begin{matrix} \uparrow \\ \mathbf{u} \end{matrix} \qquad \begin{matrix} \uparrow \\ \mathbf{v} \end{matrix} \qquad \begin{matrix} \uparrow \\ \mathbf{w} \end{matrix}$$

Therefore $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is a spanning set for $\text{Nul } A$. In the last section we observed that this set is linearly independent. Therefore $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is a basis for $\text{Nul } A$. The technique used here always provides a linearly independent set.



The Spanning Set Theorem

A basis can be constructed from a spanning set of vectors by discarding vectors which are linear combinations of preceding vectors in the indexed set.

Example

Suppose $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{v}_3 = \begin{bmatrix} -2 \\ -3 \end{bmatrix}$.

Solution: If \mathbf{x} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$, then

$$\begin{aligned} \mathbf{x} &= c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3(\text{---}\mathbf{v}_1 + \text{---}\mathbf{v}_2) \\ &= \text{-----}\mathbf{v}_1 + \text{-----}\mathbf{v}_2 \end{aligned}$$

Therefore,

$$\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}.$$



The Spanning Set Theorem

Theorem (5 The Spanning Set Theorem)

Let

$$S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$$

be a set in V and let

$$H = \text{Span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}.$$

- If one of the vectors in S - say \mathbf{v}_k - is a linear combination of the remaining vectors in S , then the set formed from S by removing \mathbf{v}_k still spans H .
- If $H \neq \{\mathbf{0}\}$, some subset of S is a basis for H .



Bases for Col A: Examples

Example

Find a basis for Col A, where

$$A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{a}_4] = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

Solution: Row reduce:

$$[\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{a}_4] \sim \cdots \sim \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = [\mathbf{b}_1 \ \mathbf{b}_2 \ \mathbf{b}_3 \ \mathbf{b}_4]$$



Bases for Col A: Examples (cont.)

Note that

$$\mathbf{b}_2 = \text{---}\mathbf{b}_1 \quad \text{and} \quad \mathbf{a}_2 = \text{---}\mathbf{a}_1$$

$$\mathbf{b}_4 = 4\mathbf{b}_1 + 5\mathbf{b}_3 \quad \text{and} \quad \mathbf{a}_4 = 4\mathbf{a}_1 + 5\mathbf{a}_3$$

\mathbf{b}_1 and \mathbf{b}_3 are not multiples of each other

\mathbf{a}_1 and \mathbf{a}_3 are not multiples of each other

Elementary row operations on a matrix do not affect the linear dependence relations among the columns of the matrix.

Therefore

$$\text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4\} = \text{Span}\{\mathbf{a}_1, \mathbf{a}_3\}$$

and $\{\mathbf{a}_1, \mathbf{a}_3\}$ is a basis for Col A.



Bases for Col A: Theorem and Example

Theorem (6)

The pivot columns of a matrix A form a basis for Col A.

Example

Let $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} -2 \\ -4 \\ 6 \end{bmatrix}$, $\mathbf{v}_3 = \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$. Find a basis for $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

Solution: Let

$$A = \begin{bmatrix} 1 & -2 & 3 \\ 2 & -4 & 6 \\ -3 & 6 & 9 \end{bmatrix}$$

and note that

$$\text{Col}A = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}.$$



Bases for Col A: Theorem and Example (cont.)

By row reduction, $A \sim \begin{bmatrix} 1 & -2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. Therefore a basis

for $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is $\left\{ \begin{bmatrix} \\ \\ \end{bmatrix}, \begin{bmatrix} \\ \\ \end{bmatrix} \right\}$.



Bases for Nul A & Col A : Review

Review

1. To find a basis for Nul A , use elementary row operations to transform $[A \ \mathbf{0}]$ to an equivalent reduced row echelon form $[B \ \mathbf{0}]$. Use the reduced row echelon form to find parametric form of the general solution to $A\mathbf{x} = \mathbf{0}$. The vectors found in this parametric form of the general solution form a basis for Nul A .
2. A basis for Col A is formed from the pivot columns of A .
Warning: Use the pivot columns of A , not the pivot columns of B , where B is in reduced echelon form and is row equivalent to A .

