

Test 1
Introduction to PDE
MATH 3363-25820 (Fall 2009)

Name and UH-ID: _____

This exam has 4 questions, for a total of 20 points.
Please answer the questions in the spaces provided on the question sheets.
If you run out of room for an answer, continue on the back of the page.

Upon finishing PLEASE write and sign your pledge below:
On my honor I have neither given nor received any aid on this exam.

1 Rules

You may only use pencils, pens, erasers, and straight edges. No calculators, notes, books or other aides are permitted. Scrap paper will be provided. Be sure to show a few key intermediate steps when deriving results - answers only will not get full marks.

2 Given

You may assume the eigenvalues of the Sturm-Liouville problem

$$\begin{aligned} X'' + \lambda X &= 0, & 0 < x < 1 \\ X'(0) &= 0, & X(1) = 0. \end{aligned}$$

are $\lambda_n = (n - \frac{1}{2})^2 \pi^2$ and $X_n(x) = \cos((n - \frac{1}{2})\pi x)$, for $n = 1, 2, \dots$, without derivation.

You may also assume the following orthogonality conditions for m, n positive integers:

$$\int_0^1 \cos((m - \frac{1}{2})\pi x) \cos((n - \frac{1}{2})\pi x) dx = \begin{cases} 1/2, & m = n, \\ 0, & m \neq n. \end{cases}$$

3 Questions

Consider the following heat problem in dimensionless variables

$$\begin{aligned}u_t &= u_{xx} + bx^2, & 0 < x < 1, & \quad t > 0 \\u_x(0, t) &= 0, & u(1, t) &= 1, & \quad t > 0 \\u(x, 0) &= u_0, & 0 < x < 1,\end{aligned}$$

where $b > 0$ and $u_0 > 0$ are constants. This is the heat equation with a source, where the rod is insulated at $x = 0$ and kept at 1 degree at $x = 1$.

- 4 points 1. Derive the steady-state (equilibrium) solution

$$u_E(x) = \frac{b}{12}(1 - x^4) + 1$$

It is insufficient to simply verify that the solution works.

- 3 points 2. (a) Using $u_E(x)$, transform the given heat problem for $u(x, t)$ into the following problem for a function $v(x, t)$:

$$\begin{aligned}v_t &= v_{xx}, & 0 < x < 1, & \quad t > 0 \\v_x(0, t) &= 0, & v(1, t) &= 0, & \quad t > 0 \\v(x, 0) &= f(x), & 0 < x < 1,\end{aligned}$$

where $f(x)$ will be determined by the transformation.

Show your work, which involves writing $v = u - u_E$ and using the information from u and u_E to derive the problem for v .

(b) State $f(x)$ in terms of u_0 , b and x .

- 10 points 3. Derive the solution

$$v(x, t) = \sum_{n=1}^{\infty} v_n(x, t) = \sum_{n=1}^{\infty} A_n \cos\left(\left(n - \frac{1}{2}\right)\pi x\right) e^{-(n - \frac{1}{2})^2 \pi^2 t}$$

and derive equations for A_n in terms of $f(x)$. Be sure to give the intermediate steps: (a) separate variables, (b) write down problems and solve for $X(x)$ (using information from the Given section), (c) solve for $T_n(t)$, put things together, (d) impose the IC, (e) use orthogonality of $\cos\left(\left(n - \frac{1}{2}\right)\pi x\right)$ (see Given section) to find A_n in terms of $f(x)$. Substitute for $f(x)$ from Part 2. You may use (without proof) the following integrals, for any integer n ,

$$\begin{aligned}\int_0^1 (1 - x^4) \cos\left(\left(n - \frac{1}{2}\right)\pi x\right) dx &= \frac{(-1)^{n+1} 96(\pi^2(2n - 1)^2 - 8)}{(2n - 1)^5 \pi^5}, \\ \int_0^1 \cos\left(\left(n - \frac{1}{2}\right)\pi x\right) dx &= \frac{2(-1)^{n+1}}{(2n - 1)\pi}\end{aligned}$$

Name and UH-ID:

3 points

4. (a) Solve for

$$u(x, t) = u_E(x) + \sum_{n=1}^{\infty} u_n(x, t)$$

using the earlier transformations. Write down precisely the functions $u_n(x, t)$.

(b) Find an approximate solution good for large time and state the limit as $t \rightarrow \infty$.

Name and UH-ID:

1.

Name and UH-ID:

2.

Name and UH-ID:

3.

Name and UH-ID:

4.