

Practice Test 1

1 Given

You may assume the eigenvalues of the Sturm-Liouville problem

$$\begin{aligned} X'' + \lambda X &= 0, & 0 < x < 1 \\ X(0) &= 0 & X(1) = 0 \end{aligned}$$

are $\lambda_n = n^2\pi^2$ and $X_n(x) = \sin(nx)$, for $n = 1, 2, \dots$, without derivation.

You may also assume the following orthogonality conditions for m, n positive integers:

$$\int_0^1 \sin(m\pi x) \sin(n\pi x) dx = \begin{cases} 1/2, & m = n \neq 0, \\ 0, & m \neq n. \end{cases}$$

$$\int_0^1 \cos(m\pi x) \cos(n\pi x) dx = \begin{cases} 1/2, & m = n \neq 0, \\ 0, & m \neq n. \end{cases}$$

2 Question

Consider the following heat problem in dimensionless variables

$$\begin{aligned} u_t &= u_{xx} + \frac{\pi^2}{4}u - b, & 0 < x < 1, & \quad t > 0 \\ u(0, t) &= 0, & u(1, t) = 0, & \quad t > 0 \\ u(x, 0) &= u_0 & 0 < x < 1. \end{aligned}$$

(a) [3 points] Explain in terms of a heated rod precisely what the problem models mathematically.

(b) [3 points] Derive the equilibrium solution

$$u_E(x) = \frac{4b}{\pi^2} \left[1 - \cos\left(\frac{\pi x}{2}\right) - \sin\left(\frac{\pi x}{2}\right) \right]$$

It is insufficient to simply verify that the solution works.

(c) [3 points] Using $u_E(x)$, transform the given heat problem for $u(x, t)$ into the following problem for a function $v(x, t)$:

$$\begin{aligned} v_t &= v_{xx} + \frac{\pi^2}{4}v, & 0 < x < 1, & \quad t > 0 \\ v(0, t) &= 0, & v(1, t) &= 0, & \quad t > 0 \\ v(x, 0) &= f(x) & 0 < x < 1. \end{aligned}$$

where $f(x)$ will be determined by the transformation.

(d) [3 points] For an appropriate value of α show that the transformation $w(x, t) = e^{\alpha t}v(x, t)$ further simplifies the problem to

$$\begin{aligned} w_t &= w_{xx}, & 0 < x < 1, & \quad t > 0 \\ w(0, t) &= 0, & w(1, t) &= 0, & \quad t > 0 \\ w(x, 0) &= f(x) & 0 < x < 1. \end{aligned}$$

(e) [8 points] Derive the solution

$$w(x, t) = \sum_{n=1}^{\infty} w_n(x, t) = \sum_{n=1}^{\infty} \frac{2}{\pi} \left(\frac{2(u_0 - 4b/\pi^2)}{2n-1} + \frac{32b(2n-1)}{\pi^2(4n-3)(4n-1)} \right) e^{-(2n-1)^2\pi^2 t} \sin((2n-1)\pi x)$$

and hence solve for $u(x, t) = u_E(x) + \sum_{n=1}^{\infty} w_n(x, t)$ using the earlier transformations.