

Math 1432 - 13209

Jeff Morgan (651) PGH - 11-noon MWF

<http://www.math.uh.edu/~jmorgan/Math1432>

Read the Syllabus!

Test 1 is available online at CourseWare.

Practice Test 1 is also available and counts as an online quiz.

Homework 1 is posted on the course homepage and due next Wednesday.

EMCF01 is due tomorrow morning at 9am. **A help video was posted.**

EMCF02 is due on Saturday morning at 9am.

Your first written quiz will be given in lab/workshop on Friday.

Online Quiz 1 is Available on CourseWare.

Poppers start week 3. Get your packet at the UC Book Store.

Access Codes are required to continue to access the online text, EMCF, quizzes, etc... See the deadline on the course homepage.



CourseWare = <http://www.casa.uh.edu>

<http://www.math.uh.edu/~jmorgan/Math1432>

Math 1432 - 13209

Jeff Morgan - jmorgan@math.uh.edu

Read the **Syllabus**

Use the **Discussion Board on CourseWare** to get and give help.

Lecture notes/videos, additional help material, course announcements, homework and EMCFs will be posted in the calendar below. **Note:** Practice Tests count the same as online quizzes.

Course Calendar

Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
<p>January 13</p> <p>Note: Practice Test 1 counts the same as an online quiz. Exam 1 counts as a major exam.</p>	<p>14</p> <p>Notes</p> <p>Exam 1, PT1 and all Online Quizzes are open</p>	<p>15</p> <p>UH events this week</p> <p>Examples from 7.1 that will help with EMCF01</p>	<p>16</p> <p>Blank Slides</p> <p>Homework 1 posted</p> <p><i>video</i></p>	<p>17</p> <p>EMCF01 due at 9am</p>	<p>18</p> <p>Quiz in lab/workshop</p>	<p>19</p> <p>EMCF02 due at 9am</p>
<p>20</p>	<p>21</p> <p>MLK Day No Class</p>	<p>22.</p> <p>Last day to add</p>	<p>23</p> <p>EMCF03 due at 9am</p> <p>Homework 1 due in lab/workshop</p> <p>Homework 2 posted</p>	<p>24</p> <p>Exam 1 and PT1 close</p>	<p>25</p> <p>EMCF04 due at 9am</p> <p>Quiz in lab/workshop</p>	<p>26</p> <p>Quiz 1 closes (7.1-7.2)</p>

We started discussing inverse functions on Monday. The notes are posted. **In addition, I posted a video that might help with EMCF01.**

Today... More Inverse Functions and Logarithms

Make sure you read sections 7.2 and 7.3.

"Review" What were you taught about Logarithms in
highschool algebra? (base 10 answers...)

$$\log_{10}(1) = 0$$

$$\log_{10}(10) = 1$$

$$\log_{10}\left(\frac{1}{10}\right) = -1$$

$$\log_{10}(x) = y \text{ iff } \underline{x = 10^y}$$

$$\log_{10}(100) = 2$$

$$\log_{10}(10^x) = x$$

"Review" Properties: If A and B are positive real numbers, and r is a real number, then

$$\textcircled{*} \log_{10}(AB) = \log_{10}(A) + \log_{10}(B)$$

$$\log_{10}\left(\frac{A}{B}\right) = \log_{10}(A) - \log_{10}(B)$$

$$\log_{10}(A^r) = r \log_{10}(A)$$

$$\log_{10}\left(\frac{1}{A}\right) = -\log_{10}(A)$$

"Review"

For arbitrary bases...

Properties: If A and B are positive real numbers, r is a real number, and n is a positive real number which IS NOT 1, then

$$\log_n(1) = 0$$

$$\log_n(n^x) = x$$

~~A~~

$$\log_n(AB) = \log_n(A) + \log_n(B)$$

$$\log_n\left(\frac{A}{B}\right) = \log_n(A) - \log_n(B)$$

Also, for $x > 0$

$$\log_n(x) = y$$

iff

$$x = n^y$$

"Review" Change of base:

$$\log_a(b) = \frac{\log_c(b)}{\log_c(a)}$$

$a, b, c > 0$ and $a, c \neq 1$

$$\log_{10}(x) = \frac{\log_2(x)}{\log_2(10)}$$

The Truth!

The defining property of logarithmic functions:



If $n > 0$ with $n \neq 1$ and $A, B > 0$ then

$$\log_n (AB) = \log_n (A) + \log_n (B)$$

All other properties follow from this property.

So...

We say that a real valued function L defined on the positive real numbers is a logarithmic function if and only if L is not the zero function and

$$L(AB) = L(A) + L(B)$$

(all other properties follow from this one)

Question: Does $L'(x)$ exist for $x > 0$? If so, what is $L'(x)$?

A:

Let $x, y > 0$. Then

$$L(xy) = L(x) + L(y)$$

$$\frac{d}{dy} L(xy) = \frac{d}{dy} [L(x) + L(y)]$$

$$L'(xy) \cdot x = 0 + L'(y)$$

Subst $y = 1$.

$$L'(x) \cdot x = L'(1)$$

$$\Rightarrow L'(x) = \frac{L'(1)}{x}$$

number

In fact, we could prove...

If L is a logarithmic function then L is differentiable and there is a constant

C so that

$$L'(x) = C \cdot \frac{1}{x}$$

$x > 0$

As a result, all logarithmic functions are invertible. Why?

The constant depends upon the base of the logarithm.

Questions:

1. Is there a function whose derivative is $\frac{1}{x}$?


$$\frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}$$

yes

$$L(x) = \int_1^x \frac{1}{t} dt$$

2. Is this a logarithmic function?

yes



3. Is there any other logarithmic function with this derivative?

No

Question : How can we show that $L(x) = \int_1^x \frac{1}{t} dt$

is a logarithmic function?

Note: $L(1) = 0$

Answer : We need to show $L(AB) = L(A) + L(B)$

for $A, B > 0$. *spse $A > 0$ and $x > 0$.*

$$\frac{d}{dx} L(Ax) = L'(Ax) \cdot A = \frac{1}{Ax} \cdot A = \frac{1}{x}$$

$$\frac{d}{dx} [L(A) + L(x)] = 0 + \frac{1}{x}$$

← equal.

∴ there is a value C so that

$$L(Ax) = L(A) + L(x) + C.$$

Subs $x=1, A=1$.

$$0 = 0 + 0 + C \\ \Rightarrow C = 0$$

$$\therefore L(Ax) = L(A) + L(x). \quad \#$$

We refer to this special logarithmic function as the natural logarithm, and we denote it as $\ln(x)$.

$\ln(x)$ satisfies the logarithmic properties and...

Note:

$$\ln(|u|) = \begin{cases} \ln(-u), & u < 0 \\ \ln(u), & u > 0 \end{cases}$$

$$\frac{d}{du} \ln(|u|) = \begin{cases} \frac{1}{-u} \cdot (-1), & u < 0 \\ \frac{1}{u}, & u > 0 \end{cases}$$

$$= \begin{cases} \frac{1}{u}, & u < 0 \\ \frac{1}{u}, & u > 0 \end{cases}$$

$$\ln(x) = \int_1^x \frac{1}{t} dt, \quad x > 0$$

$$\frac{d}{dx} \ln(x) = \frac{1}{x}$$

$$\frac{d}{dx} \ln(u(x)) = \frac{1}{u(x)} \cdot u'(x)$$

$$\int \frac{1}{u} du = \ln(|u|) + C$$

Example: Solve the equation $\ln((3x+3)(x+5)) = 2\ln(x+5)$

$$\ln \left(\underbrace{(3x+3)(x+5)} \right) = \ln \left(\underbrace{(x+5)^2} \right)$$

$\therefore (3x+3)(x+5) = (x+5)^2$

Note: ~~$x = -5$~~ works.

But $x = -5$ is not allowed
in the original equation
b/c $\ln(0)$ d.n.e.

$$3x + 3 = x + 5$$

$$2x = 2 \Rightarrow x = 1.$$

$$\ln((3x+3)(x+5)) = 2\ln(x+5)$$

Subst $x = 1.$

$$\ln(6 \cdot 6) \stackrel{?}{=} 2 \ln(6)$$

Yes. $\ln(6^2)$

\therefore The sol'n is $\boxed{x=1.}$

Examples:

$$\frac{d}{dx} \ln(x^2 + 1) = \frac{1}{x^2 + 1} \cdot 2x = \frac{2x}{x^2 + 1}$$

$$\frac{d}{dx} \ln(u(x)) = \frac{1}{u(x)} u'(x)$$

$$\int \frac{1}{2x+3} dx = \frac{1}{2} \int \frac{1}{2x+3} \cdot 2 dx$$

$$\int \frac{1}{u} du = \ln(|u|) + C = \frac{1}{2} \ln(|2x+3|) + C$$

$$\int \frac{x}{3x^2+5} dx = \frac{1}{6} \int \frac{1}{3x^2+5} \cdot 6x dx$$

$$= \frac{1}{6} \ln(|3x^2+5|) + C$$

$$= \frac{1}{6} \ln(3x^2+5) + C$$

Since $3x^2+5 > 0$.

Example: Give the y-intercept of the tangent line to the graph of $f(x) = \ln(3x^2 - 2)$ at $x = 1$.

1. Get the tangent line.

$$\text{Point: } (1, f(1)) = (1, 0)$$

$$\text{slope: } f'(1) = 6$$

$$f'(x) = \frac{6x}{3x^2 - 2} \Rightarrow$$

Equation: $y = 6(x - 1)$

$$y = 6x - 6$$

2. Get y-intercept. -6

$$(0, -6)$$

Can we graph $y = \ln(x)$?

Domain: $x > 0$

Behavior near ∞ : $\lim_{x \rightarrow \infty} \ln(x) = \infty$

Behavior near 0: $\lim_{x \rightarrow 0^+} \ln(x) = -\infty$

Range: $(-\infty, \infty)$

Slope and Concavity: $\frac{d}{dx} \ln(x) = \frac{1}{x} > 0$

x-intercept:


$$\ln(1) = 0$$

$$(1, 0)$$


$$\frac{d^2}{dx^2} \ln(x) = -\frac{1}{x^2} < 0$$

Concave down.

But it grows very slowly.



The Graph...



Note: $\ln(x)$ is an invertible function.

We will study the inverse of this function in a few days.

What is the "base" of $\ln(x)$?

$$\ln(x) = \log_e(x)$$

$e = 2.718281828459045235360287471352662$
 $49775724709369995957496696762772407\dots$

(irrational)

<http://www.math.uh.edu/~jmorgan/Math1432>