

Reminders...

Online Quiz 3 closes this Saturday. Other **Online Quizzes** are available.

Test 2 is coming soon. **Review problems** and **video solutions** are posted, and there is a **Practice Test 2** on CourseWare.

EMCFs are due on each MWF.

Homework is posted and due on Monday.

There is a written **Quiz** in lab on Friday.

**Please tell your high school friends and former
teachers about our Mathematics Contest**

(middle school and high school students are welcome)

**February 9th
University of Houston**

Free

<http://mathcontest.uh.edu>

Today

Final comments for integrating products
of sine and cosine.

(Integrating Powers and Products of
Secant and Tangent Functions

Examples of simple products with different arguments:

$$\int \sin(\underline{2x}) \cos(\underline{3x}) dx = \int \frac{1}{2} (\sin(5x) + \sin(-x)) dx$$

$$= \frac{1}{2} \int (\sin(5x) - \sin(x)) dx$$

$$\int \cos(\underline{2x}) \cos(\underline{3x}) dx = \frac{1}{2} \cdot \left(-\frac{1}{5}\right) \cos(5x) + \frac{1}{2} \cos(x) + C$$

$$= -\frac{1}{10} \cos(5x) + \frac{1}{2} \cos(x) + C$$

$$\int \sin(2x) \sin(3x) dx =$$

$$\frac{1}{2} (\sin(A+B) + \sin(A-B)) = \sin(A) \cos(B)$$

$$\frac{1}{2} (\cos(A+B) + \cos(A-B)) = \cos(A) \cos(B)$$

$$= \int \frac{1}{2} (\cos(5x) + \cos(-x)) dx$$

$$= \frac{1}{2} \int (\cos(5x) + \cos(x)) dx$$

$$= \frac{1}{10} \sin(5x) + \frac{1}{2} \sin(x) + C$$

Useful Identities:

$$\cos(A+B) = \underbrace{\cos(A)\cos(B)} - \underbrace{\sin(A)\sin(B)}$$

$$\begin{aligned}\cos(A-B) &= \cos(A)\cos(-B) - \sin(A)\sin(-B) \\ &= \underbrace{\cos(A)\cos(B)} + \underbrace{\sin(A)\sin(B)}\end{aligned}$$

ADD $\cos(A+B) + \cos(A-B) = 2\cos(A)\cos(B)$

$$\frac{1}{2}(\cos(A+B) + \cos(A-B)) = \cos(A)\cos(B)$$

$$\sin(A+B) = \underbrace{\sin(A)\cos(B)} + \underbrace{\sin(B)\cos(A)}$$

$$\begin{aligned}\sin(A-B) &= \sin(A)\cos(-B) + \sin(-B)\cos(A) \\ &= \underbrace{\sin(A)\cos(B)} - \underbrace{\sin(B)\cos(A)}\end{aligned}$$

ADD

$$\sin(A+B) + \sin(A-B) = 2\sin(A)\cos(B)$$

$$\frac{1}{2}(\sin(A+B) + \sin(A-B)) = \sin(A)\cos(B)$$

Even Powers of secant and tangent:

Examples: $\int \sec^4(x) dx =$

$$\int \tan^4(x) dx =$$

Fundamental Identity:

$$1 + \tan^2(x) = \sec^2(x)$$

Example: $\int \tan^2(x) \sec^4(x) dx =$

$$= \int \underbrace{\tan^2(x) \sec^2(x)} \underbrace{\sec^2(x) dx}_{\text{derivative of } \tan(x)}$$

↓
convert to
tan(x) terms.

$$= \int \tan^2(x) (1 + \tan^2(x)) \underbrace{\sec^2(x) dx}$$

$$= \int \tan^2(x) \sec^2(x) dx + \int \tan^4(x) \sec^2(x) dx$$

$$= \frac{1}{3} \tan^3(x) + \frac{1}{5} \tan^5(x) + C$$

General strategy for calculating

$$\int \tan^n(x) \sec^m(x) dx$$

when m is even.

(peel off $\sec^2(x)$, and then turn everything else into powers of $\tan(x)$)

Odd Powers Part 3

Example: $\int \sec^3(x) dx = \int \underbrace{\sec(x)}_u \underbrace{\sec^2(x) dx}_{dv}$

$u = \sec(x)$	$du = \sec(x) \tan(x) dx$
$dv = \sec^2(x) dx$	$v = \tan(x)$

$$= \sec(x) \tan(x) - \int \sec(x) \tan^2(x) dx$$

$$\Rightarrow \int \sec^3(x) dx = \sec(x) \tan(x) - \int \sec(x) (\sec^2(x) - 1) dx$$
$$= \sec(x) \tan(x) - \int \sec^3(x) dx + \int \sec(x) dx$$

$$2 \int \sec^3(x) dx = \sec(x) \tan(x) + \ln(|\sec(x) + \tan(x)|) + \tilde{C}$$

$$\Rightarrow \int \sec^3(x) dx = \frac{1}{2} \sec(x) \tan(x) + \frac{1}{2} \ln(|\sec(x) + \tan(x)|) + C$$

General strategy for calculating

$$\int \tan^n(x) \sec^m(x) dx$$

when n is even and m is odd.

(the hardest case - this typically involves parts)

General strategy for calculating

$$\int \tan^n(x) \sec^m(x) dx$$

when n is odd and m is odd.

(peel off a $\sec(x)\tan(x)$, and then turn everything else into powers of $\sec(x)$)

Example: $\int \tan^3(x) \sec^3(x) dx =$

$$= \int \tan^2(x) \sec^2(x) \sec(x) \tan(x) dx$$

$$= \int (\sec^2(x) - 1) \sec^2(x) \sec(x) \tan(x) dx$$

$$= \int \sec^4(x) \sec(x) \tan(x) dx - \int \sec^2(x) \sec(x) \tan(x) dx$$

$$= \frac{1}{5} \sec^5(x) - \frac{1}{3} \sec^3(x) + C$$