

Recall: Polar Coordinates

Standard form


$$r = \sqrt{x^2 + y^2}$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

(x, y) in Q1 or Q4

Always


$$x = r \cos(\theta)$$
$$y = r \sin(\theta)$$




Popper 09

1. 0.125

Polar equations for 3 different types of circles.



II. $r = 2a \sin(\theta)$



$$x^2 + (y-a)^2 = a^2$$

$$r^2 \cos^2(\theta) + (r \sin(\theta) - a)^2 = a^2$$

$$\underline{r^2 \cos^2(\theta) + r^2 \sin^2(\theta) - 2ar \sin(\theta) + a^2 = a^2}$$

$$r^2 - 2ar \sin(\theta) = 0$$

$$\underline{r^2 = 2ar \sin(\theta)}$$


$\boxed{r = 2a \sin(\theta)}$

in polar coord.

III.

$$(x-a)^2 + y^2 = a^2$$

$\boxed{r = 2a \cos(\theta)}$




Ex: Graph the polar curve

$$r = 3 \sin(\theta)$$

$r = 2a \sin(\theta)$

circle of radius $|a|$ centered at $(0, a)$

↓
Circle of radius $\frac{3}{2}$
centered at $(0, \frac{3}{2})$



Polar Flowers

Example:

Plot the polar curve

$$r = \sin(3\theta)$$

Popper 09

2. 0

we are not in

Kansas
any more.

Step 1

Create a plot in the θr plane.
(this is NOT the polar plot)



Graph $r = \sin(3\theta)$
here

$$\bullet \Leftrightarrow \underline{\underline{0}}$$

Step 2

Interpret the plot above to create
the polar plot.

3 petal
flower



Geogebra Exploration:

$$\begin{cases} r = a \cos(m\theta) \\ r = a \sin(m\theta) \end{cases}$$



See the applets
linked from the
course homepage
after class.

Polar graphs
that produce
flowers

$$\begin{cases} r = a \cos(m\theta) \\ r = a \sin(m\theta) \end{cases}$$


$a \nearrow$ is a real #, $a \neq 0$,
and m is a positive integer.

Fundamental Question: How do the values
 a and m effect the graph?


m odd \Rightarrow m petals | m even \Rightarrow
 $2m$ petals.

$|a|$ gives the distance of each
tip from the origin.


The petals are equally spaced.




$$r = 2 \cos(2\theta)$$



$$r = 2 \cos(5\theta)$$



$$r = 2 \sin(4\theta)$$



$$r = 2 \sin(3\theta)$$

Popper 09

3. Give the number of petals on the flower

$$r = 3 \cos(4\theta).$$

4. Give the number of petals on the flower

$$r = 4 \cos(5\theta).$$

5. Give the furthest distance ^{from the origin} to a tip of a petals on the


$$\text{flower } r = 6 \cos(4\theta).$$

Example:


Plot the polar curve

$$r = 3 + 4\cos(\theta)$$


Step 1
Create a plot in the θr plane.
(this is NOT the polar plot)



Step 2
Interpret the plot above to create
the polar plot.



$$r = 3 + 4\cos(\theta)$$








Example: Geogebra Investigation

Investigate

$$r = 3 + b \cos(\theta)$$

See the link from the course homepage after class.

for b between 1 and 5.



You can create similar graphs.

Geogebra Exploration:

$$r = a + b \cos(\theta)$$

and

you

$$r = a + b \sin(\theta)$$

See the applets
linked from the
course homepage
after class.

Polar curves of the form

$$r = a + b \cos(\theta)$$

and

$$r = a + b \sin(\theta)$$

$$|a| = |b|$$


$$|a| > |b|$$

are Cardioids, Limacons with dimples, and Limacons with inner loops.


$$|a| < |b|$$

Look Below...


$$r = a + b \cos(\theta)$$



Cardioid
 $|a|=|b|$



Limacon with dimple
 $|a|>|b|$ (dent)



Limacon with inner loop
 $|a|<|b|$

Note: The cosine versions can be reflected across the y axis if b is negative.