

Answers to Odd-Numbered Problems

CHAPTER 1

Exercises 1.1

1. (a) ordinary, first order
(c) partial, second order
(e) ordinary, third order
(g) ordinary, second order
3. Both y and z are solutions.
5. Both y and z are solutions.
7. Both u_1 and u_2 are solutions.
9. u_1 is a solution; u_2 is not a solution.
11. $y = 16x^2 + C_1x + C_2$
13. $y = Ce^{3x}$.
15. $r = 2, -2$; $y_1(x) = e^{2x}$ and $y_2(x) = e^{-2x}$ are solutions.
17. $r = 3$; $y(x) = e^{3x}$ is a solution.
19. No real values of r ; $r = 1 \pm 2i$ are complex values.
21. $r = 3, -3$; $y_1(x) = x^3$ and $y_2(x) = x^{-3}$ are solutions.

Exercises 1.3

1. (b) $y = 2e^{5x}$.
3. (b) $y = \frac{e}{e - 2e^x}$.
5. (b) $y = -\sin 3x + \frac{1}{3} \cos 3x$.
7. (b) $y = -\frac{17}{4}x + 9\sqrt{x}$.
(c) y' and y'' are not defined at $x = 0$; there is no solution to $y'(0) = 2$.
11. $xy' - 3y + 3 = 0$.

13. $y' = \frac{2y^3 - 6}{3xy^2}$
15. $y' - 2y = -4e^{-2x}$.
17. $y'' = 0$.
19. $y'' - 4y' + 4y = 0$.
21. $x^2y'' + xy' - y = 0$.
23. $y'' + 9y = 0$.
25. $y''' = 0$.

CHAPTER 2

Exercises 2.1

1. $y = -\frac{1}{2} + Ce^{2x}$.
3. $y = 1 + Ce^{-x^2}$.
5. $y = e^{-x} + Ce^x$.
7. $y = x^{-2} \sin x + Cx^{-2}$.
9. $y = \frac{2}{9}(x+1)^{5/2} + C(x+1)^{-2}$.
11. $y = \sin x \cos x + C \cos x = \frac{1}{2} \sin 2x + C \cos x$.
13. $y = e^x + \frac{C}{x}$.
15. $y = x(\ln x)^2 + Cx$.
17. $y = 1 + Ce^{-e^x}$.
19. $y = x - 1 + 2e^{-x}$.
21. $y = \frac{\ln(1+e^x)}{e^x} + (e - \ln 2)e^{-x}$.
23. $y = \frac{5 - \cos 2x}{2 \sin x}$.

Exercises 2.2

1. $y = \left(\frac{x^2}{4} + C\right)^2.$
3. $\tan^{-1} y = x^3 + C$ or $y = \tan(x^3 + C).$
5. $\cot y = \ln \sqrt{\frac{1-x}{1+x}} + C.$
7. $e^{-y} = e^x - xe^x + C.$
9. $\sqrt{y^2 - 1} = \frac{x}{1 + Cx}.$
11. $y^2 = C(\ln x)^2 - 1.$
13. $\ln |y| = -\ln |x| - \frac{1}{x} - 1.$
15. $y = xe^{x^2-1}.$
17. $y + \ln |y| = \frac{1}{3}x^3 - x - 5.$
19. $y = \frac{x + C}{1 - Cx}$

Exercises 2.3

1. $y = \frac{2}{Cx - 3x^3}.$
3. $y = (Ce^{2x} - e^2)^2.$
5. $y = \frac{1}{\sqrt[3]{Cx^3 - 2x^3 \ln x}}.$
7. $y^2 = Cx + x^2.$
9. $x \ln x + \frac{x + y}{e^{y/x}} = Cx.$
11. $\csc(y/x) - \cot(y/x) = Cx.$
13. $y^2 = \frac{C}{1 + x^2} - 1.$
15. $y = \frac{\ln |\sec x + \tan x| + C}{x}.$
17. $y + \ln |1 - y| = C - x - \ln |1 + x|.$
19. $y = -x \ln(C - \ln x).$

21. $y = C(3x^2 + 1)^{1/3} - 3.$

23. $y = \frac{1}{Cx + \ln x + 1}.$

25. $2y^3 = x^3 - Cx.$

27. (a) $u = \sin y$ (b) $\sin y = e^{-x^2}(4x + C).$

Exercises 2.4.1

1. $x^2 + 3y^2 = C.$

3. $\frac{x^2}{2} + y^2 - 4y = C.$

5. $y^2 = \ln(\sin^2 x) + C.$

7. $y = -\frac{1}{2}x^2 + C.$

9. $\frac{x^2}{2} + y^2 = C$; ellipses, center at the origin, major axis horizontal.

11. $x^2 + y^2 - Cy = 0.$

Exercises 2.4.2

1. (a) $A(t) = 50 \left(\frac{9}{10} \right)^{t/2} \approx 50e^{-0.05268t}.$ (b) $A(4) = 50 \left(\frac{9}{10} \right)^2 = 40.5$ grams.

(c) $T \approx 13.16$ hours.

3. $t = \frac{2 \ln 10}{\ln 2} \approx 6.64$ hours.

5. (a) $P(t) \approx 0.25e^{0.0421t}$ (b) ≈ 1.6573 square centimeters (c) ≈ 16.464 hours

7. (a) $P(t) \approx 4.5e^{0.01438t}.$ (b) 48.19 years (c) ≈ 6.17 billion.

Exercises 2.4.3

1. (a) $40.1^\circ.$ (b) 1.62 minutes.

3. (a) $u(t) = 150 - 100e^{\frac{t}{10} \ln(3/4)} = 150 - 100 \left(\frac{3}{4} \right)^{t/10}$

(b) $t = \frac{10 \ln(1/2)}{\ln(3/4)} \approx 24.09$ minutes

(c) The temperature will never reach 200° ; $\lim_{t \rightarrow \infty} u(t) = 150$

5. (a) Approximately 12:12 (b) Approximately 12:48

Exercises 2.4.4

1. (a) $v = \left(v_0 + \frac{g}{r}\right) e^{-rt} - \frac{g}{r}$ (b) $\lim_{t \rightarrow \infty} v = -\frac{g}{r}$.

(c) $y = y_0 + \frac{1}{r} \left(v_0 + \frac{g}{r}\right) (1 - e^{-rt}) - \frac{g}{r}t$

3. $k \approx 17.8$

Exercises 2.4.5

1. (a) $A(t) = 10,000 (1 - e^{-t/200})$ (b) $t = 200 \ln 5 \approx 322$ minutes

3. (a) $A(t) = \frac{9}{2} (1 - e^{-t/150})$ (b) $t = 150 \ln 3 \approx 165$ minutes

5. (a) $A(t) = \frac{3}{20} t(100 - t)$ (b) $\max = A(50) = 375$

Exercises 2.4.6

1. (a) 3259 people. (b) ≈ 6.89 days.

3. (a) $\frac{d^2y}{dt^2} = k \frac{dy}{dt} (M - 2y)$; $\frac{dy}{dt} > 0$ for $0 < y < M/2$, $\frac{dy}{dt} < 0$ for $y > M/2$.

dy/dt has a maximum when $y = M/2$

5. $k \approx 0.0006$

CHAPTER 3

Exercises 3.2

1. Yes

3. Yes

5. Yes

7. (a) $r = -1$, $r = 4$.

(b) Fundamental set: $y_1(x) = x^{-1}$, $y_2(x) = x^4$; general solution: $y = C_1 x^{-1} + C_2 x^4$.

(c) $y = \frac{9}{5} x^{-1} + \frac{1}{5} x^4$.

(d) The trivial solution: $y \equiv 0$.

9. $y'' - 2y' - 3y = 0$.
11. $y'' = 0$.
13. $x^2 y'' - 2x y' + 2y = 0$.
15. $W[y_1, y_2](x) = e^{-\int_a^x p(t) dt} \neq 0$ for all x .
17. $\{y_1(x) = x, y_2(x) = x^2\}$.
19. $\{y_1(x) = e^{x^2}, y_2(x) = e^{-x^2}\}$.
21. $\alpha\delta - \beta\gamma \neq 0$.
23. $W[y_1 + y_2, y_1 - y_2] = -2W[y_1, y_2]$.
25. Set $u(x) = \frac{y_2(x)}{y_1(x)}$. Then

$$u'(x) = \frac{y_1 y_2' - y_2 y_1'}{y_1^2} = \frac{W[y_1, y_2]}{y_1^2} \equiv 0.$$

therefore, $u \equiv \lambda$ constant, which implies that $y_2 = \lambda y_1$.

Exercises 3.3

1. $y = C_1 e^{2x} + C_2 e^{-4x}$.
3. $y = C_1 e^{5x} + C_2 x e^{5x}$.
5. $y = e^{-2x} [C_1 \cos 3x + C_2 \sin 3x]$.
7. $y = C_1 + C_2 e^{-2x}$.
9. $y = C_1 e^{2\sqrt{3}x} + C_2 e^{-2\sqrt{3}x}$.
11. $y = e^x [C_1 \cos x + C_2 \sin x]$.
13. $y = C_1 e^{6x} + C_2 e^{-5x}$.
15. $y = e^{-x/2} [C_1 \cos x/2 + C_2 \sin x/2]$.
17. $y = C_1 e^{4x} + C_2 x e^{4x}$.
19. $y = 2e^{2x} - e^{3x}$.
21. $y = -3e^{-x} - 2xe^{-x}$.
23. $y = -e^x \cos x$.
25. $y'' + 3y' - 10y = 0$.

27. $y'' + 4y = 0.$

29. $y'' - \frac{5}{2}y' + y = 0.$

31. $y'' + 2y' + 10y = 0.$

33. $y'' + 16y = 0.$

35. $y = (1 + \beta)e^{x/2} + (1 - \beta)e^{-x/2}; \beta = -1.$

37. If the roots of $r^2 + ar + b = 0$ are real (real and unequal, or real and equal), then they are negative; r negative implies $e^{rx} \rightarrow 0$ and $xe^{rx} \rightarrow 0$ as $x \rightarrow \infty$. If the roots are complex conjugates, then they have negative real part and α negative implies $e^{\alpha x} \cos \beta x \rightarrow 0$ and $e^{\alpha x} \sin \beta x \rightarrow 0$ as $x \rightarrow \infty$.

39. Suppose that $a > 0$ and $b = 0$. The general solution of the differential equation is

$$y = C_1 + C_2 e^{-ax} \quad \text{and} \quad \lim_{x \rightarrow \infty} y = C_1.$$

The solution that satisfies the initial conditions is: $y = \left(\alpha + \frac{\beta}{a}\right) - \frac{\beta}{a} e^{-ax}; k = \alpha + \frac{\beta}{a}.$

41. $r_1, r_2 = \frac{-a \pm \sqrt{a^2 - 4b}}{2} = \frac{-a}{2} \pm \frac{\sqrt{a^2 - 4b}}{2} = \alpha \pm \beta.$

General solution:

$$\begin{aligned} y = C_1 e^{(\alpha+\beta)x} + C_2 e^{(\alpha-\beta)x} &= C_1 e^{\alpha x} e^{\beta x} + C_2 e^{\alpha x} e^{-\beta x} \\ &= e^{\alpha x} \left[(C_1 + C_2) \frac{e^{\beta x} + e^{-\beta x}}{2} + (C_1 - C_2) \frac{e^{\beta x} - e^{-\beta x}}{2} \right] \\ &= e^{\alpha x} (K_1 \cosh \beta x + K_2 \sinh \beta x). \end{aligned}$$

43. $y = C_1 x^{-2} + C_2 x^4.$

45. $y = C_1 x^2 + C_2 x^2 \ln x.$

Exercises 3.4

1. $z(x) = x^2 \ln x + \frac{1}{2}; \quad y = C_1 x^2 + C_2 x^{-1} + x^2 \ln x + \frac{1}{2}.$

3. $z(x) = -x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2; \quad y = C_1 x + C_2 x^2 - x^2 \ln x + \frac{1}{2} x^2 (\ln x)^2.$

5. $z(x) = -(1 + x^2); \quad y = C_1 x + C_2 e^x - (1 + x^2).$

7. $y = C_1 e^{-x} + C_2 e^{2x} - \frac{2}{3} x e^{-x}.$

9. $y = C_1 \cos 2x + C_2 \sin 2x - \frac{1}{4} \cos 2x \ln(\cos 2x) + \frac{1}{2}x \sin 2x.$
11. $y = C_1 e^x + C_2 x e^x - e^x \cos x.$
13. $y = C_1 e^{-2x} + C_2 x e^{-2x} - e^{-2x} \ln x.$
15. $y = C_1 \cos 3x + C_2 \sin 3x + \sin 3x \ln(\sec 3x + \tan 3x) - 1.$
17. $y = C_1 x + C_2 x^{-1} + x \ln x.$
19. $y = C_1 x + C_2 x \ln x + x^2.$

Exercises 3.5

1. $y = C_1 e^{-x} + C_2 e^{3x} - e^{2x}.$
3. $y = C_1 e^{-3x} + C_2 x e^{-3x} + \frac{1}{4}e^{3x}.$
5. $y = C_1 e^{-2x} + C_2 - \frac{1}{2} \cos 2x - \frac{1}{2} \sin 2x.$
7. $y = C_1 e^{-x/2} + C_2 e^{-x} + x^2 - 6x + 14 - \frac{9}{10} \cos x - \frac{3}{10} \sin x.$
9. $y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{1}{2}x + \frac{1}{4}.$
11. $y = C_1 e^{-2x} + C_2 e^{-4x} + \frac{3}{2}x e^{-2x}.$
13. $y = C_1 \cos 3x + C_2 \sin 3x + \frac{2}{3} + \frac{1}{162} (9x^2 - 6x + 1) e^{3x}.$
15. $y = e^x (C_1 \cos 2x + C_2 \sin 2x) - \frac{1}{10} e^{-x} \cos 2x - \frac{1}{20} e^{-x} \sin 2x.$
17. $y = e^x - \frac{1}{2} e^{-2x} - x - \frac{1}{2}.$
19. $y = \frac{13}{15} e^{-x} + \frac{1}{12} e^{2x} + \frac{1}{20} \cos 2x - \frac{3}{20} \sin 2x.$
21. $z = A + (Bx^2 + Cx)e^{-x} + D \cos 3x + E \sin 3x.$
23. $z = Ax^2 + Bx + C + Dx \cos x + Ex \sin x.$
25. $z = (Ax^3 + Bx^2)e^{2x} + Cx^2 + Dx + E + (Fx + G) \cos 2x + (Hx + I) \sin 2x.$
27. $z = Ae^{-x} + Bxe^{-x} \cos x + Cxe^{-x} \sin x + D.$
29. $y = C_1 e^{2x} + C_2 x e^{2x} + \frac{8}{25} \cos x + \frac{6}{25} \sin x + 3x e^{2x} \ln x.$
31. $y = C_1 \cos 3x + C_2 \sin 3x + \frac{3}{8} \cos x - \sin 3x \ln(\sec 3x + \tan 3x) + 1.$

33. $y_1 - y_2$ is a solution of the reduced equation $y'' + ay' + by = 0$ with $a, b > 0$. As shown in Exercises 3.4, Problem 37, $y_1 - y_2 \rightarrow 0$ as $x \rightarrow \infty$. If $a = 0, b > 0$, then all solutions of the reduced equation are bounded (Problem 38, Exercises 3.4).

Exercises 3.6

1. The equation of motion is $y(t) = \sin(8t + \frac{1}{2}\pi)$. The amplitude is 1 and the frequency is $8/2\pi = 4/\pi$.
3. The velocity at the equilibrium point is: $\pm 2\pi A/T$.
5. (a) $A \sin(\omega t + \phi_0) = A \cos(\omega t + \phi_0 - \frac{\pi}{2})$; take $\phi_1 = \phi_0 - \frac{1}{2}\pi$.
 (b) $A \sin(\omega t + \phi_0) = A \cos \phi_0 \sin \omega t + A \sin \phi_0 \cos \omega t = B \sin \omega t + C \cos \omega t$.
7. Assume that $r_1 > r_2$. If $C_1 = 0$ or $C_2 = 0$, then $y = C_1 e^{r_1 t} + C_2 e^{r_2 t}$ can never be zero. If both C_1 and C_2 are nonzero, then $C_1 e^{r_1 t} + C_2 e^{r_2 t} = 0$ implies $e^{(r_1 - r_2)t} = -\frac{C_2}{C_1}$. Since $e^{(r_1 - r_2)t}$ is an increasing function ($r_1 > r_2$), it can take the value $-\frac{C_2}{C_1}$ at most once. By the same reasoning, $x'(t) = C_1 r_1 e^{r_1 t} + C_2 r_2 e^{r_2 t}$ can be zero at most once. Therefore the motion can change direction at most once.
9. If $\gamma \neq \omega$, we try $z = A \cos \gamma t + B \sin \gamma t$ as a particular solution of $y'' + \omega^2 y = \frac{F_0}{m} \cos \gamma t$.
 Substituting z into the equation, we get $-\gamma^2 z + \omega^2 z = \frac{F_0}{m} \cos \gamma t$, giving

$$z = \frac{F_0/m}{\omega^2 - \gamma^2} \cos \gamma t.$$

11. If $\gamma = \omega$, we try $z = At \cos \omega t + Bt \sin \omega t$ as a particular solution of

$$y'' + \omega^2 y = \frac{F_0}{m} \cos \omega t.$$

Substituting z into the equation, we have

$$(2B\omega - A\omega^2 t) \cos \omega t - (2A\omega + B\omega^2 t) \sin \omega t + \omega^2 (At \cos \omega t + Bt \sin \omega t) = \frac{F_0}{m} \cos \omega t,$$

which gives $A = 0, B = \frac{F_0}{2\omega m}$, as required.

Exercises 3.7

1. $y = C_1 e^x + C_2 e^{2x} + C_3 e^{3x}$.

3. $y = C_1 e^{2x} + C_2 e^{-2x} + e^x [C_3 \cos 2x + C_4 \sin 2x].$
5. $y = C_1 \cos x + C_2 \sin x + e^{2x} [C_3 \cos 3x + C_4 \sin 3x].$
7. $y = C_1 + C_2 x + C_3 e^x + C_4 e^{-x} + C_5 \cos x + C_6 \sin x.$
9. $y = 2x.$
11. $y^{(4)} - 8y''' + 31y'' - 78y' + 90y = 0.$
13. $y^{(5)} - 2y^{(4)} - 2y''' - 2y'' - 3y' = 0.$
15. $y^{(5)} - 2y^{(4)} + y''' - 2y'' = 0.$
17. $y^{(4)} - y'' = 0.$
19. $y = C_1 e^{-x} + C_2 \cos x + C_3 \sin x + \frac{1}{4}e^x + 4.$
21. $y = C_1 \cos x + C_2 \sin x + C_3 x \cos x + C_4 x \sin x + 6 + \frac{1}{9} \cos 2x.$
23. $y = \frac{1}{72}e^{-x} [e^{3x}(2x - 1) + 3 \cos \sqrt{3}x + \sqrt{3} \sin \sqrt{3}x].$

Chapter 4

Exercises 4.1

1. $F(s) = \frac{1}{s^2}.$
3. $F(s) = \frac{1}{s^2 + 1}.$
5. $F(s) = \frac{1}{s^2 - 1}.$
7. $F(s) = \frac{s - a}{(s - a)^2 + b^2}.$
11. $F(s) = \frac{s}{s^2 + 1}.$
13. $F(s) = \frac{\beta}{s^2 + \beta^2}.$

Exercises 4.2

1. $F(s) = \frac{3}{s} - \frac{2}{s^2} + \frac{2}{s^3}.$
3. $F(s) = \frac{3}{s} + \frac{4}{s-3} - \frac{2s}{s^2+4}.$
5. $F(s) = \frac{10}{s^3} - \frac{4}{(s+3)^2+4}.$
7. $F(s) = \frac{2s}{(s^2+1)^2} + \frac{2(s^2-4)}{(s^2+4)^2}.$
9. $\sinh \beta x = \frac{e^{\beta x} - e^{-\beta x}}{2}.$
11. $F(s) = \frac{1}{2} \left[\frac{1}{s-3} + \frac{1}{s-2} - \frac{1}{s-1} + \frac{1}{s+4} \right].$
15. $Y(s) = \frac{1}{s-2}.$
17. $Y(s) = \frac{2}{(s-2)(s+4)} - \frac{9}{(s^2+9)(s+4)} - \frac{3}{s+4}.$
19. $Y(s) = \frac{2}{(s+3)^2}.$
21. $Y(s) = \frac{3}{s(s-5)(s+3)} + \frac{4}{(s-5)(s+3)^2} + \frac{s-5}{(s-5)(s+3)}.$
23. Set $g(x) = \int_0^x f(t) dt$. Then $g'(x) = f(x)$ and $g(0) = 0$.

$$F(s) = \mathcal{L}[f(x)] = \mathcal{L}[g'(x)] = s\mathcal{L}[g(x)] - g(0) = s\mathcal{L}[g(x)].$$

$$\text{Therefore, } \mathcal{L}[g(x)] = \frac{1}{s}F(s).$$

Exercises 4.3

1. $f(x) = 6e^{-7x}.$
3. $f(x) = \frac{1}{5} \sin 5x.$
5. $f(x) = e^{-4x} \cos x.$
7. $f(x) = e^{-2x} \cos 2x + e^{-2x} \sin 2x.$
9. $f(x) = 2xe^{-2x} - e^x \cos x - e^x \sin x.$

11. $f(x) = \frac{1}{2}e^{-x} - \frac{1}{2}\cos x + \frac{1}{2}\sin x.$
13. $f(x) = \frac{1}{4} - \frac{1}{4}\cos 2x.$
15. $f(x) = \frac{1}{2} - e^x + \frac{3}{2}e^{-2x}.$
17. $f(x) = e^{2x} - 4e^x + 2x + 3.$
19. $y = \frac{2}{3}e^{-2x} + \frac{1}{3}e^x.$
21. $y = \frac{3}{2}e^{-x} - \frac{1}{2}\cos x + \frac{1}{2}\sin x.$
23. $y = e^x \sin x.$
25. $y = \frac{3}{4}e^{-x} + \frac{1}{4}e^x + \frac{1}{2}xe^x.$
27. $y = \frac{1}{4}e^x + xe^{-x} + x - 2.$
29. $y = e^{-2x} + e^x.$
31. $y = -\frac{1}{5}e^{-2x}\cos 2x - \frac{1}{10}e^{-2x}\sin 2x + \frac{1}{5}e^{-x}$
33. $\alpha = \frac{1}{4}.$
35. $\beta = -\frac{26}{5}.$
37. $y = \frac{7}{4}e^{2(x-1)} - 3e^{x-1} + \frac{1}{2}x + \frac{3}{4}.$

Exercises 4.4

1. $F(s) = \frac{2e^{-5s}}{s}.$
3. $F(s) = \frac{2}{s^2} - 2e^{-3s}\frac{1}{s^2} - 5e^{-3s}\frac{1}{s}.$
5. $f(x) = 0 + 5u(x-4); \quad F(s) = 5e^{-4s}\frac{1}{s}.$
7. $f(x) = 0 + (x-2)u(x-2) + 2u(x-2); \quad F(s) = e^{-2s}\frac{1}{s^2} + 2e^{-2s}\frac{1}{s}.$
9. $f(x) = 8 + 2(x-5)u(x-5) + 2u(x-5); \quad F(s) = \frac{8}{s} + 2e^{-5s}\frac{1}{s^2} + 2e^{-5s}\frac{1}{s}.$
11. $f(x) = x^2 - (x-3)^2u(x-3) - 3(x-3)u(x-3); \quad F(s) = \frac{2}{s^3} - e^{-3s}\frac{2}{s^3} - 3e^{-3s}\frac{1}{s^2}.$
13. $f(x) = x - 1 - (x-2)u(x-2) - u(x-2) + e^{-2}e^{-(x-2)}u(x-2);$

$$F(s) = \frac{1}{s^2} - \frac{1}{s} - e^{-2s}\frac{1}{s^2} - e^{-2s}\frac{1}{s} + e^{-2}e^{-2s}\frac{1}{s+1}.$$

$$15. \quad f(x) = \sin 2x - \sin 2(x - \pi)u(x - \pi) + (x - \pi)u(x - \pi) + \pi u(x - \pi);$$

$$F(s) = \frac{2}{s^2 + 4} - e^{-\pi s} \frac{2}{s^2 + 4} + e^{-\pi s} \frac{1}{s^2} + \pi e^{-\pi s} \frac{1}{s}.$$

$$17. \quad f(x) = x - (x - 2)u(x - 2) - 2u(x - 2) + (x - 4)^2 u(x - 4);$$

$$F(s) = \frac{1}{s^2} - e^{-2s} \frac{1}{s^2} - 2e^{-2s} \frac{1}{s} + e^{-4s} \frac{2}{s^3}.$$

$$19. \quad f(x) = 1 - u(x - 2) + (x - 2)u(x - 2) - (x - 4)u(x - 4) - 2u(x - 4) + e^{-(x-4)}u(x - 4);$$

$$F(s) = \frac{1}{s} - e^{-2s} \frac{1}{s} + e^{-2s} \frac{1}{s^2} - e^{-4s} \frac{1}{s^2} - 2e^{-4s} \frac{1}{s} + e^{-4s} \frac{1}{s + 1}.$$

Exercises 4.5

$$1. \quad f(x) = 2 + 2u(x - 3) = \begin{cases} 2, & 0 \leq x < 3 \\ 4, & x \geq 3 \end{cases}.$$

$$3. \quad f(x) = \sin x - \sin x u(x - \pi) = \begin{cases} \sin x, & 0 \leq x < \pi \\ 0, & x \geq \pi. \end{cases}$$

$$5. \quad f(x) = \cos x - \cos x u(x - \pi) + \sin x u(x - \pi) = \begin{cases} \cos x, & 0 \leq x < \pi \\ \sin x, & x \geq \pi. \end{cases}$$

$$7. \quad f(x) = \cos \pi x - \sin \pi x u(x - 2) = \begin{cases} \cos \pi x, & 0 \leq x < 2 \\ \cos \pi x - \sin \pi x & x \geq 2. \end{cases}$$

$$9. \quad f(x) = 3e^{3(x-3)}u(x - 3) - 2e^{2(x-3)}u(x - 3) = \begin{cases} 0, & 0 \leq x < 3 \\ 3e^{3(x-3)} - 2e^{2(x-3)}, & x \geq 3. \end{cases}$$

$$11. \quad f(x) = 2 + e^{(x-1)}u(x - 1) - e^2 e^{(x-2)}u(x - 2) = \begin{cases} 2, & 0 \leq x < 1 \\ 2 + e^{(x-1)}, & 1 \leq x < 2 \\ 2 + e^{(x-1)} - e^x, & x \geq 2. \end{cases}$$

$$13. \quad f(x) = \cos 2x - 1 + u(x - 2) - \cos 2(x - 2)u(x - 2) = \begin{cases} \cos 2x - 1, & 0 \leq x < 2 \\ \cos 2x - \cos 2(x - 2) & x \geq 2. \end{cases}$$

$$15. \quad f(x) = 2e^\pi e^{-2x} \cos 3x u(x - \pi/2) - e^\pi e^{-2x} \sin 3x u(x - \pi/2)$$

$$= \begin{cases} 0, & 0 \leq x < \pi/2 \\ 2e^\pi e^{-2x} \cos 3x - e^\pi e^{-2x} \sin 3x & x \geq \pi/2. \end{cases}$$

Exercises 4.6

1. $y = -\frac{1}{2} + \frac{5}{2}e^{2x} + u(x-1)\left[-\frac{1}{2} + \frac{1}{2}e^{2(x-1)}\right].$

$$= \begin{cases} -\frac{1}{2} + \frac{5}{2}e^{2x}, & 0 \leq x < 1 \\ -1 + \frac{5}{2}e^{2x} + \frac{1}{2}e^{2(x-1)}, & x \geq 1 \end{cases}$$

3. $y = 1 - \cos x + \sin x - u(x-1)[\cos(x-1) - 1].$

$$= \begin{cases} 1 - \cos x + \sin x, & 0 \leq x < \pi \\ 2 \cos x, & x \geq \pi \end{cases}$$

5. $y = 1 - e^{-x} - xe^{-x} + u(x-2)\left[x - 4 + xe^{-(x-2)}\right].$

$$= \begin{cases} 1 - e^{-x} - xe^{-x}, & 0 \leq x < 2 \\ -3 - e^{-x} - xe^{-x} + x + xe^{-(x-2)}, & x \geq 2 \end{cases}$$

7. $y = -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x + u(x-1)\left[\frac{1}{3} + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{x-1}\right].$

$$= \begin{cases} -\frac{1}{3} - \frac{1}{6}e^{3x} + \frac{1}{2}e^x, & 0 \leq x < 1 \\ -\frac{1}{6}e^{3x} + \frac{1}{2}e^x + \frac{1}{6}e^{3(x-1)} - \frac{1}{2}e^{(x-1)}, & x \geq 1 \end{cases}$$

9. $y = \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + u(x-1)\left[xe^{-(x-1)} - 1\right].$

$$= \begin{cases} \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x}, & 0 \leq x < 1 \\ \frac{1}{4}e^x + \frac{11}{4}e^{-x} + \frac{3}{2}xe^{-x} + xe^{-(x-1)} - 1, & x \geq 1 \end{cases}$$

Chapter 5

Exercises 5.2

1. $x = 4, y = 1.$

3. $x = 4 - 2a, y = a, a$ any real number.

5. $x = -3, y = 1.$

7. No solution.

9. $x = 2a - 3, y = a, a$ any real number.

11. $x = \frac{3}{7}a + 1$, $y = \frac{5}{7}a - 1$, $z = a$, a any real number.

13. No solution.

15. $x = 2$, $y = 1$, $z = 1$.

Exercises 5.3

1. matrix of coefficients: 3; augmented matrix: 3; $x = 5$, $y = 3$, $z = -1$.

3. matrix of coefficients: 2; augmented matrix: 2; $x = 4 - 2a$, $y = a$, $z = -2$, a an real number.

5. matrix of coefficients: 3; augmented matrix: 3; $x_1 = -1$, $x_2 = -1 - 2a$, $x_3 = 3 + a$, $x_4 = a$, a any real number.

7. matrix of coefficients: 3; augmented matrix: 3; $x_1 = 8 + 2a - 3b$, $x_2 = a$, $x_3 = 3 - 1 - 2b$, $x_4 = b$, $x_5 = -3$, a , b any real numbers.

9. $x = 2$, $y = 5$.

11. $x = -3 - a$, $y = 2 + 2a$, $z = a$, a any real number.

13. $x = \frac{10}{7}$, $y = \frac{2}{7}$, $z = \frac{3}{2}$.

15. $x_1 = 11 - 2a + b$, $x_2 = a$, $x_3 = 3 - b$, $x_4 = b$, a , b any real numbers.

17. $x_1 = -2$, $x_2 = -5$, $x_3 = -1$, $x_4 = 5$.

19. $x_1 = 3 - 2a$, $x_2 = a$, $x_3 = 2$, $x_4 = 1$.

21. (i) $k \neq -3$, 2 (ii) $k = -3$ (iii) $k = 2$.

23. (i) $a \neq -3$, 3 (ii) $a = -3$ (iii) $a = 3$.

25. The system has at least one solution if $b = 2a$, a any real number.

27. (a) No (b) No (c) Yes

29. $y = \frac{4}{3}e^x - \frac{1}{3}e^{-2x}$.

31. $y = -\frac{7}{2} + \frac{1}{2}e^{2x} + 4e^{-x/2}$.

Exercises 5.4

1. Yes

3. No

5. No. The leading 1 in the last column is not the only nonzero in its column.
7. Yes
9. $x = 10, y = -9, z = -7$
11. $x = -3 - a, y = 2 + 2a, z = a, a$ any real number.
13. $x_1 = 11 - 2a + b, x_2 = a, x_3 = 3 - b, x_4 = b, a, b$ any real numbers.
15. $x_1 = 7 - 2a - b, x_2 = 1 + 3a - 4b, x_3 = a, x_4 = b, a, b$ any real numbers.
17. $x = y = 0$.
19. $x = y = z = 0$.
21. $x_1 = 2a - b, x_2 = -a + 4b, x_3 = a, x_4 = b, a, b$ any real numbers.
23. $x_1 = x_2 = x_3 = x_4 = 0$.
25. Consider the system

$$\begin{aligned}x + y &= 0 \\2x + 2y &= 0 \\3x + 3y &= 0\end{aligned}$$

This system has the solutions $x = -a, y = a, a$ any real number.

27. $a = 4$.
29. (a) $x = 1 + a, y = -1 - a, z = a, a$ any real number.
- (b) $(x, y, z) = (1 + a, -1 - a, a) = (1, -1, 0) + (a, -a, a), a$ any real number;
 $x = 1, y = -1, z = 0$ is a solution of the nonhomogeneous system, $x = a, y = -a, z = a$ is the set of all solutions of the corresponding homogeneous system.

Exercises 5.5

1. (a) $\begin{pmatrix} 0 & 4 \\ 3 & 5 \\ 1 & -1 \end{pmatrix}.$
- (c) $\begin{pmatrix} 8 & -4 \\ 1 & 4 \end{pmatrix}.$

$$(e) \begin{pmatrix} 2 & 8 \\ 2 & 8 \\ 5 & -3 \end{pmatrix}.$$

$$3. (a) \begin{pmatrix} -4 & -3 \\ 28 & -6 \\ -20 & 24 \end{pmatrix}.$$

(c) Not defined.

$$(e) \begin{pmatrix} 1 & 3 \\ -3 & -12 \\ -41 & 21 \end{pmatrix}.$$

$$5. (a) c_{32} = 2 \quad (b) c_{13} = 34 \quad (c) d_{21} = 5 \quad (d) d_{22} = 1.$$

$$7. (a) d_{22} = 6 \quad (b) d_{12} = -4 \quad (c) d_{23} = -18.$$

$$11. (a) AB = \begin{pmatrix} 4 & 7 & 10 \\ 0 & -5 & -14 \end{pmatrix}, \quad BA \text{ not defined.}$$

$$(b) AC = \begin{pmatrix} 14 & 5 \\ -2 & -3 \end{pmatrix}, \quad CA = \begin{pmatrix} -1 & 14 \\ 5 & 12 \end{pmatrix}.$$

$$(c) AD = DA = \begin{pmatrix} 4 & 4 \\ -2 & 2 \end{pmatrix}.$$

$$13. A(BD) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}$$

$$(AB)D = \begin{pmatrix} 0 & -2 \\ 3 & 5 \end{pmatrix} \begin{pmatrix} -2 & 3 & -2 \\ 3 & 0 & -4 \end{pmatrix} = \begin{pmatrix} -6 & 0 & 8 \\ 9 & 9 & -26 \end{pmatrix}.$$

$$15. (a) 3 \times 3 \quad (c) \text{ Does not exist} \quad (e) 2 \times 3.$$

Exercises 5.6

$$1. A^{-1} = \begin{pmatrix} 1/2 & 0 \\ -3/2 & 1 \end{pmatrix}.$$

$$3. A^{-1} = \begin{pmatrix} -2 & 1 \\ 3/2 & -1/2 \end{pmatrix}.$$

$$5. A^{-1} = \begin{pmatrix} -11 & 2 & 2 \\ -4 & 0 & 1 \\ 6 & -1 & -1 \end{pmatrix}.$$

- 7. No inverse.
- 9. No inverse.
- 11. $\det A = \pm 1$.
- 13. $x = 5, y = 0$.
- 15. $x = \frac{9}{2}, y = -5$.
- 17. $x = \frac{7}{9}, y = \frac{1}{3}, z = -\frac{5}{9}$.
- 19. -31 .
- 21. -45 .
- 23. 30 .
- 25. -21 .
- 27. -18 .
- 29. 26 .
- 31. $x = 0, 1, -3$.
- 33. $y = -\frac{25}{37}$.
- 35. Cramer's rule does not apply.
- 37. $x = 0$.
- 39. $\lambda = -4, 7$.

Exercises 5.7

- 3. Dependent; $(-4, 8, 9) = 2(1, -2, 3) + 3(-2, 4, 1)$.
- 5. Dependent; $(-2, 6, 3) = (1, -1, 3) + 2(0, 2, 3) - 3(1, -1, 2)$.
- 7. Dependent; $(7, -4, 1) = 3(1, -2, 1) + 2(2, 1, -1)$.
- 9. Dependent; $(4, -2, 0, 2) = 2(2, -1, 0, 1)$.
- 11. $b \neq -\frac{1}{3}$.
- 13. $b = 0, -7$.
- 17. No; a linearly dependent set can have linearly independent subsets. For example,

$\{(1, -2, 3), (-2, 4, 1)\}$ is a linearly independent subset of $\{(1, -2, 3), (-2, 4, 1)\}, (-4, 8, 9)$.

19. $W(x) = -a$; linearly independent.

21. $W(x) = -2x^{-6}$; linearly independent.

23. $W(x) = e^{2x}(x - 2)$; linearly independent.

25. (a) False (b) True (c) True.

Exercises 5.8

1. $2, \begin{pmatrix} 1 \\ 0 \end{pmatrix}; \quad 3, \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$

3. $-1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}; \quad 4, \begin{pmatrix} 2 \\ 3 \end{pmatrix}.$

5. $1, 1, \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$

7. $2, 2, \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$

9. $2 + i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} + i \begin{pmatrix} 0 \\ -1 \end{pmatrix}; \quad 2 - i, \begin{pmatrix} 1 \\ 0 \end{pmatrix} - i \begin{pmatrix} 0 \\ -1 \end{pmatrix}.$

11. $8, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}; \quad 1, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}; \quad 2, \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.$

13. $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}.$

15. $1 + i, \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} + i \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}; \quad 1 - i, \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} - i \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}; \quad 0, \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}.$

17. $1, 1, 1, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$

$$19. \quad 2 + 3i, \begin{pmatrix} -5 \\ 3 \\ 2 \end{pmatrix} + i \begin{pmatrix} 3 \\ 3 \\ 0 \end{pmatrix}; \quad 2 - 3i, \begin{pmatrix} -5 \\ 3 \\ 2 \end{pmatrix} - i \begin{pmatrix} 3 \\ 3 \\ 0 \end{pmatrix}; \quad 2, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}.$$

$$21. \quad 2, \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}; \quad 2, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}; \quad 6, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}; \quad 4, \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}.$$

Chapter 6

Exercises 6.1

$$1. \quad \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -3 & t \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 0 \\ \sin 2t \end{pmatrix}.$$

$$3. \quad \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ e^t \end{pmatrix}.$$

$$5. \quad \begin{aligned} x'_1 &= 2x_1 - x_2 + e^{2t} \\ x'_2 &= 3x_1 + 2e^{-t} \end{aligned}.$$

$$7. \quad \begin{aligned} x'_1 &= 2x_1 + 3x_2 - x_3 + e^t \\ x'_2 &= -2x_1 + x_3 + 2e^{-t} \\ x'_3 &= 2x_1 + 3x_2 + e^{2t} \end{aligned}.$$

$$9. \quad \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} \sin t \\ -2 \cos t \end{pmatrix}.$$

$$11. \quad \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 2 & 1 & 3 \\ 1 & -3 & 0 \\ 2 & -1 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 3e^{2t} \\ -2 \cos t \\ t \end{pmatrix}.$$

Exercises 6.2

1. Independent

3. Independent

5. Dependent

7. Dependent

9. Dependent

11. (c) $\mathbf{x}(t) = c_1\mathbf{u} + c_2\mathbf{v}$, where c_1, c_2 are arbitrary constants.

(d) $\mathbf{x}(t) = -2 \begin{pmatrix} e^{2t} \\ e^{2t} \end{pmatrix} + \begin{pmatrix} 3e^{3t} \\ 2e^{3t} \end{pmatrix}.$

13. (b) $\mathbf{x}(t) = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix} - \begin{pmatrix} 4te^{-t} \\ e^{-t} \\ 0 \end{pmatrix}.$

Exercises 6.3

1. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-3t} \begin{pmatrix} 4 \\ -1 \end{pmatrix}.$

3. $\mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + C_2 e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}; \quad \mathbf{x}(t) = -3e^{3t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + 4e^{2t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$

5. $\mathbf{x}(t) = C_1 e^{5t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 2 \\ -1 \end{pmatrix}.$

7. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}.$

9. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + C_3 e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix};$

$$\mathbf{x}(t) = -e^{-t} \begin{pmatrix} 3 \\ -2 \\ 12 \end{pmatrix} + 2e^t \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}.$$

11. $\mathbf{x}(t) = C_1 e^{10t} \begin{pmatrix} 9 \\ 7 \\ 4 \end{pmatrix} + C_2 e^{6t} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + C_3 e^{5t} \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}.$

13. (a) $\mathbf{x}' = \begin{pmatrix} 0 & 1 \\ -b & -a \end{pmatrix} \mathbf{x};$ (b) $\lambda^2 + a\lambda + b = 0;$ (c) The are the same.

Exercises 6.4

1. $\mathbf{x}(t) = C_1 \left[\cos 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right] + C_2 \left[\cos 2t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ -1 \end{pmatrix} \right].$

3. $\mathbf{x}(t) = C_1 e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + C_2 \left[e^t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right].$

5. $\mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} + C_2 \left[e^{-t} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + t e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} \right].$

7. $\mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + C_2 e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + C_3 e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix};$

$$\mathbf{x}(t) = e^{-t} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + e^{-t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$$

9. $\mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} + C_2 \left[\cos 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} - \sin 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] + C_3 \left[\cos 2t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \sin 2t \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} \right].$

11. $\mathbf{x}(t) = C_1 e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + C_3 e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix};$

$$\mathbf{x}(t) = \frac{7}{2} e^{3t} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \frac{9}{2} e^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \frac{3}{2} e^t \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}$$

13. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + C_3 \left[e^t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + t e^t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$

15. $\mathbf{x}(t) = C_1 e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + C_2 e^t \left[\cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + C_3 e^t \left[\cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right];$

$$\mathbf{x}(t) = e^{2t} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + e^t \left[\cos t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \sin t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right] + 3e^t \left[\cos t \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} + \sin t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right].$$

$$17. \quad \mathbf{x}(t) = C_1 e^{6t} \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} + C_2 e^{-2t} \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} + C_3 e^{-2t} \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix}.$$

Exercises 6.5

$$1. \quad \mathbf{x}(t) = \begin{pmatrix} e^{3t} & e^{-t} \\ e^{3t} & 5e^{-t} \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \begin{pmatrix} \frac{1}{4}te^{-t} \\ \frac{5}{4}te^{-t} \end{pmatrix}.$$

$$3. \quad \mathbf{x}(t) = \begin{pmatrix} -2e^{-t} & -1 \\ 3e^{-t} & 1 \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \begin{pmatrix} 2t^2 - t + 2 \\ -2t^2 + 3t - 3 \end{pmatrix}.$$

$$5. \quad \mathbf{x}(t) = \begin{pmatrix} -e^{-5t} & e^{-2t} \\ 2e^{-5t} & e^{-2t} \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \begin{pmatrix} -\frac{27}{50} + \frac{1}{4}e^{-t} + \frac{6}{5}t \\ -\frac{21}{50} + \frac{1}{2}e^{-t} + \frac{3}{5}t \end{pmatrix}.$$

$$7. \quad \mathbf{x}(t) = \begin{pmatrix} -e^t \sin t & e^t \cos t \\ e^t \cos t & e^t \sin t \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \begin{pmatrix} te^t \cos t \\ te^t \sin t \end{pmatrix}.$$

$$9. \quad \mathbf{x}(t) = \begin{pmatrix} 0 & e^{2t} & -1 \\ 0 & e^{2t} & 1 \\ e^{3t} & 0 & 0 \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \\ C_3 \end{pmatrix} + \begin{pmatrix} \frac{1}{2}te^{2t} - \frac{1}{4}e^{2t} \\ -e^t + \frac{1}{4}e^{2t} + \frac{1}{2}te^{2t} \\ \frac{1}{2}t^2e^{3t} \end{pmatrix}.$$

$$11. \quad \mathbf{x}(t) = \begin{pmatrix} e^{4t} \\ -e^{4t} \end{pmatrix} + \begin{pmatrix} -e^{2t} [1 - 2t - e^{2t} + 2te^{2t}] \\ e^{2t} [1 + 2t + e^{2t} + 2te^{2t}] \end{pmatrix}.$$