## Math 1313 Section 3.5 Section 3.5: The Inverse of a Matrix

Over the set of real number we have what we call the **multiplicative inverse** or **reciprocal**. The multiplicative inverse of a number is a second number that when multiplied by the first number yields the **multiplicative identity 1**.

This is where the Identity Matrix comes in.

Let A be a square matrix of size n and another square matrix  $A^{-1}$  of size n such that  $AA^{-1} = A^{-1}A = I_n$  is called the **inverse of A**.

Note: Not every square matrix has an inverse. A matrix with no inverse is called **singular**.

## Finding the Inverse of a Matrix

Given the n x n matrix *A*:

- 1. Adjoin the n x n identity matrix I to obtain the augmented matrix (A | I)
- 2. Use the Gauss-Jordan elimination method to reduce (A | I) to the form (I | B), if possible.

The matrix B is the inverse of A.

**Example 1:** Find the inverse, if possible and check:

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

Math 1313Section 3.5**Example 2:** Find the inverse of a 3 x 3 matrix.(Use Gauss-Jordan)

$$\mathbf{C} = \begin{pmatrix} 1 & 4 & -1 \\ 2 & 3 & -2 \\ -1 & 2 & 3 \end{pmatrix}$$

Math 1313 Section 3.5 **Example 3:** Find the inverse.

$$\mathbf{B} = \begin{pmatrix} 4 & 2 & 2 \\ -1 & -3 & 4 \\ 3 & -1 & 6 \end{pmatrix}$$

## **Matrices That Have No Inverses**

If there is a row to the left of the vertical line in the augmented matrix containing all zeros, then the matrix does not have an inverse. Example 3 has this problem and does not have an inverse.

Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . Suppose D = ad - bc is not equal to zero. Then  $A^{-1}$  exists and is given by  $A^{-1} = \frac{1}{D} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ 

**Example 4:** Find the inverse of the following matrices.

**a.**  $A = \begin{pmatrix} -5 & 10 \\ 2 & 7 \end{pmatrix}$ 

**b.** 
$$B = \begin{pmatrix} 8 & -4 \\ -4 & 2 \end{pmatrix}$$

## **Matrix Representation**

A system of linear equations may be written in a compact form with the help of matrices.

**Example 5:** Given the following system of equations, write it in matrix form.

2x-4y+z=6-3x+6y-5z=-1x-3y+7z=0

Math 1313 Section 3.5

**Example 6:** Write each system of equations as a matrix equation and then solve the system using the inverse of the coefficient matrix.

2x + 3y = 53x + 5y = 8

The use of inverses to solve systems of equations is advantageous when we are required to solve more than one system of equations AX = B, involving the same coefficient matrix, A, and different matrices of constants, B.

**Example 7:** A performance theatre has 10,000 seats. The ticket prices are either \$25 or \$35, depending on the location of the seat. Assume every seat can be sold.

a. How many tickets of each type should be sold to bring in a return of \$275,000? b. How many tickets of each type should be sold to bring in a return of \$300,000? Let x = number of \$25 tickets and y = number of \$35 tickets