

**The 60th William Lowell Putnam Mathematical Competition**  
**Saturday, December 4, 1999**

A-1 Find polynomials  $f(x), g(x)$ , and  $h(x)$ , if they exist, such that for all  $x$ ,

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

A-2 Let  $p(x)$  be a polynomial that is nonnegative for all real  $x$ . Prove that for some  $k$ , there are polynomials  $f_1(x), \dots, f_k(x)$  such that

$$p(x) = \sum_{j=1}^k (f_j(x))^2.$$

A-3 Consider the power series expansion

$$\frac{1}{1 - 2x - x^2} = \sum_{n=0}^{\infty} a_n x^n.$$

Prove that, for each integer  $n \geq 0$ , there is an integer  $m$  such that

$$a_n^2 + a_{n+1}^2 = a_m.$$

A-4 Sum the series

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{m^2 n}{3^m (n3^m + m3^n)}.$$

A-5 Prove that there is a constant  $C$  such that, if  $p(x)$  is a polynomial of degree 1999, then

$$|p(0)| \leq C \int_{-1}^1 |p(x)| dx.$$

A-6 The sequence  $(a_n)_{n \geq 1}$  is defined by  $a_1 = 1, a_2 = 2, a_3 = 24$ , and, for  $n \geq 4$ ,

$$a_n = \frac{6a_{n-1}^2 a_{n-3} - 8a_{n-1} a_{n-2}^2}{a_{n-2} a_{n-3}}.$$

Show that, for all  $n$ ,  $a_n$  is an integer multiple of  $n$ .

B-1 Right triangle  $ABC$  has right angle at  $C$  and  $\angle BAC = \theta$ ; the point  $D$  is chosen on  $AB$  so that  $|AC| = |AD| = 1$ ; the point  $E$  is chosen on  $BC$  so that  $\angle CDE = \theta$ . The perpendicular to  $BC$  at  $E$  meets  $AB$  at  $F$ . Evaluate  $\lim_{\theta \rightarrow 0} |EF|$ .

B-2 Let  $P(x)$  be a polynomial of degree  $n$  such that  $P(x) = Q(x)P''(x)$ , where  $Q(x)$  is a quadratic polynomial and  $P''(x)$  is the second derivative of  $P(x)$ . Show that if  $P(x)$  has at least two distinct roots then it must have  $n$  distinct roots.

B-3 Let  $A = \{(x, y) : 0 \leq x, y < 1\}$ . For  $(x, y) \in A$ , let

$$S(x, y) = \sum_{\frac{1}{2} \leq \frac{m}{n} \leq 2} x^m y^n,$$

where the sum ranges over all pairs  $(m, n)$  of positive integers satisfying the indicated inequalities. Evaluate

$$\lim_{(x,y) \rightarrow (1,1), (x,y) \in A} (1 - xy^2)(1 - x^2y)S(x, y).$$

B-4 Let  $f$  be a real function with a continuous third derivative such that  $f(x), f'(x), f''(x), f'''(x)$  are positive for all  $x$ . Suppose that  $f'''(x) \leq f(x)$  for all  $x$ . Show that  $f'(x) < 2f(x)$  for all  $x$ .

B-5 For an integer  $n \geq 3$ , let  $\theta = 2\pi/n$ . Evaluate the determinant of the  $n \times n$  matrix  $I + A$ , where  $I$  is the  $n \times n$  identity matrix and  $A = (a_{jk})$  has entries  $a_{jk} = \cos(j\theta + k\theta)$  for all  $j, k$ .

B-6 Let  $S$  be a finite set of integers, each greater than 1. Suppose that for each integer  $n$  there is some  $s \in S$  such that  $\gcd(s, n) = 1$  or  $\gcd(s, n) = s$ . Show that there exist  $s, t \in S$  such that  $\gcd(s, t)$  is prime.