- B2 ('23) For each positive integer n, let k(n) be the number of B2 ('16) Define a positive integer n to be *squarish* if either n ones in the binary representation of $2023 \cdot n$. What is the minimum value of k(n)?
- B2 ('22) Let \times represent the cross product in \mathbb{R}^3 . For what positive integers n does there exist a set $S \subset \mathbb{R}^3$ with exactly n elements such that

$$S = \{v \times w : v, w \in S\}?$$

B2 ('21) Determine the maximum value of the sum

$$S = \sum_{n=1}^{\infty} \frac{n}{2^n} (a_1 a_2 \cdots a_n)^{1/n}$$

over all sequences a_1, a_2, a_3, \cdots of nonnegative real numbers satisfying

$$\sum_{k=1}^{\infty} a_k = 1.$$

B2 ('20) Let k and n be integers with $1 \le k < n$. Alice and Bob play a game with k pegs in a line of n holes. At the beginning of the game, the pegs occupy the k leftmost holes. A legal move consists of moving a single peg to any vacant hole that is further to the right. The players alternate moves, with Alice playing first. The game ends when the pegs are in the k rightmost holes, so whoever is next to play cannot move and therefore loses. For what values of n and k does Alice have a winning strategy?

B2 For all $n \ge 1$, let

$$a_n = \sum_{k=1}^{n-1} \frac{\sin\left(\frac{(2k-1)\pi}{2n}\right)}{\cos^2\left(\frac{(k-1)\pi}{2n}\right)\cos^2\left(\frac{k\pi}{2n}\right)}$$

Determine

$$\lim_{n \to \infty} \frac{a_n}{n}$$

- B2 ('18) Let n be a positive integer, and let $f_n(z) = n + (n 1)z + (n 2)z^2 + \cdots + z^{n-1}$. Prove that f_n has no roots in the closed unit disk $\{z \in \mathbb{C} : |z| \le 1\}$.
- B2 ('17) Suppose that a positive integer N can be expressed as the sum of k consecutive positive integers

$$N = a + (a + 1) + (a + 2) + \dots + (a + k - 1)$$

for k = 2017 but for no other values of k > 1. Considering all positive integers N with this property, what is the smallest positive integer a that occurs in any of these expressions?

16) Define a positive integer n to be *squarish* if either n is itself a perfect square or the distance from n to the nearest perfect square is a perfect square. For example, 2016 is squarish, because the nearest perfect square to 2016 is $45^2 = 2025$ and 2025 - 2016 = 9 is a perfect square. (Of the positive integers between 1 and 10, only 6 and 7 are not squarish.)

For a positive integer N, let S(N) be the number of squarish integers between 1 and N, inclusive. Find positive constants α and β such that

$$\lim_{N \to \infty} \frac{S(N)}{N^{\alpha}} = \beta,$$

or show that no such constants exist.

- B2 ('15) Given a list of the positive integers 1, 2, 3, 4, ..., take the first three numbers 1, 2, 3 and their sum 6 and cross all four numbers off the list. Repeat with the three smallest remaining numbers 4, 5, 7 and their sum 16. Continue in this way, crossing off the three smallest remaining numbers and their sum, and consider the sequence of sums produced: 6, 16, 27, 36, Prove or disprove that there is some number in the sequence whose base 10 representation ends with 2015.
- B2 ('14) Suppose that f is a function on the interval [1,3] such that $-1 \le f(x) \le 1$ for all x and $\int_1^3 f(x) dx = 0$. How large can $\int_1^3 \frac{f(x)}{x} dx$ be?
- B2 ('13) Let $C = \bigcup_{N=1}^{\infty} C_N$, where C_N denotes the set of 'cosine polynomials' of the form $f(x) = 1 + \sum_{n=1}^{N} a_n \cos(2\pi nx)$ for which:
 - (i) $f(x) \ge 0$ for all real x, and

(ii) $a_n = 0$ whenever *n* is a multiple of 3.

Determine the maximum value of f(0) as f ranges through C, and prove that this maximum is attained.

- B2 ('12) Let P be a given (non-degenerate) polyhedron. Prove that there is a constant c(P) > 0 with the following property: If a collection of n balls whose volumes sum to V contains the entire surface of P, then $n > c(P)/V^2$.
- B2 ('11) Let S be the set of all ordered triples (p, q, r) of prime numbers for which at least one rational number x satisfies $px^2 + qx + r = 0$. Which primes appear in seven or more elements of S?
- B2 ('10) Given that A, B, and C are noncollinear points in the plane with integer coordinates such that the distances AB, AC, and BC are integers, what is the smallest possible value of AB?

B2 ('85) Define polynomials $f_n(x)$ for $n \ge 0$ by $f_0(x) = 1$, B2 ('93) Consider the following game played with a deck of 2n $f_n(0) = 0$ for $n \ge 1$, and

$$\frac{d}{dx}f_{n+1}(x) = (n+1)f_n(x+1)$$

for n > 0. Find, with proof, the explicit factorization of $f_{100}(1)$ into powers of distinct primes.

B2 ('86) Prove that there are only a finite number of possibilities for the ordered triple T = (x - y, y - z, z - x), where x, y, z are complex numbers satisfying the simultane- B2 ('94) For which real numbers c is there a straight line that ous equations

$$x(x-1) + 2yz = y(y-1) + 2zx + z(z-1) + 2xy,$$

and list all such triples T.

B2 ('87) Let r,s and t be integers with $0\leq r,0\leq s$ and $r+s\leq$ t. Prove that

$$\frac{\binom{s}{0}}{\binom{t}{r}} + \frac{\binom{s}{1}}{\binom{t}{r+1}} + \dots + \frac{\binom{s}{s}}{\binom{t}{r+s}} = \frac{t+1}{(t+1-s)\binom{t-s}{r}}$$

- B2 ('88) Prove or disprove: If x and y are real numbers with B2 ('96) Show that for every positive integer n, $y \ge 0$ and $y(y+1) \le (x+1)^2$, then $y(y-1) \le x^2$.
- B2 ('89) Let S be a non-empty set with an associative operation that is left and right cancellative (xy = xz implies y =z, and yx = zx implies y = z). Assume that for every be a group?
- B2 ('90) Prove that for |x| < 1, |z| > 1,

$$1 + \sum_{j=1}^{\infty} (1 + x^j) P_j = 0,$$

where P_i is

$$\frac{(1-z)(1-zx)(1-zx^2)\cdots(1-zx^{j-1})}{(z-x)(z-x^2)(z-x^3)\cdots(z-x^j)}.$$

B2 ('91) Suppose f and g are non-constant, differentiable, real- B2 ('99) Let P(x) be a polynomial of degree n such that P(x) =valued functions defined on $(-\infty, \infty)$. Furthermore, suppose that for each pair of real numbers x and y,

$$f(x+y) = f(x)f(y) - g(x)g(y), g(x+y) = f(x)g(y) + g(x)f(y).$$

- If f'(0) = 0, prove that $(f(x))^2 + (g(x))^2 = 1$ for all B2 ('00) Prove that the expression x.
- B2 ('92) For nonnegative integers n and k, define Q(n, k) to be the coefficient of x^k in the expansion of $(1 + x + x^2 +$ $(x^3)^n$. Prove that

$$Q(n,k) = \sum_{j=0}^{k} \binom{n}{j} \binom{n}{k-2j},$$

where $\binom{a}{b}$ is the standard binomial coefficient. (Reminder: For integers a and b with $a \ge 0$, $\binom{a}{b} = \frac{a!}{b!(a-b)!}$ for $0 \le b \le a$, with $\binom{a}{b} = 0$ otherwise.)

- cards numbered from 1 to 2n. The deck is randomly shuffled and n cards are dealt to each of two players. Beginning with A, the players take turns discarding one of their remaining cards and announcing its number. The game ends as soon as the sum of the numbers on the discarded cards is divisible by 2n + 1. The last person to discard wins the game. Assuming optimal strategy by both A and B, what is the probability that A wins?
- intersects the curve

$$x^4 + 9x^3 + cx^2 + 9x + 4$$

in four distinct points?

- B2 ('95) An ellipse, whose semi-axes have lengths a and b, rolls without slipping on the curve $y = c \sin\left(\frac{x}{a}\right)$. How are a, b, c related, given that the ellipse completes one revolution when it traverses one period of the curve?

$$\left(\frac{2n-1}{e}\right)^{\frac{2n-1}{2}} < 1 \cdot 3 \cdot 5 \cdots (2n-1) < \left(\frac{2n+1}{e}\right)^{\frac{2n+1}{2}}.$$

a in S the set $\{a^n : n = 1, 2, 3, ...\}$ is finite. Must S B2 ('97) Let f be a twice-differentiable real-valued function satisfying

$$f(x) + f''(x) = -xg(x)f'(x),$$

where $g(x) \ge 0$ for all real x. Prove that |f(x)| is bounded.

- B2 ('98) Given a point (a, b) with 0 < b < a, determine the minimum perimeter of a triangle with one vertex at (a, b), one on the x-axis, and one on the line y = x. You may assume that a triangle of minimum perimeter exists.
 - Q(x)P''(x), where Q(x) is a quadratic polynomial and P''(x) is the second derivative of P(x). Show that if P(x) has at least two distinct roots then it must have ndistinct roots.

$$\frac{gcd(m,n)}{n}\binom{n}{m}$$

is an integer for all pairs of integers $n \ge m \ge 1$.

B2 ('01) Find all pairs of real numbers (x, y) satisfying the system of equations

$$\frac{1}{x} + \frac{1}{2y} = (x^2 + 3y^2)(3x^2 + y^2)$$
$$\frac{1}{x} - \frac{1}{2y} = 2(y^4 - x^4).$$

B2 ('02) Consider a polyhedron with at least five faces such that B2 ('06) Prove that, for every set $X = \{x_1, x_2, \dots, x_n\}$ of n exactly three edges emerge from each of its vertices. Two players play the following game:

> Each player, in turn, signs his or her name on a previously unsigned face. The winner is the player who first succeeds in signing three faces that share a common vertex.

Show that the player who signs first will always win by playing as well as possible.

- B2 ('03) Let n be a positive integer. Starting with the sequence $1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}$, form a new sequence of n-1 entries $\frac{3}{4}, \frac{5}{12}, \dots, \frac{2n-1}{2n(n-1)}$ by taking the averages of two consecutive entries in the first sequence. Repeat the averaging of neighbors on the second sequence to obtain a third sequence of n-2 entries, and continue until the final sequence produced consists of a single number x_n . Show that $x_n < 2/n$.
- B2 ('04) Let m and n be positive integers. Show that

$$\frac{(m+n)!}{(m+n)^{m+n}} < \frac{m!}{m^m} \frac{n!}{n^n}.$$

B2 ('05) Find all positive integers n, k_1, \ldots, k_n such that $k_1 +$ $\cdots + k_n = 5n - 4$ and

$$\frac{1}{k_1} + \dots + \frac{1}{k_n} = 1.$$

real numbers, there exists a non-empty subset S of Xand an integer m such that

$$\left|m + \sum_{s \in S} s\right| \le \frac{1}{n+1}$$

B2 ('07) Suppose that $f : [0,1] \to \mathbb{R}$ has a continuous derivative and that $\int_0^1 f(x) dx = 0$. Prove that for every $\alpha \in$ (0, 1),

$$\left| \int_{0}^{\alpha} f(x) \, dx \right| \le \frac{1}{8} \max_{0 \le x \le 1} |f'(x)|.$$

B2 ('08) Let $F_0(x) = \ln x$. For $n \ge 0$ and x > 0, let $F_{n+1}(x) =$ $\int_0^x F_n(t) dt$. Evaluate

$$\lim_{n \to \infty} \frac{n! F_n(1)}{\ln n}.$$

B2 ('09) A game involves jumping to the right on the real number line. If a and b are real numbers and b > a, the cost of jumping from a to b is $b^3 - ab^2$. For what real numbers c can one travel from 0 to 1 in a finite number of jumps with total cost exactly c?